## Brownian Motion of Grains and Negative Friction in Dusty Plasmas S.A. Trigger

Joint Institute for High Temperatures, Russian Academy of Sciences, 13/19 Izhorskaya Str., Moscow 127412, Russia; Humboldt University, 110 Invalidenstr., D-10115 Berlin, Germany

Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine

14B Metrolohichna Str., Kiev 03143, Ukraine

e-mail: azagorodny@gluk.org

Within the approximation of dominant charging collisions the explicit microscopic calculations of the Fokker-Planck kinetic coefficients for highly-charged grains moving in plasma are performed. It is shown that due to ion absorption by grain the friction coefficient can be negative and the appropriate threshold value of the grain charge is found. The stationary solutions of the Fokker-Planck equation with the velocity dependent kinetic coefficient are obtained and the considerable deviation of such solutions from the Maxwellian distribution is established.

PACS number(s) : 52.27.Lw, 52.20.Hv, 52.25.Fi

In the systems close to equilibrium Brownian particles keep stationary random motion under the action of stochastic forces which are compensated by the particle friction and thus, the work produced by the Langevin sources is equal to the energy dissipated in course of the Brownian particle motion. This energy balance is described by the fluctuation-dissipation theorem in the form of the Einstein law. Obviously, the fluctuationdissipation theorem and the Einstein relation can be violated in the case of non-equilibrium systems (even in the stationary case), in particular in the open systems. Starting from the classical Lord Rayleigh work [1] many studies of the non-equilibrium motion of Brownian particles with additional (inner, or external) energy supply have been performed. In particular, such studies are of great importance for physico-chemical [2, 3] and biological [4] systems in which non-equilibrium Brownian particle motion is referred as the motion of active Brownian particles. Recently the dynamical and energetic aspects of motion for the active Brownian particles have been described on the basis of the Langevin equation and the appropriate Fokker-Planck equation [5, 6]. The possibility of negative friction (negative values of friction coefficient) of the Brownian particles was regarded, as a result of energy pumping. For some phenomenological dependences of the friction coefficient as a function of the grain's velocity the one-particle stationary non-Maxwellian distribution function was found.

The traditional formulations of the non-equilibrium Brownian motion are based on some phenomenological expressions for the friction and diffusion coefficients. In particular, it means that deviations from the Einstein relation, as well as the velocity dependence of these coefficients are postulated and high level of uncertainty for the application of such models to the real systems takes place. Here we will consider another situation, when the kinetic coefficients can be calculated exactly on the basis of the microscopically derived Fokker-Planck equation for dusty plasmas [7,8]. It will be shown that in the case of strong interaction parameter  $\Gamma \equiv e^2 Z_a Z_i / a T_i \gg 1$ (here  $Z_q$ ,  $Z_i$  are the charge numbers for the grains and ions respectively, a is the grain radius,  $T_i$  is the ion temperature) the negative friction coefficient appears for some velocity domain. If the charging collisions are dominant, this domain is determined by the inequalities:  $v^2 < 10v_{\text{Ti}}^2(\Gamma - 1)/(3\Gamma - 1)$  for  $\Gamma \ge 1$ , but  $(\Gamma - 1) \ll 1, (v_{\text{Ti}}^2 \equiv T_i/m_i)$ ; and  $v^2 < 2v_{\text{Ti}}^2\Gamma$  for  $\Gamma \gg 1$ . The physical reason for manifestation of negative friction is clear: the cross-section for ion absorption by grain increases with the relative velocity between the ion and grain decrease due to the charge-dependent part of the cross-section. Therefore, for a moving highly-charged grain ( $\Gamma \gg 1$ ) the momentum transfer from ions to the grain in the direction of grain velocity is higher than in the opposite direction.

Naturally, the Coulomb scattering and particle friction related to the ion-grain and neutral-grain elastic collisions will increase the threshold for negative friction and even can suppress it for some plasma parameters. However, the latter are rather sophisticated processes, and their theoretical description within various approximations can shadow the physics of the phenomenon. Some estimate of the influence of such processes will be also done below, but the detailed description of their influence should be a matter of further consideration. On the other hand, the problem of microscopic investigation of negative friction in dusty plasmas is so fundamental, that it deserves a description of its simplest manifestation (which occurs in the case of dominant charging collisions) to be considered in this Letter.

We start from the Fokker-Planck equation for the spherical grains in dusty plasma with typical narrow charge distribution around a negative value  $q = e_e Z_g$ , that permits to put all grain charges to be equal. We also ignore the increase of the grain mass [9, 10] assuming that neutral atoms generated in the course of the surface electron-ion recombination escape from the grain surface into a plasma. Than, for the conditions  $l_i \gg \lambda_D$ , a, where  $l_i$  is the ion mean free path length and  $\lambda_D$  is the plasma screening (Debye) length, the friction and diffusion coefficients in the Fokker-Planck kinetic equation  $\beta(q, \mathbf{v})$  and  $D(q, \mathbf{v})$  [8] are given by:

$$\beta(q, \mathbf{v}) = -\sum_{\alpha=e,i} \frac{m_{\alpha}}{m_g} \int d\mathbf{v}' \frac{\mathbf{v} \cdot \mathbf{v}'}{v^2} \sigma_{\alpha g}(q, |\mathbf{v} - \mathbf{v}'|) |\mathbf{v} - \mathbf{v}'| \times f_{\alpha}(\mathbf{r}, \mathbf{v}', t),$$

A.G. Zagorodny

$$D(q, \mathbf{v}) = \sum_{\alpha=e,i} \frac{1}{2} \left(\frac{m_{\alpha}}{m_{g}}\right)^{2} \int d\mathbf{v}' \frac{(\mathbf{v} \cdot \mathbf{v}')^{2}}{v^{2}}$$
(1)  
 
$$\times \sigma_{\alpha g}(q, |\mathbf{v} - \mathbf{v}'|) |\mathbf{v} - \mathbf{v}'| f_{\alpha}(\mathbf{r}, \mathbf{v}', t),$$
  
 
$$\sigma_{g\alpha}(q, v) = \pi a^{2} \left(1 - \frac{2e_{\alpha}q}{m_{\alpha}v^{2}a}\right) \theta \left(1 - \frac{2e_{\alpha}q}{m_{\alpha}v^{2}a}\right).$$

Here  $\sigma_{g\alpha}(q, \mathbf{v})$  is the cross-section for grain charging within the orbital motion limited (OML) theory. In this approximation all electrons and ions approaching to the grain on the distance less than *a* are assumed to be absorbed. Subscript  $\alpha = e, i$  labels plasma particle species. The rest of notation is traditional. To include the processes of the electron, ion and atom scattering we have to summarize the appropriate coefficients on the different type of the processes. In the case of dominant charging collisions integration in Eq. (1) can be performed explicitly. The ion part of  $\beta$  (which exceeds the electron one at least in  $(m_i T_e/m_e T_i)^{1/2}$  times) is the following

$$\beta_i(q, \mathbf{v}) = -\sqrt{2\pi} \frac{m_i}{m_g} a^2 n_i v_{\mathrm{Ti}} \left[ I_1(\eta) + I_2(\eta, \Gamma) \right], \qquad (2)$$

$$I_1(\eta) = \left( -\frac{1}{2} + \frac{1}{4\eta} \right) \sqrt{\frac{\pi}{\eta}} \mathrm{Erf} \sqrt{\eta} - \frac{1}{2\sqrt{\eta}} e^{-\eta},$$

$$I_2(\eta, \Gamma) = \frac{\Gamma}{\eta} \left[ \frac{1}{2} \sqrt{\frac{\pi}{\eta}} \mathrm{Erf} \sqrt{\eta} - e^{-\eta} \right].$$

Here  $n_a$  are the densities of the electrons and ions,  $\operatorname{Erf}\sqrt{\eta}$ is the error function and  $\eta \equiv \eta(v) = v^2/2v_{\mathrm{Ti}}^2$ . It is easy to see that the terms  $I_1(\eta)$  and  $I_2(\eta, \Gamma)$  describe the parts of  $\beta_i(q, \mathbf{v})$  related to purely geometrical and charge-dependent collecting cross-sections, respectively. The integration for  $D_i(q, \mathbf{v})$  leads to the expression:

$$D(q, \mathbf{v}) = \frac{4}{3}\sqrt{2\pi} \left(\frac{m_i}{m_g}\right) a^2 n_i v_{\mathrm{Ti}} \left(\frac{T_i}{m_g}\right) \left[K_1(\eta) \qquad (3) + K_2(\eta, \Gamma)\right],$$

$$K_1(\eta) = \frac{3}{16\eta} \left[2(\eta - 1)e^{-\eta} + (2\eta^2 + \eta + 1)\sqrt{\frac{\pi}{\eta}} \mathrm{Erf}\sqrt{\eta}\right],$$

$$K_2(\eta, \Gamma) = \frac{3\Gamma}{8\eta}(\eta + 1) \left[-2e^{-\eta} + \sqrt{\frac{\pi}{\eta}} \mathrm{Erf}\sqrt{\eta}\right].$$

As follows from the physical reason and directly from Eq. (3), the coefficient  $D(q, \mathbf{v})$  is always positive. At the same time for some values of  $\eta$  and  $\Gamma$  the friction coefficient  $\beta_i(q, \mathbf{v})$  can be negative. To find the root  $\eta(\Gamma)$  of the equation  $\beta_i(\eta, \Gamma) = 0$  let us consider for two limiting cases  $\eta \ll 1$  and  $\eta \gg 1$ . For  $\eta \ll 1$  Eq. (2) gives

$$\beta_i(\eta, \Gamma) = 2A \left[ 1 - \Gamma - \frac{\eta}{5} (1 - 3\Gamma) \right], \qquad (4)$$

where

$$A = \frac{1}{3}\sqrt{2\pi} \left(\frac{m_i}{m_g}\right) a^2 n_i v_{\rm Ti}.$$

Equation (4) has a root  $\eta(\Gamma)$ , which exists and is small (according to the conditions of applicability for this expansion) only for  $\Gamma > 1$ , but  $\Gamma - 1 \ll 1$ :

$$\eta(\Gamma) = 5 \frac{(\Gamma - 1)}{3\Gamma - 1} \simeq \frac{5}{2} (\Gamma - 1).$$
(5)

Therefore the function  $\beta_i(\eta, \Gamma)$  is negative at  $\eta < \eta(\Gamma)$ , if  $\Gamma > 1$ . Eq. (4) shows that for  $\Gamma < 1$ , when  $\beta_i$  is positive for all  $\eta$ , the derivative  $(d\beta_i(\eta, \Gamma)/d\eta)|_{\eta=0}$  changes its sign at  $\Gamma = 1/3$ . Near  $\eta = 0$  the friction  $\beta_i$  decreases as function  $\eta$ , if  $\Gamma < 1/3$  and increases, if  $\Gamma > 1/3$ .



FIG. 1. Numerical solution of the equation  $\beta(\eta, \Gamma) = 0$ separating the positive and negative values of the friction coefficient.

For  $\eta \gg 1$  and arbitrary values of  $\Gamma$  Eq. (2) gives

$$\beta_i(\eta, \Gamma) = \frac{3}{2} A \sqrt{\frac{\pi}{\eta}} \left[ 1 - \frac{1+2\Gamma}{2\eta} \right]. \tag{6}$$

It means that for  $\Gamma \gg 1$  and large  $\eta$  the equation  $\beta_i(\eta_1\Gamma) = 0$  has a root  $\eta_1(\Gamma) \simeq (1+2\Gamma)/2 \sim \Gamma$ . Naturally, for all  $\Gamma > 1$  there exists appropriate  $\eta(\Gamma)$ , which is the root of that equation. This conclusion is confirmed by the exact numerical calculations of  $\eta(\Gamma)$  (Fig. 1). Asymptotically (for  $\eta \gg \max(1,\Gamma)$ )  $\beta_i(\eta,\Gamma)$  tends to zero as  $\sqrt{\eta^{-1}}$  and is positive for  $\eta > \eta(\Gamma)$ . For the case of the negative friction ( $\Gamma > 1$ ) the maximum of the coefficient  $\beta_i(\eta,\Gamma)$  is located at the point  $\eta_m(\Gamma) \gg 1$ :

$$\eta_m \simeq \frac{3}{2} \left( 1 + 2\Gamma \right), \qquad \beta_i(\eta_m, \Gamma) \simeq A \sqrt{\frac{2\pi}{3(1+2\Gamma)}}.$$
 (7)

For the diffusion coefficient  $D_i(\eta, \Gamma)$  the expansions for  $\eta \ll 1$  and  $\eta \gg 1$  lead to

$$D_i(\eta, \Gamma) = 4A\left(\frac{T_i}{m_g}\right) \left[1 + \frac{\Gamma}{2} + \frac{\eta}{10}(1+2\Gamma)\right], \ \eta \ll 1 \quad (8)$$
$$D_i(\eta, \Gamma) = 4A\left(\frac{T_i}{m_g}\right) \frac{3}{8}(\pi\eta)^{1/2} \left[1 + \frac{1}{2\eta}(1+2\Gamma)\right], \ \eta \gg 1.$$

The typical behaviour of  $\beta_i(\eta, \Gamma)$  and  $D_i(\eta, \Gamma)$  calculated numerically on the basis of Eqs. (2), (3) is shown in Figs. 2, 3.



FIG. 2. The velocity dependences of the dimensionless friction coefficient  $\tilde{\beta}(\eta, \Gamma) = I_1(\eta) + I_2(\eta, \Gamma)$  for the different values of  $\Gamma$ .



FIG. 3. The same dependences for the dimensionless diffusion coefficient  $\widetilde{D}(q, v) = K_1(\eta) + K_2(\eta, \Gamma)$ .

The stationary solution of the Fokker-Planck equation with the kinetic coefficients (2), (3) for the grain distribution function  $f_g(q, v)$  is

$$f_g(q,v) = \frac{C}{D_i(q,v)} \exp\left[-\int_0^v dv v \frac{\beta_i(q,v)}{D_i(q,v)}\right],\qquad(9)$$

where C is a constant, providing normalization  $\int d\mathbf{v} f_g(q, v) = 1$ . The velocity dependence of this solution for different values of  $\Gamma$  is shown in Fig. 4.

In order to get some analytical estimates let us consider the vicinity of the point  $\Gamma = 1$ . In such case the integration in Eq. (9) leads to the non-Maxwellian distribution function, which for  $\Gamma > 1$  possesses a maximum at  $v \neq 0$ :

$$f_g(q, \mathbf{v}) = \left(\frac{m_i}{2\pi T_i}\right)^{3/2} \frac{\sqrt{\pi}}{2Y_{3/2}(1 + \Gamma/2)} \left[1 \qquad (10)\right]$$

$$+ \frac{m_i v^2}{2T_i^*} \frac{1}{5} (1+2\Gamma) \Big]^{-1} \exp\left\{-\frac{m_g v^2}{2T_i^*} \Big[ (1-\Gamma) \right]^{-1} \exp\left\{-\frac{m_g v^2}{2T_i^*} \Big]^{-1} + \frac{1}{5} \left[ (1-1) \right]^{-1} + \frac{1}{5} \left[$$

$$+ \frac{m_i v^2}{2T_i^*} \frac{1}{10} \left( 5\Gamma^2 + 4\Gamma - 3 \right) \right] \bigg\} \,.$$

Here

$$Y_{\nu}(\Gamma) = \int_{0}^{\infty} d\eta \eta^{\nu-1} \exp\left[-\alpha\eta(1-\Gamma) - \gamma\eta^{2}\right]$$
(11)  
$$= (2\gamma)^{-\frac{\nu}{2}} \frac{\sqrt{\pi}}{2} \exp\left[\frac{\alpha^{2}(1-\Gamma)^{2}}{8\gamma}\right] D_{-\nu} \left(\frac{\alpha(1-\Gamma)}{\sqrt{2\gamma}}\right)$$
$$T_{i}^{*} = 2T_{i} \left(1 + \frac{\Gamma}{2}\right), \qquad \alpha = \frac{m_{g}}{2m_{i}} \left(1 + \frac{\Gamma}{2}\right),$$
$$\gamma = \frac{\alpha(5\Gamma^{2} + 4\Gamma - 3)}{10(2+\Gamma)}$$

and  $D_{\nu}$  is the cylindrical parabolic function. Eq. (10) is relevant, if the integral over  $\eta$  converges, what is valid for positive  $\gamma(\Gamma > 0, 472)$ . At the same time, as was mentioned above, for the applicability of the expansions (4), (8) the inequality  $\Gamma < 1 + \delta$  with  $\delta \ll 1$  is required. It is clear from Eqs. (6), (8), that the asymptotic behaviour of the distribution function is non-exponential  $f_g(\eta)|_{\eta\to\infty} \sim \eta^{-(m_g/m_i)}$ , but for the values of  $\Gamma$  under consideration it is not essential for calculations of the averages, due to the rapid convergence of the integrals over  $\eta$ . In particular, the average kinetic energy K of grains is

$$K(\Gamma) = \left(\frac{m_g}{m_i}\right) T_i \frac{Y_{5/2}(\Gamma)}{Y_{3/2}(\Gamma)}$$
(12)

that gives  $K = 2,86(m_g/m_i)^{1/2}T_i$  for  $\Gamma = 1$ . Thus, the ion absorption by grains can lead to grain heating and grain average kinetic energy (or their effective temperature) can be much higher, than the electron and ion temperatures. This effect can be applied to the explanation of the experimental data [11, 12], as it was already suggested in [7,8] on the basis of the velocity independent approximation for  $\beta$  and D.

Let us consider now the case  $\Gamma \gg 1$ . In this case the ratio  $Q = -(T_i/m_i)\beta(\eta,\Gamma)/D(\eta,\Gamma)$  can be represented for all values of  $\eta$  with a good accuracy by the function  $Q = (m_q/m_i)(\Gamma - \eta)/[(1 + \Gamma)(\eta + \Gamma)]$  that gives

$$f_g = \frac{C}{D(\eta, \Gamma)} \exp\left\{\frac{m_g}{m_i} \left[\frac{(3\Gamma+1)}{(2\Gamma-1)} \ln \frac{\Gamma(\eta+1)}{\Gamma+\eta} - \frac{1}{2} \ln \frac{\eta^2 + (\Gamma+1)\eta + \Gamma}{\Gamma}\right]\right\},$$
(13)

where

$$D_i(\eta, \Gamma)|_{\Gamma \gg 1} \simeq \frac{3}{2} A \frac{T_i}{m_g} \sqrt{\pi \eta} \left[ 1 + \frac{\Gamma}{(\eta + \frac{3}{4}\sqrt{\pi \eta})} \right]. \quad (14)$$

It is evident, that  $f_g(\eta, \Gamma)$  given by Eq. (13) is non-exponential.

Finally for the domain  $\Gamma < 0,472$  we can omit the term  $\sim v^4$  in Eq. (10) and the distribution  $f_g(q, v)$  becomes Maxwellian with the effective grain temperature:

$$T_{\rm eff} = \frac{2T_i^*}{1-\Gamma} = \frac{2T_i(1+\frac{\Gamma}{2})}{1-\Gamma}.$$
 (15)

The above conclusions are in a good agreement with the results of numerical calculations of  $f_g(\eta, \Gamma)$  on the basis of Eq. (9) (Fig. 4).



FIG. 4. The same dependences for the distribution function  $f_g(\eta, \Gamma)$ .

If we take into account the processes of atom-grain and ion-grain elastic scattering the coefficients  $\beta(\eta, \Gamma)$ and  $D(\eta, \Gamma)$  should include additional terms. To estimate, for example, contribution of atom-grain and iongrain scattering we can consider  $\beta(\eta)$  calculated on the basis of the Eq. (2) with the appropriate transport crosssections. For the case  $\eta \ll 1$  the friction coefficient is

$$\beta(\eta, \Gamma) = 2A \left[ 1 - \Gamma + 3 \frac{n_a}{n_i} \left( \frac{T_a m_a}{T_i m_i} \right)^{1/2} - \frac{\eta}{5} (1 - 3\Gamma) + 6\Gamma^2 \ln \Lambda \right].$$
(16)

The negative friction exists for small  $\eta$ , if the Coulomb scattering is strongly suppressed, when the Coulomb logarithm ln  $\Lambda$  is small [13–15], that is typical for strong interaction. The root of the function  $\beta(\eta, \Gamma)$  is shifted to the region of large,  $\Gamma$ , which is determined by the atom density  $n_a$ . The result of rough estimate for the case  $\lambda_{\rm Li} > \lambda_D > a$  ( $\lambda_{\rm Li}$  is the Landau length) gives

$$\eta(\Gamma) = \frac{5}{3\Gamma - 1} \left[ \Gamma - 1 - 3\sqrt{\frac{Z_g\Gamma}{\pi n_i a^3}} - \frac{3n_a}{n_i} \left( \frac{T_a m_a}{T_i m_i} \right)^{1/2} \right].$$
(17)

In general, a more detail consideration of scattering processes and the effects of strong interactions between the grains [15], is needed for the exact description of the region of negative friction. In conclusion, ion absorption by grains can generate negative friction and provide the substantial increase of the average grain kinetic energy in comparison with the temperatures of the other plasma components. Microscopical justification of negative friction on kinetic level is presented and deviation of the grain distribution function from the Maxwellian distribution is found. For dusty plasma, as for an open system, the fluctuation-dissipation theorem in the form of Einstein relation is not applicable.

## ACKNOWLEDGMENT

The authors thank W. Ebeling, U. Erdmann, L. Schimansky-Geier, and P.P.J.M. Schram for valuable discussions and the Netherlands Organization for Scientific Research (NWO) for the support of this work. We also appreciate V. Kubaichuk for the help in numerical calculations.

- J.W. Rayleigh. *The Theory of Sound*, vol. I, 2nd edition (Dover, New-York) 1945.
- [2] P-G. de Gennes. Rev. Mod. Phys. 57 827 (1985).
- [3] A.S. Mikhailov, and D. Meinköhn. In: Stochastic Dynamics, edited by L. Schimansky- Geier, T. Pöschel, vol. 484 of Lectures Notes in Physics (Springer, Berlin, 1997) p. 334.
- [4] M. Scheinbein, and H. Gruler. Bull. Math. Biology 55 585 (1993).
- [5] F. Schweitzer, W. Ebeling, and B. Tilch. Phys. Rev. Lett. 80 5044 (1998).
- [6] U. Erdmann, W. Ebeling, L. Schimansky-Geier, and F. Schweitzer. Eur. Phys. J. B15 105 (2000).
- [7] A.G. Zagorodny, P.P.J.M. Schram, and S.A. Trigger. Phys. Rev. Lett. 84 3594 (2000).
- [8] P.P.J.M. Schram, A.G. Sitenko, S.A. Trigger, and A.G. Zagorodny. Phys. Rev. E 63 016403 (2000).
- [9] S.A. Trigger. Contrib. Plasma Phys. **41** 331 (2001).
- [10] A.M. Ignatov, S.A. Trigger, W. Ebeling, and P.P.J.M. Schram. Phys. Lett. A293 141 (2001).
- [11] A. Melzer, A. Homann, and A. Piel. Phys. Rev. E 53 2757 (1996).
- [12] H.M. Thomas, and G.E. Morfill. Nature **379** 806 (1996).
- [13] M.D. Kilgore, J.E. Daugherty, R.K. Porteous, and D.B. Graves, J. Apply Phys. **73** 7195 (1993).
- [14] S.A. Khrapak, A.V. Ivlev, G.E. Morfill, and H.M. Thomas. *Ion drag force in complex plasmas*. In: 29 EPS Conference, abstract and report, Montreux, June 2002, to be published.
- [15] S.A. Trigger, G.M.W. Kroesen, P.P.J.M. Schram, E. Stoffels, and W.W. Stoffels. *Ion drag and Brownian motion in dusty plasmas*. In: 29 EPS Conference, abstract and report, Montreux, June 2002, to be published.