Distances and classification of amino acids for different protein secondary structures

Xin Liu[†],Li-mei Zhang^{*}, Shan Guan[‡], Wei-Mou Zheng[†]

[†]Institute of Theoretical Physics, China, Beijing 100080, China *School of Science at North Jiaotong University, Beijing 100044, China [‡]Center of Bioinformations at Peking University, Beijing 100871, China

Abstract

Window profiles of amino acids in protein sequences are taken as a description of the amino acid environment. The relative entropy or Kullback-Leibler distance derived from profiles is used as a measure of dissimilarity for comparison of amino acids and secondary structure conformations. Distance matrices of amino acid pairs at different conformations are obtained, which display a non-negligible dependence of amino acid similarity on conformations. Based on the conformation specific distances clustering analysis for amino acids is conducted.

PACS number(s): 87.10.+e,02.50.-r

1 Introduction

The similarity of amino acids(aa) is the basis of protein sequence alignment, protein design and protein structure prediction. Several scoring schemes have been proposed based on amino acid similarity. The mutation data matrices of Dayhoff [6] and the substitution matrices of Henikoff [1] are standard choices of scores for sequence alignment and amino acid similarity evaluation. However, these matrices, focusing on the whole protein database, pay little attention on protein secondary structures(*ss*). How the amino acid similarity is influenced by different secondary structures is an interesting question. Furthermore, understanding the differences can help us in protein sequence analysis. Despite efforts in uncovering the information encoded in the primary structure, we still cannot read the language describing the final 3D fold of an active biological macromolecule. Compared with the DNA sequence, a protein sequence is generally much shorter, but the size of the alphabet is five times larger. A proper coarse-graining of the 20 amino acids into fewer clusters for different conformation is important for improving the signal-to-noise ratio when extracting information by statistical means.

It is our purpose to propose a simple scheme to study amino acid similarity from amino acid string statistics. Information about the environment for an amino acid at a certain conformation state may be provided by statistics of residue strings or windows centered at the amino acid. The success of window-based approaches such as GOR [2] for secondary structure prediction validates the use of such statistics. We shall derive a measure for the difference of amino acid pairs based on the distance of probability distributions, and investigate how the difference is dependent on conformations.

2 Amino acid distances

Our discussion will be heavily based on the distance between two probability distributions. A well defined measure of the distance is the Kullback-Leibler (KL) distance or relative entropy [7, 8, 9], which, for two distributions $\{p_i\}$ and $\{q_i\}$, is given by

$$d(\{p_i\},\{q_i\}) = \sum_{i} p_i \log(p_i/q_i).$$
(1)

It corresponds a likelihood ratio, and, if p_i is expanded around q_i , its leading term is the χ^2 distance:

$$d_{\chi}(\{p_i\},\{q_i\}) = \sum_{i} (p_i - q_i)^2 / p_i.$$
⁽²⁾

It is often to use the following symmetrized form for the KL distance

$$D(\{p_i\},\{q_i\}) = \frac{1}{2}[d(\{p_i\},\{q_i\}) + d(\{q_i\},\{p_i\})].$$
(3)

The distributions to be considered here come from window statistics. For a given amino acid residue $a_i = x$ at the conformation state α in a sequence $a_1 a_2 \cdots a_i \cdots$, we take the string $a_{-n+i}a_{-n+i+1} \cdots a_i \cdots a_{i+n}$ of width (2n + 1) as a window. Denote by $N_k(y|x, \alpha)$ the count of residue y at the k-th site from the center of such windows. As in GOR, only the conformation of the central residue is concerned. A quantity derived from $N_k(y|x, \alpha)$ is

$$N(x,\alpha) = \sum_{y} N_k(y|x,\alpha),\tag{4}$$

which, as the total count of residue x at the conformation α , is independent of k. The conditional probability distribution $P_k(y|x, \alpha)$ is estimated as

$$P_k(y|x,\alpha) = \frac{N_k(y|x,\alpha)}{N(x,\alpha)}.$$
(5)

The weight matrix $\mathbf{M}_{20 \times 2n}$ with its entries being $P_k(y|x, \alpha)$ is the so-called residue profile of x at α . Such profiles are used in window-based approaches, e.g. GOR and artificial neural network algorithm [12].

We expect that on average the correlation between the central residue and an outer site decays when they become far apart in sequence. To examine the correlation, we consider a large window width of 21, i.e. n = 10, and take the 'noise' background to be the following average:

$$Q(y|x,\alpha) = \frac{1}{6} \left[\sum_{k=-10}^{-8} P_k(y|x,\alpha) + \sum_{k=8}^{10} P_k(y|x,\alpha) \right].$$
 (6)

The KL distance $D_{k;x,\alpha}(\{P_k(y|x,\alpha)\}, \{Q(y|x,\alpha)\})$ provides a measure of the correlation between the central site and site k. As we shall see, for our purpose of amino acid comparison a narrow window of a strong correlation with width of 7 is used to describe amino acid environment.

Using distribution $P_k(y|x, \alpha)$ from window statistics to characterize amino acid residues, we define the distance of residue pair x and y at the same conformation α as the following sum of KL distances

$$D_{xy;\alpha} = \sum_{k=\pm 1,\pm 2,\pm 3} D(\{P_k(z|x,\alpha)\}, \{P_k(z|y,\alpha)\}).$$
(7)

Similarly, to explore the difference of the same residue x at different conformations α and β , we may define the distance

$$D_{\alpha\beta;x} = \sum_{k=\pm 1,\pm 2,\pm 3} D(\{P_k(z|x,\alpha)\}, \{P_k(z|x,\beta)\}).$$
(8)

By means of the residue pair distances we can further study the classification of amino acids. With the KL distance, we may define the cluster distance in a way consistent with that for residue pairs. For example, we characterize the cluster consisting of residues x and y by the 'coarse-grained' probability

$$P_k(z|x\&y,\alpha) = \frac{N_k(z|x,\alpha) + N_k(z|y,\alpha)}{N(x,\alpha) + N(y,\alpha)}.$$
(9)

We then may define the distance between this cluster and some other residues or clusters. With cluster distance defined, the cluster analysis can be used to reduce amino acid alphabets.

3 Results

Our analysis is performed on a data set taken from the database PDB_SELECT[3, 4] of nonredundant protein sequences with known structures. The sequences share amino acid identity less than 25%. We keep only the non-membrane sequences with their lengths between 80 and 420. The secondary structure assignment is taken from the DSSP database [5]. As in GOR, we use the following reduction of the 8 DSSP states to 4 states of helix(h), sheet(e), coil(c) and turn(t): $H, G, I \rightarrow h, E \rightarrow e, X, S, B \rightarrow c$ and $T \rightarrow t$. The counts of each amino acid at the reduced four different conformation states are given in Table 1.

We first estimate probability distributions of residues for each central residue at a given conformation. At this step, the window width is 21. We then calculate distances $D_{k;x,\alpha}(\{P_k(y|x,\alpha)\}, \{Q(y|x,\alpha)\})$ of these distributions to their corresponding noise distributions. The results are shown in Figs. 1 to 4, each of which is for one conformation of the central residue. The 20 curves in each figure correspond to 20 central amino acids. Due to the sample size difference, curves are not directly comparable. (Roughly speaking, under the null hypothesis of identical distribution the χ^2 distance should be scaled with the sample size, so a small sample size would give a relatively large distance.) However, a decay is clearly seen when the site k become far away from the center. For more discussions on correlations we refer reader to [10, 11]. As seen from most curves of the figures, distances at the 6 sites nearest to the center are significantly larger than those at window border sites. We shall use window width of 7 for further comparison of amino acids.

It is natural to expect that similar residues would have similar window statistics. Thus, the KL distance between two residue profiles provides a measure of their similarity, i.e. a small KL distance implies a large similarity. We calculate the KL distance matrices $D_{xy;\alpha}$ for residue pairs at different conformations with formula (7). The results are given in Tables 2 and 3, where entries have been multiplied by a factor 200. With the distributions (9) defined for clusters, we further perform the simplest bottom-up approach of hierachical clustering for residues, by starting from 20 clusters of single residues, and then joining two nearest clusters step by step until a single cluster is obtained. The results of clustering are given in Tables 4 to 7. Since the dendritic trees returned from clustering are less informative, for visualization we introduce graphs where vertices are the 20 amino acids, and an edge exists between a pair of amino acids if and only if their distance is below some preset threshold. Graphs obtained from the distance matrices are shown in Figs. 5 to 8, where vertices with no connecting edges are neglected.

In sequence pair alignment we often do not have structure information of both sequences. With the

structure information ignored, we have the mixed counts

$$N_k(y|x) = \sum_{\alpha} N_k(y|x,\alpha), \tag{10}$$

from which we calculate the residue pair distances averaged over conformations. The distance matrix obtained is given in Table 8. We have also calculated distances (8) to compare different conformations. Distances between any two conformations for various residues are listed in Table 9.

4 Discussions

Figures 1 to 4 illustrate the dependence of outer sites in a window on the center. Although in the KL distance we sum up effects on individual residues from the center, we still can see the tendency that the center is generally more strongly correlated with the C-terminal sites than N-terminal sites. Furthermore, we may divide the 20 amino acids into two groups with M, I, L, V, F, Y and W in one, and the remainders in the other. They roughly correspond to hydrophobic and hydrophilic groups. It is seen that for the coil and turn conformations a hydrophobic center exhibits a stronger correlation with outer sites than a hydrophilic center, while for the sheet conformation a hydrophilic center exhibits a stronger correlation.

It is interesting to make a comparison between the distance matrices obtained here with the commonly used BLOSUM62 similarity score matrix. A small distance implies a large similarity score. There are many evidences showing the consistency between the distances and scores. For example, residue pairs VI, IL, VL and ST have positive BLOSUM scores and at the same time small distances. The graphs in Figs. 5 to 8 contain two connected subgraphs: one consists of I, L, V, F, Y, and the other consists of S, T. This is another evidence of the consistency. Generally, the averaged distance matrix is closer to BLOSUM62 than the conformation specific ones. However, there do exist some remarkable differences. For example, residue pairs GT, QA, FV with negative scores have rather small distances in either the conformation helix, or sheet or coil, while pairs YH and NH with positive scores have rather large distances in the helix conformation. Moreover, YH has a large distance in all the four conformations.

BLOSUM matrices are derived from conserved amino acid patterns called blocks. It is expected that for most score entries we should see the consistency in at least one conformation specific distance matrix. For a given residue pair, if residue profiles of an amino acid center are very dissimilar for different conformations, after averaging over conformations the pair distance would generally become smaller. In this case, BLOSUM scores and conformation specific distance need not be consistent since the former contains no structure information.

Our results show some strong dependence of residue behavior on conformations. For example, the distances of pairs CD and SI in helix are about twice higher than in sheet. There are many residue pairs displaying strong dependence of distances on conformations. Table 9 views the conformation dependence from conformation pair comparison. Indeed, the table indicates that for any conformation pairs there are certain residues which behave very differently in the two conformations. However, generally speaking, coil and turn are quite similar.

In comparison of physicochemical properties of amino acids, the abundance of amino acids is not taken into consideration. This is also the case for the above defined distances. Other statistical variables including the effect of sample size may be introduced. One candidate is the χ^2 statistic for identical distributions. The analysis using this new statistic is under study.

We expect that algorithms using multiple conformation specific matrices should work better in sequence alignment. The popular Needleman-Wunsch algorithm can be modified to include putative conformation for each residue. This will be discussed elsewhere.

This work was supported in part by the Special Funds for Major National Basic Research Projects and the National Natural Science Foundation of China.

References

- [1] S. Henikoff and J.G. Henikoff, Proc. Natl. Acad. Sci. (USA), 89, 10915 (1992).
- [2] J. Garnier, D. Osguthorpe and B. Robson, J.Mol.Biol. 120, 97 (1978).
- [3] U. Hobohm and C. Sander, Protein Science 3,522 (1994).
- [4] ftp.embl-heidelberg.de/pub/databases/protein_extras/pdb_select/old/2001_sep.25
- [5] W. Kabsch and C. Sander, Biopolymers, 22, 2577 (1983).
- [6] M. O. Dayhoff and R. V. Eck, Atlas of Protein Sequence and Structure (Natl. Biolmed. Res. Found., Silver Spring, MD), 3, 33 (1968).
- [7] S. Kullback, J.C. Keegel and J.H. Kullback, Information Theory and Statistics, Wiley, New York (1959).

- [8] S. Kullback, Topics in Statistical Information Theory, Springer, Berlin (1987).
- [9] T.Sakamoto, M. Ishiguro and G.Kitagawa, Akaike Information Criterion Statistics, KTK Scientific, Tokyo (1986).
- [10] O. Weiss and H. Herzel, J. Theor. Biol, **190**, 341 (1998).
- [11] O. Weiss, M.A. Jiménez-Montaño and H. Herzel, J. Theor. Biol, 206, 379 (2000).
- [12] P. Stolorz, A. Lapedes and Y. Xia, J.Mol.Biol, 225, 363 (1992).

Table 1. Sample sizes of each amino acid residue in different protein secondary structures.

-	h	е	с	t
С	690	732	822	224
\mathbf{S}	2841	1764	3538	1179
Т	2350	2288	3112	762
Р	1173	624	3648	1302
Α	5950	2019	2651	1122
G	1795	1633	4328	3090
Ν	1904	922	2692	1388
D	2841	1029	3621	1424
\mathbf{E}	4773	1514	2325	1172
\mathbf{Q}	2757	1008	1532	653
Η	1132	794	1148	426
\mathbf{R}	3108	1469	1948	771
Κ	3861	1579	2645	1187
Μ	1390	693	679	223
Ι	3169	3333	1719	368
\mathbf{L}	6262	3307	2952	850
\mathbf{V}	3233	4461	2330	487
\mathbf{F}	2225	1948	1545	444
Y	1806	1773	1303	459
W	827	632	536	173

5	Table 2.	Am	ino aci	id dist	ance 1	natrice	es for	helices	(bott	om-lef	t) and	turns	(top-	right).	Entr	ies hav	ve beer	n mult	tiplied	by a
facto	r 200.																			
C		106	116	135	118	145	134	132	121	129	111	118	124	154	134	121	119	104	123	215
S	64		23	52	29	59	35	33	36	37	54	26	36	78	61	38	49	45	40	100
T	63	13		61	33	74	40	35	39	46	62	33	37	92	63	40	46	45	38	93
P	81	48	49		44	99	71	69	54	62	82	47	55	106	89	71	71	62	66	132
A	45	21	17	63		64	38	39	32	36	58	29	33	63	64	34	46	48	43	98
G	57	15	20	52	25		32	39	54	55	57	47	52	81	79	61	88	75	70	115
N	82	14	22	67	33	26		18	30	31	44	29	31	72	68	38	63	53	36	96
D	101	17	26	56	39	32	16		25	34	44	30	29	77	58	36	54	49	37	91
E	82	20	25	56	27	36	22	14		33	53	37	23	73	65	46	51	59	43	106
Q	70	16	21	60	19	28	17	21	14		51	32	38	79	66	51	62	62	49	100
\tilde{H}	55	23	24	55	26	26	33	35	34	28		51	58	90	79	54	81	70	55	113
R	69	21	22	59	21	30	22	28	24	13	28		30	71	69	38	54	49	48	101
K	80	21	25	67	28	38	22	27	23	19	38	13		81	63	38	49	51	47	102
M	48	57	45	85	23	56	75	82	64	51	50	50	60		93	62	85	93	78	141
Ι	43	81	65	104	35	78	104	116	88	76	66	73	79	22		47	55	55	54	104
L	35	65	52	90	26	62	83	99	73	59	53	56	67	15	10		49	36	31	85
V	37	59	44	81	22	55	77	90	67	53	52	51	60	16	12	09	-	46	58	99
\overline{F}	34	67	53	87	30	61	90	99	79	69	54	66	75	22	17	12	15		48	100
\overline{Y}	44	43	35	77	23	47	64	71	55	47	34	47	54	26	29	21	$\frac{1}{22}$	16		- 90
Ŵ	49	61	53	82	35	64	87	92	72	58	57	60	71	31	35	27	29	25	24	00
	C	S	T	P	A	G	N	D	Ē	$\overset{\circ\circ}{Q}$	H	R	K	M	I	Ĺ	V	F	Y	W

Table 3. Amino acid distance matrices for sheets (bottom-left) and coils (top-right). Entries have been multiplied by a factor 200.

C		43	51	70	36	47	51	66	50	51	46	48	54	55	51	44	42	47	48	66
S	42		10	24	17	17	16	21	20	14	30	15	19	32	38	27	24	30	32	45
T	49	15		28	19	24	16	21	17	16	34	17	17	26	32	24	22	28	31	46
P	68	42	46		37	28	25	22	31	28	48	28	31	52	67	59	49	62	61	71
A	33	20	24	42		16	22	29	16	16	30	17	21	23	27	17	15	24	25	41
G	35	29	37	62	16		18	23	31	28	36	17	31	31	44	17	27	31	32	44
N	51	23	27	46	30	37		14	19	19	30	19	20	34	42	34	31	31	34	54
D	54	24	31	46	32	42	23		22	23	39	23	22	46	56	48	41	46	51	65
E	60	21	19	48	32	47	26	24		14	32	14	11	25	30	25	17	25	24	39
Q	52	20	17	53	28	41	29	30	22		32	14	14	30	33	26	21	26	25	38
H	50	27	26	54	28	33	34	33	30	28		29	36	51	44	41	39	44	40	67
R	46	21	20	44	20	33	32	31	21	23	22		16	31	33	24	21	28	29	47
K	62	29	20	52	30	47	35	34	20	23	35	24		34	34	28	21	29	26	50
M	38	45	44	65	24	33	52	62	50	46	44	38	52		28	22	22	27	30	39
Ι	32	38	36	62	24	35	56	57	49	41	40	36	43	23		12	15	16	20	34
L	27	37	34	58	19	29	50	55	45	41	37	32	43	20	09		10	12	17	33
V	31	35	32	58	19	27	51	57	46	40	36	32	38	22	09	10		14	17	29
F	29	45	44	71	25	33	62	67	59	47	49	42	56	28	14	12	15		18	33
Y	32	35	32	64	24	33	51	54	47	34	33	31	42	29	13	13	15	14		31
W	46	57	58	71	47	60	69	76	62	52	54	57	66	48	39	39	38	33	37	
	C	S	T	P	Α	G	N	D	E	Q	H	R	K	M	Ι	L	V	F	Y	W

Table 4. Clustering of amino acid alphabets for helices. The first column indicates the number of amino acid groups.

Table 5. Clustering of amino acid alphabets for sheets. The first column indicates the number of amino acid groups.

19	AGFILVY	М	D	Ε(QS	Т	R	Κ	Η	Ν	С	W	Ρ
18	A G F ILV Y	М	D	ΕO	QS	Т	R	Κ	Η	Ν	С	W	Ρ
17	A G FILV Y	М	D	Ε(QS	Т	R	Κ	Η	Ν	С	W	Ρ
16	A G FILVY	М	D	ΕO	QS	Т	R	Κ	Η	Ν	С	W	Ρ
15	A G FILVY	М	D	ΕO	Q SI	[R	Κ	Η	Ν	С	W	Ρ
14	A G FILVY	М	D	ΕO	QST		R	Κ	Η	Ν	С	W	Ρ
13	A G FILVY	М	D	EQ	ST		R	Κ	Η	Ν	С	W	Ρ
12	A G FILVY	М	D	EQ	STR			Κ	Η	Ν	С	W	Ρ
11	A G FILVY	М	D	EQ	STRF	ζ			Η	Ν	С	W	Ρ
10	AG FILVY	М	D	EQ	STRF	ζ			Η	Ν	С	W	Ρ
9	AGFILVY	М	D	EQ	STRF	ζ			Η	Ν	С	W	Ρ
8	AGFILVYM		D	EQ	STRF	ζ			Η	Ν	С	W	Ρ
7	AGFILVYM		D	EQ	STRF	ΚH				Ν	С	W	Ρ
6	AGFILVYM		D	EQ	STRF	KHN	1				С	W	Ρ
5	AGFILVYM		DF	EQS	TRKI	IN					С	W	Ρ
4	AGFILVYMDEQS'	ΓRI	KHI	N							С	W	Ρ
3	AGFILVYMDEQS	ΓRI	KHI	NC								W	Ρ
2	AGFILVYMDEQS'	ΓRI	KHI	NCW									Ρ

Table 6. Clustering of amino acid alphabets for coils. The first column indicates the number of amino acid groups.

Table 7. Clustering of amino acid alphabets for turns. The first column indicates the number of amino acid groups.

19 18	A DN E K S T R Q L A DN E K ST R Q L	Y Y	F F	V V	H H	G G	I	P P	M M	W W	C C
17	A DN EK ST RQL	Y	F	V	Н	G	Ι	Ρ	М	W	C
16	ADNEK ST RQL	Y	F	V	Η	G	Ι	Ρ	М	W	С
15	A DNEKST R Q L	Y	F	V	Η	G	Ι	Ρ	М	W	С
14	A DNEKSTR Q L	Y	F	V	Η	G	Ι	Ρ	М	W	С
13	ADNEKSTR Q L	Y	F	V	Η	G	Ι	Ρ	М	W	С
12	ADNEKSTRQ L	Y	F	V	Η	G	Ι	Ρ	М	W	С
11	ADNEKSTRQL	Y	F	V	Η	G	Ι	Ρ	М	W	С
10	ADNEKSTROLY		F	V	Η	G	Ι	Ρ	М	W	С
9	ADNEKSTRQLYF			V	Η	G	Ι	Ρ	М	W	С
8	ADNEKSTROLYFV				Η	G	Ι	Ρ	М	W	С
7	ADNEKSTROLYFVH					G	Ι	Ρ	М	W	С
6	ADNEKSTROLYFVHG						Ι	Ρ	М	W	Ċ
5	ADNEKSTROLYFVHGI							Ρ	М	W	Ċ
4	ADNEKSTROLYFVHGIP								М	W	Ċ
3	ADNEKSTRÖLYFVHGIPM	[Ŵ	С
2	ADNEKSTRQLYFVHGIPM	W									С



C																				
S	21																			
T	25	5																		
P	25	9	11																	
A	29	12	12	16																
G	21	8	11	11	11															
N	25	$\overline{7}$	9	13	12	8														
D	32	9	9	15	10	11	6													
E	40	18	18	21	11	18	14	9												
Q	34	12	12	18	8	14	10	9	8											
H	21	13	14	17	18	14	12	15	23	17										
R	31	11	13	16	7	13	11	10	9	5	15									
K	35	15	14	18	12	16	10	9	8	10	22	8								
M	33	19	16	20	10	17	18	18	19	16	24	15	18							
Ι	25	16	13	16	12	14	16	17	20	18	19	16	15	10						
L	26	16	14	17	9	14	16	17	19	15	20	14	15	8	4					
V	24	10	9	13	8	9	11	12	15	13	17	12	12	10	6	6				
F	22	13	11	16	13	11	14	16	20	18	18	16	15	12	6	6	6			
Y	24	9	9	13	13	10	11	14	19	15	14	15	14	13	8	9	7	5		
W	32	20	19	20	21	17	22	25	29	23	24	24	27	18	14	13	13	10	12	
	C	S	T	P	A	G	N	D	E	Q	H	R	K	M	Ι	L	V	F	Y	W



Figure 1: KL distances (doubled) of outer sites from their corresponding noise background. Each curve is for an amino acid at the center labeled 0, whose conformation is turn. For clarity, the curves for M,I,L,V,F,Y and W have been shifted up by multiplying an extra factor 100.

Table 9. Conformation pair distances for each amino acid. Entries have been multiplied by a factor 200. (h: Helix, e: Sheet, c: coil, t: Turn.)

	he	hc	ht	ec	\mathbf{et}	ct
С	133	185	163	127	197	139
\mathbf{S}	93	129	124	93	148	73
Т	98	120	131	103	175	96
Р	172	118	121	89	233	116
Α	112	148	127	122	149	73
G	79	101	80	91	107	57
Ν	126	145	118	106	152	76
D	149	137	149	93	174	81
Ε	159	152	138	109	192	73
\mathbf{Q}	130	157	133	93	143	93
Η	100	150	110	117	152	98
R	131	146	128	91	144	85
Κ	137	149	128	93	155	88
Μ	130	161	147	126	156	135
Ι	138	180	134	118	130	110
\mathbf{L}	143	162	113	127	148	98
V	114	151	151	98	147	101
\mathbf{F}	120	150	111	107	115	88
Υ	95	147	96	111	117	80
W	120	181	201	123	173	111



Figure 2: KL distances (doubled) of outer sites from their corresponding noise background. Each curve is for an amino acid at the center labeled 0, whose conformation is coil. For clarity, the curves for M,I,L,V,F,Y and W have been shifted up by multiplying an extra factor 100.



Figure 3: KL distances (doubled) of outer sites from their corresponding noise background. Each curve is for an amino acid at the center labeled 0, whose conformation is sheet. For clarity, the curves for M,I,L,V,F,Y and W have been shifted up by multiplying an extra factor 100.



Figure 4: KL distances (doubled) of outer sites from their corresponding noise background. Each curve is for an amino acid at the center labeled 0, whose conformation is helix. For clarity, the curves for M,I,L,V,F,Y and W have been shifted up by multiplying an extra factor 100.



Figure 5: Connecting graph of amino acids in helix. Edges exist only between vertices with a scaled distance not greater than 20. Vertices without any connecting edges are not shown.



Figure 6: Connecting graph of amino acids in sheet. Edges exist only between vertices with a scaled distance not greater than 20. Vertices without any connecting edges are not shown.



Figure 7: Connecting graph of amino acids in coil. Edges exist only between vertices with a scaled distance not greater than 17. Vertices without any connecting edges are not shown.



Figure 8: Connecting graph of amino acids in turn. Edges exist only between vertices with a scaled distance not greater than 35. Vertices without any connecting edges are not shown.