

# FAST CALCULATIONS IN NONLINEAR COLLECTIVE MODELS OF BEAM/PLASMA PHYSICS

**Antonina N. Fedorova, Michael G. Zeitlin**

*IPME RAS, St. Petersburg, V.O. Bolshoj pr., 61, 199178, Russia*  
*e-mail: zeitlin@math.ipme.ru*  
*e-mail: anton@math.ipme.ru*  
*<http://www.ipme.ru/zeitlin.html>*  
*<http://www.ipme.nw.ru/zeitlin.html>*

We consider an application of variational-wavelet approach to nonlinear collective models of beam/plasma physics: Vlasov/Boltzmann-like reduction from general BBGKY hierarchy. We obtain fast convergent multiresolution representations for solutions which allow to consider polynomial and rational type of nonlinearities. The solutions are represented via the multiscale decomposition in nonlinear high-localized eigenmodes (waveletons).

Presented at VIII International Workshop on  
Advanced Computing and Analysis Techniques in Physics Research,  
Section III "Simulations and Computations in  
Theoretical Physics and Phenomenology"  
ACAT'2002, June 24-28, 2002, Moscow

# Fast Calculations in Nonlinear Collective Models of Beam/Plasma Physics

A. N. Fedorova and M. G. Zeitlin <sup>a</sup>

<sup>a</sup>IPME, RAS, V.O. Bolshoj pr., 61, 199178, St. Petersburg, Russia

E-mail: zeitlin@math.ipme.ru, <http://www.ipme.ru/zeitlin.html>, <http://www.ipme.nw.ru/zeitlin.html>

We consider an application of variational-wavelet approach to nonlinear collective models of beam/plasma physics: Vlasov/Boltzmann-like reduction from general BBGKY hierarchy. We obtain fast convergent multiresolution representations for solutions which allow to consider polynomial and rational type of nonlinearities. The solutions are represented via the multiscale decomposition in nonlinear high-localized eigenmodes (waveletons).

We consider applications of numerical-analytical technique based on variational-wavelet approach to nonlinear collective models of beam/plasma physics, e.g. some forms of Vlasov/Boltzmann-like reductions from general BBGKY hierarchy. These equations are related to the modeling of propagation of intense charged particle beams in high-intensity accelerators and transport systems [1]. In our approach we use fast convergent multiresolution variational-wavelet representations, which allows to consider polynomial and rational type of nonlinearities [2], [3]. The solutions are represented via the multiscale decomposition in nonlinear high-localized eigenmodes, which corresponds to the full multiresolution expansion in all underlying hidden time/space or phase space scales. In contrast with different approaches we don't use perturbation technique or linearization procedures. We consider representation (3) below, where each term corresponds to the contribution from the scale  $i$  in the full underlying multiresolution decomposition as nonlinear multiscale generalization of old  $\delta F$  approach [1]. As a result, fast scalar/parallel modeling demonstrates appearance of high-localized coherent structures (waveletons) and different pattern formation in systems with complex collective behaviour. Let  $M$  be the phase space of ensemble of  $N$  particles ( $\dim M = 6N$ ) with coordinates  $x_i = (q_i, p_i)$ ,  $i = 1, \dots, N$ ,  $q_i = (q_i^1, q_i^2, q_i^3) \in R^3$ ,  $p_i = (p_i^1, p_i^2, p_i^3) \in R^3$  and  $F_N(x_1, \dots, x_N; t)$  be the  $N$ -particle distribution

functions. For  $i = 1, 2$  we have from general BBGKY hierarchy:

$$\begin{aligned} \frac{\partial F_1(x_1; t)}{\partial t} + \frac{p_1}{m} \frac{\partial}{\partial q_1} F_1(x_1; t) \\ = \int dx_2 L_{12} F_2(x_1, x_2; t) \end{aligned} \quad (1)$$

$$\begin{aligned} \frac{\partial F_2(x_1, x_2; t)}{\partial t} + \left( \frac{p_1}{m} \frac{\partial}{\partial q_1} + \frac{p_2}{m} \frac{\partial}{\partial q_2} - L_{12} \right) \\ F_2(x_1, x_2; t) = \int dx_3 (L_{13} + L_{23}) F_3(x_1, x_2, x_3, t) \end{aligned} \quad (2)$$

where partial Liouvillean operators and details are described in [3]. We are interested in the cases when  $F_k(x_1, \dots, x_k; t) = \prod_{i=1}^k F_1(x_i; t) + G_k(x_1, \dots, x_k; t)$  where  $G_k$  are correlators, really have additional reductions as in case of Vlasov-like systems. Then we have in (1), (2) polynomial type of nonlinearities (more exactly, multilinearity). Our goal is the demonstration of advantages of the following representation

$$F_j = \sum_{i \in Z} \delta^i F_j, \quad (3)$$

for the full exact solution for the systems related to equations (1), (2). It is possible to consider (3) as multiscale nonlinear generalization of old  $\delta F$  approach [1]. In (3) each  $\delta^i F_j$  term corresponds to the contribution from the scale  $i$  in the full underlying multiresolution decomposition  $\dots \subset V_{-1} \subset V_0 \subset V_1 \subset V_2 \dots$  of the proper function




Figure 1.  $N = 1$  eigenmode contribution to (3).

space to which  $F_j$  is really belong. It corresponds to the following decompositions:

$$\{F_k(t)\} = \bigoplus_{-\infty < j < \infty} W_j, \quad \overline{\{F_k(t)\}} = V_0 \bigoplus_{j=0}^{\infty} W_j \quad (4)$$

in case when  $V_0$  is the coarsest scale of resolution and where  $V_{j+1} = V_j \oplus W_j$  and bases in scale spaces  $W_i(V_j)$  are generated from base functions  $\psi(\varphi)$  by action of affine group of translations and dilations (the so called “wavelet microscope”). Our constructions are based on variational approach which provides the best possible fast convergence properties in the sense of combined norm  $\|F^{N+1} - F^N\| \leq \varepsilon$  introduced in [3]. Our five basic points after functional space choice are:

1. Ansatz-oriented choice of the (multidimensional) bases related to some polynomial tensor algebra. Some examples related to general BBGKY hierarchy are considered in [3].
2. The choice of proper variational principle. A few projection or Galerkin-like principles for constructing (weak) solutions are considered in [2], [3]. It should be noted advantages of formulations related to biorthogonal (wavelet) decomposition.
3. The choice of bases functions in scale spaces  $W_j$  from wavelet zoo. They correspond to high-localized (non-linear) oscillations/excitations, coherent (non-linear) resonances, etc. Besides fast convergence properties of the corresponding variational-wave-




Figure 2. Stable pattern/waveleton.

let expansions it should be noted minimal complexity of all underlying calculations, especially in case of choice of wavelet packets which minimize Shannon entropy.

4. Operators representations providing maximum sparse representations for arbitrary (pseudo)differential/integral operators  $df/dx$ ,  $d^n f/dx^n$ ,  $\int T(x,y)f(y)dy$ , etc [3].

5. (Multi)linearization. Besides variation approach we consider a different method to deal with (polynomial) nonlinearities, which is based on the para-product structures [3].

So, after application of points 1-5 above, we arrive to explicit numerical-analytical realization of representations (3), (4). Fig.1 demonstrates the first contribution to the full solution (3) while Fig.2 presents (stable) pattern (waveleton) as solution of system (2), (3). We evaluate accuracy of calculations according to norm mentioned above [3].

## REFERENCES

1. R.C. Davidson, Phys. Plasmas, 6, 298, 1999; The Physics of High-Brightness Beams, Ed. J. Rozenzweig, L. Serafini, World Sci., 2001.
2. A.N. Fedorova, M.G. Zeitlin, Quantum Aspects of Beam Physics, 527-538, 539-550, World Scientific, 2002.
3. A.N. Fedorova, M.G. Zeitlin, arXiv preprints: physics/0206049, 0206050, 0206051, 0206052, 0206053, 0206054, 0212066, nlin/0206024.