## Shortcut to a Fermi-Degenerate Gas of Molecules via Cooperative Association

Olavi Dannenberg,<sup>1</sup> Matt Mackie,<sup>1</sup> and Kalle-Antti Suominen<sup>1,2</sup>

<sup>1</sup>Helsinki Institute of Physics, PL 64, FIN-00014 Helsingin yliopisto, Finland

<sup>2</sup>Department of Physics, University of Turku, FIN-20014 Turun yliopisto, Finland

(May 22, 2019)

The creation of a Fermi-degenerate gas molecules using either photoassociation or the Feshbach resonance is theoretically examined. This problem raises an interest because, unlike bosons, fermions in general do not behave cooperatively, so that the collective association of, say, two million atoms into one million molecules is not to be expected. Nevertheless, we find that the coupled Fermi system displays collective Rabi-like oscillations and adiabatic passage between atoms and molecules, thereby mimicking Bose-Einstein statistics. Cooperative association of a degenerate mixture of Bose and Fermi gases could therefore serve as a shortcut to a degenerate gas of Fermi molecules.

Pacs number(s): 03.75.Ss, 05.30.Fk, 34.10.+x

Because of the large number of ro-vibrational degrees of freedom in a molecule, laser cooling is difficult if not impossible to apply in practice, so that a laser-coolingbased approach [1] to a Fermi-degenerate gas of atoms will not work for molecules. Moreover, buffer-gas cooling [2] and Stark-deceleration [3] techniques could take years to reach quantum degeneracy. Here we report that a degenerate gas of Fermi molecules could be formed by applying either photoassociation [4] or a Feshbach resonance [5] to a degenerate mixture of Bose and Fermi atoms [6–8]. The collective Fermi-atom-molecule system is predicted to undergo Rabi-like oscillations, as well as adiabatic passage, from atoms to molecules, thereby mimicking cooperative behavior that was previously attributed to Bose statistics [9,10].

Photoassociation occurs when two atoms absorb a laser photon, thereby jumping from the free two-atom continuum to a bound molecular state. If the initial atoms form a Bose-Einstein condensate (BEC), then a photoassociation laser could conceivably be used to collectively convert a BEC of atoms into a BEC of molecules [11–13]. As it happens, collective free-bound photoassociation is theoretically identical to magnetoassociation, whereby a molecular condensate can be created when one atom from a condensed pair spin flips in the presence of a magnetic field tuned near a Feshbach resonance [14–16]. Intuition developed in one instance is therefore generally applicable to the other, and we will often refer simply to collective association.

In analogy with coherent optical transients in few level atomic systems [17], photoassociation of a BEC has been predicted to facilitate Rabi-like oscillations between atomic and molecular condensates [11–13], whereby an entire gas of, say, two million Bose-condensed atoms are collectively converted into a million molecules that are, in turn, collectively converted back into (roughly) two million atoms, ad infinitum. Another interesting possibility arises because the ground state of the system is all atoms for large positive detunings and all molecules for extremely negative detunings, so that a slow sweep of the laser detuning from one extreme to the other will collectively convert a BEC of atoms into a BEC of molecules [11]. The Feshbach resonance was used in a similar manner to observe [18] collective Ramsey fringes between an atomic condensate and a small fraction of molecular condensate dressed by dissociated atom pairs [19,20].

The statistics of neutral atoms is determined by the number of neutrons in the nucleus, which must be odd for fermionic atoms and even for bosonic atoms. The sum of the total number of neutrons in the nucleus of the constituent atoms similarly determines the statistics of neutral molecules. Accordingly, molecules formed by free-bound association of two fermions will necessarily result in a boson, whereas fermionic molecules are born from the union of a boson and a fermion. Given a degenerate mixture of Bose and Fermi gases [6–8], is it possible that collective free-bound association could serve as a source of Fermi molecules?

To address this question, we model a degenerate Bose-Fermi mixture of atoms [6–8] coupled by either a Feshbach resonance or photoassociation to a Fermi-degenerate gas of molecules. The initial bosonic [fermionic] atoms are denoted by the field  $\phi(\mathbf{r},t)$ [ $\psi_{-}(\mathbf{r},t)$ ], and the fermionic molecules by the field  $\psi_{+}(\mathbf{r},t)$ . Neglecting particle-particle interactions, the Hamiltonian density for such an untrapped system can be written  $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_I$ , where

$$\frac{\mathcal{H}_0}{\hbar} = \phi^{\dagger} \left[ -\frac{\hbar \nabla^2}{2m_B} - \mu_B \right] \phi + \psi_-^{\dagger} \left[ -\frac{\hbar \nabla^2}{2m_-} - \mu_- \right] \psi_- 
+ \psi_+^{\dagger} \left[ -\frac{\hbar \nabla^2}{2m_+} + \delta_0 - \mu_+ \right] \psi_+,$$
(1a)

$$\frac{\mathcal{H}_I}{\hbar} = -\frac{1}{2}\mathcal{K}\left(\psi_+^{\dagger}\psi_-\phi + \phi^{\dagger}\psi_-^{\dagger}\psi_+\right). \tag{1b}$$

Here  $m_B (m_-)$  is the mass of a bosonic (fermionic) atom,  $m_+ = m_B + m_-$  is the mass of a molecule,  $\mu_{B(\pm)}$  is the so-called chemical potential that implicitly accounts for particle trapping,  $\delta_0$  is the binding energy (detuning) of the molecular state relative to the dissociation threshold, and  $\mathcal{K}$  is the (real) atom-molecule coupling.

Without loss of generality, we now make some simplifications. First, the number of atoms, and the traps that contain them, can be adjusted so that  $\mu_B = \mu_- = \mu$ . Second is that atom-molecule conversion occurs on a timescale much faster than the motion of the atoms in the trap, allowing us to neglect the kinetic energies, and justifying our omission of an explicit trapping potential for the particles. Third, since the total number of particles,  $N = \int d\mathbf{r} \left[ |\phi(\mathbf{r})|^2 + |\psi_{-}(\mathbf{r})|^2 + 2|\psi_{+}(\mathbf{r})|^2 \right]$ , is a conserved quantity, we may add the term  $\mu N$  to the Hamiltonian without altering the physics; upon doing so, we also redefine the detuning as  $\delta = \delta_0 - (\mu_+ - 2\mu)$ . Fourth is that, whatever the Fermi energy associated with the fermionic components, it lies within the Wigner threshold regime [20], so that the coupling  $\kappa$  can be taken as the same for all modes. Finally, we switch to momentum space, but retain only the  $\mathbf{k}$  = atomic condensate modes since these transitions are favored over  $\mathbf{k} \neq 0$  modes due to Bose stimulation. The simplified Hamiltonian reads

$$H = \sum_{\mathbf{k}} \left[ \delta c_{\mathbf{k}}^{\dagger} c_{\mathbf{k}} - \frac{1}{2} \kappa (c_{\mathbf{k}}^{\dagger} b_{\mathbf{k}} a + a^{\dagger} b_{\mathbf{k}}^{\dagger} c_{\mathbf{k}}) \right], \qquad (2)$$

where a annihilates a bosonic atom with wavevector  $\mathbf{k} = 0$ ,  $b_{\mathbf{k}}$  ( $c_{\mathbf{k}}$ ) annihilates an atom (molecule) with wavevector  $\mathbf{k}$ , and  $\kappa = \mathcal{K}/\sqrt{V}$ .

The system described by the Hamiltonian (2) is a particular nonlinear version of a two-level atom driven by an external field. Of course, the two-level atom is well known to undergo Rabi flopping [17], and similar behavior observed for the present system would in principle answer question of whether collective association will provide a shortcut to the formation of a Fermi-degenerate gas of molecules. We therefore consider an on-resonance system, for which  $\delta = 0$  and the Heisenberg equations of motion are

$$i\dot{a} = \kappa \sum_{\mathbf{k}} b^{\dagger}_{\mathbf{k}} c_{\mathbf{k}},$$
 (3a)

$$i\dot{b}_{\mathbf{k}} = \kappa a^{\dagger} c_{\mathbf{k}},$$
 (3b)

$$i\dot{c}_{\mathbf{k}} = \kappa b_{\mathbf{k}}a.$$
 (3c)

If the system starts out with  $N_B$  bosonic atoms, the operator a will have a characteristic size  $\sqrt{N_B}$ . On the other hand, while the fermionic operators  $b_{\mathbf{k}}$  and  $c_{\mathbf{k}}$  are characteristically of order unity, the sum over momentum states will ultimately introduce the initial number of fermionic atoms. Hence, for equal numbers of bosons and fermions,  $N_B = N_F = N$ , it is intuitively obvious that the equations of motion (3) yield a system that evolves with the characteristic frequency  $\Omega = \sqrt{N\kappa}$ .

Said intuition is confirmed by solving the timedependent Schrödinger equation numerically, the results of which are shown in Fig. 1. For the simplest case of N = 1, the system undergoes complete conversion from a doubly-degenerate Fermi-Bose system of atoms to a Fermi-degenerate gas of molecules in a time  $\Omega^{-1}$ . However, considering a larger number of initial atoms, N = 5, we see that quantum many-body fluctuations not only frustrate complete conversion, but also adjust the oscillation frequency and lead to a series of collapse and revivals. Increasing the initial particle number to N = 10 brings the amplitude of the first half-oscillation closer to the ideal case. This behavior is exactly analogous to the single-component bosonic case [11,13]. Although limited computational resources preclude the explicit investigation of large particle number, based on the bosonic analogy [11,13] we fully expect the first half oscillation to be complete for large N, i.e.,  $N \gg 1$  initial Fermi atoms should be converted to N Fermi molecules over a timescale  $\tau \sim \Omega^{-1}$ .

Nevertheless, collisions between particles will shift the energy of a state with a given number of atoms and molecules in a manner that depends nonlinearly on the numbers, making it difficult to maintain exact resonance. Furthermore, once a the system has Rabi flopped to molecules, the depopulated atomic condensate makes Bose stimulation, and thus neglect of the  $\mathbf{k} \neq 0$  BEC modes, questionable. And so we investigate the more robust possibility of rapid adiabatic passage [17]. From the Hamiltonian (2), it should be clear that the system will favor all atoms for large positive detunings, while favoring all molecules for large negative detunings. With  $\Omega$ established as the characteristic frequency for collective atom-molecule conversion, then changes in the detuning that are of the order of  $\Omega$ , and occur over a time of order  $\Omega^{-1}$ , should qualify as adiabatic. Hence, if the detuning is swept in a linear fashion according to  $\delta = -\xi \Omega^2 t$ , then dimensionless sweep rates  $\xi \stackrel{<}{\sim} 1$  should enable rapid adiabatic passage to a degenerate gas of Fermi molecules.

Our suspicions are again corroborated by a numerical solution to the Schrödinger equation, shown in Fig. 2. While many-body effects appear to rather seriously affect the efficiency of a marginally adiabatic sweep ( $\xi = 1$ ) compared to the N = 1 case, the difference between N = 5 and N = 10 (not shown) is in fact small. Overall, many-body effects are expected to be weak for near-adiabatic sweeps ( $\xi \sim 1$ ), and vanishing for sweeps that are deep-adiabatic ( $\xi \lesssim 0.1$ ), again in agreement with our BEC experience [11]. On the matter of transitions to noncondensate modes, we note that, in the all-boson case [24], such transitions can be neglected for a sweep directed as in Fig. 2, i.e., for  $\dot{\delta} < 0$ . Given the success of the analogy so far, similar results are expected for a Fermi system.

Before closing, we estimate some explicit numbers. We eschew photoassociation because of the losses associated with the electronically-excited state [25], and focus on the atom-molecule coupling provided by the <sup>87</sup>Rb-<sup>40</sup>K Feshbach resonance located at  $B_0 = 534$  G [21], which has a width  $\Delta_R = 4$  G and an associated zero-field Fermi-Bose-atom scattering length  $a_{aa}^{FB} = -17.8$  nm. Accordingly, the atom-molecule coupling is  $\mathcal{K} \approx (4\pi |a_{aa}^{FB}|\mu_{Bohr}\Delta_R/m_r)^{1/2} = 0.14 \text{ cm}^{3/2} \text{ s}^{-1}$ , where we have estimated the difference between the fermionic-

atom and -molecule magnetic moments to be equal to the Bohr magneton  $\mu_{Bohr}$ . Assuming  $N_B = 10^5$  condensate atoms in a trap with respective radial and axial frequencies  $\omega_r/2\pi = 215 \,\text{Hz}$  and  $\omega_a/2\pi = 16.3 \,\text{Hz}$  [8], the density of bosons is  $\rho_B = 8.1 \times 10^{13} \,\mathrm{cm}^{-3}$ . As for the fermions, we assume a modest number, say,  $N_F = 10^3$ , which two consequences: (i) the atomic BEC will act as a reservoir, thus absorbing any heat created by holes in the Fermi sea [26]: (ii) barring an unfortunately large scattering length for Bose-atom and Fermi-molecule collisions, we can neglect the possibility of any Fermi-Bose collapse instabilities [8]. For  $N_B \gg N_F$ , the timescale for atommolecule conversion is  $\tau_{a2m} \sim (\sqrt{\rho_B} \kappa)^{-1} = 8.2 \times 10^{-7} \text{ s.}$ This timescale is safely below the fastest timescale for trapped-atom motion  $\tau_t = (\omega_r/2\pi)^{-1} = 4.7 \times 10^{-3} \,\mathrm{s},$ justifying our neglect of trap dynamics and the kinetic energy. Similarly, the timescale for Bose atom-atom and Fermi-Bose atom-molecule collisions are roughly equal,  $\tau_c \sim 0.1 \,\mathrm{ms}$ , and are also reasonably neglected.

Lastly, a problem with the Feshbach resonance is noted: the molecules it creates, while translationally and rotationally ultracold, are vibrationally very hot, and are therefore highly susceptible to vibrational relaxation induced by collisions with unconverted atoms (moleculemolecule collisions are Pauli blocked). There is no known handle on such transitions for the present system. Using the sodium numbers [14,16] for comparison, Fermi-Bose [Fermi-Fermi] collisions are estimated to cause vibrational relaxation on a timescale  $\tau_r^{FB} \sim (\sqrt{\rho_B \rho_F} \times 10^{-10})^{-1} \text{ s} = 1 \text{ ms } [\tau_r^{FF} \sim (\rho_F \times 10^{-10})^{-1} \text{ s} = 10 \text{ ms}],$ which is also negligible. If collision-induced vibrational relaxation of the molecules does turn out to be a problem, the trap could be expanded to further reduce the rate of collisions with unconverted atoms.

In conclusion, we highlight that the term  $\Omega = \sqrt{N} \kappa$  was previously referred to as the Bose-enhanced freebound coupling, and the detuning sweep, e.g., referred to as Bose-stimulated rapid adiabatic passage from atoms to molecules. However, this behavior is now clearly independent of statistics, so that Bose stimulation of free-bound association has nothing whatsoever to do with Bose statistics, but is instead a many-body cooperative effect that applies equally well to Fermi-degenerate systems. To four-wave mixing in a Fermi gas of atoms [22,23], we therefore add the above two examples of fermions mimicking an effect that was previously attributed to Bose-Einstein statistics. Such behavior should allow for a shortcut to a degenerate gas of Fermi molecules via photoassociation or the Feshbach resonance.

The authors acknowledge Juha Javanainen for helpful discussions. This work supported by the Academy of Finland (MM and KAS, project 50314), and the Magnus Ehrnrooth foundation (OD).

- [1] B. DeMarco and D. S. Jin, Science 285, 1703 (1999).
- [2] J. D. Weinstein *et al.*, Nature **395**, 148 (1998).
- [3] H. L. Bethlem *et al.*, Nature **406**, 491 (2000).
- [4] J. Weiner, V. S. Bagnato, S. Zilio, and P. S. Julienne, Rev. Mod. Phys. **71**, 1 (1999).
- [5] H. Feshbach, *Theoretical Nuclear Physics* (Wiley, New York, 1992).
- [6] A. G. Truscott et al., Science 291, 2570 (2001).
- [7] F. Schreck *et al.*, Phys. Rev. A **64**, 011402(R) (2001).
- [8] G. Modugno et al., Science 287, 2240 (2002).
- [9] For a discussion of the role of Bose enhancement in photoassociation, see M. Mackie and J. Javanainen, J. Mod. Opt. 47, 2645 (2000).
- [10] For a brief review of cooperative fermion behavior, see J. Javanainen, Nature 412, 689 (2002).
- [11] J. Javanainen and M. Mackie, Phys. Rev. A 59, R3186 (1999).
- [12] D. J. Heinzen, R. Wynar, P. D. Drummond, and K. V. Kheruntsyan, Phys. Rev. Lett. 84, 5029 (2000).
- [13] A. Vardi, V. A. Yurovsky, and J. R. Anglin, Phys. Rev. A 64, 063611 (2001).
- [14] F. A. van Abeelen and B. J. Verhaar, Phys. Rev. Lett. 83, 1550 (1999).
- [15] E. Timmermans, P. Tommasini, R. Côté, M. Hussein, and A. Kerman, Phys. Rev. Lett. 83, 2691 (1999).
- [16] V. A. Yurovsky, A. Ben-Reuven, P. S. Julienne, and C. J. Williams, Phys. Rev. A 60, R765 (1999).
- [17] L. Allen and J. H. Eberly, Optical Resonance and Two-Level Atoms, (Dover, New York, 1987).
- [18] E. A. Donley, N. R. Claussen, S. T. Thompson, and C. E. Weiman, Nature **417**, 529 (2002).
- [19] S. J. J. M. F. Kokkelmans and M. J. Holland, Phys. Rev. Lett. 89, 180401 (2002).
- [20] M. Mackie, K.-A. Suominen, and J. Javanainen Phys. Rev. Lett. 89, 180403 (2002).
- [21] A. Simoni *et al.*, (cond-mat/0301159).
- [22] M. G. Moore and P. Meystre, Phys. Rev. Lett. 86, 4199 (2001).
- [23] W. Ketterle and S. Inouye, Phys. Rev. Lett. 86, 4203 (2001).
- [24] V.A.Yurovsky and A.Ben-Reuven, (cond-mat/0205267);
   M. Mackie *et al.*, (cond-mat/0210131)].
- [25] A two-color photoassociation scheme, whereby electronically excited molecules are transferred to a stable molecule state, would of course avert such losses, and the work herein indicates that "Bose-stimulated" Raman adiabatic passage from atoms to molecules [M. Mackie *et al.*, Phys. Rev. Lett. **84**, 3803 (2000)] will in principle work for fermions.
- [26] E. Timmermans, Phys. Rev. Lett. 87, 240403 (2001).



FIG. 1. Rabi-like oscillations in the fraction of Fermi-Bose atoms converted to Fermi molecules. The initial number of bosonic and fermionic atoms are equal, i.e.,  $N_B = N_F = N$ . (a) The oscillations are complete for an initial number of atoms N = 1, while for N = 5 many-body effects lead to frequency-shifted oscillations that are incomplete and collapse. Better short-time agreement with the N = 1 result is obtained for N = 10. (b) The oscillations eventually revive.



FIG. 2. Rapid adiabatic passage from a Fermi-Bose gas of atoms to Fermi molecules. The detuning is swept as  $\delta(t) = -\xi \Omega^2 t$ , and  $N_B = N_F = N$ . (a) For borderline adiabaticity,  $\xi = 1$ , increasing the number of initial atoms from N = 1 to N = 5 indicates that many-body effects reduce the efficiency. (b) For N = 5 and  $\xi = 0.1$ , near-unit conversion is still possible, despite many-body effects.