# Brussels-Austin Nonequilibrium Statistical Mechanics in the Later Years: Large Poincare System s and Rigged Hilbert Space

Robert C.Bishop

Department of Philosophy, Logic and Scienti c M ethod The London School of Economics, Houghton St. London, W C 2A 2A E, UK

#### A bstract

This second part of a two-part essay discusses recent developments in the Brussels-Austin G roup after the m id 1980s. The fundamental concems are the same as in their similarity transformation approach (see Part I), but the contemporary approach utilizes rigged H ilbert space (whereas the older approach used H ilbert space). While the emphasis on nonequilibrium statisticalmechanics remains the same, the use of similarity transformations shifts to the background. In its place arose an interest in the physical features of large Poincare systems, nonlinear dynamics and the mathematical tools necessary to analyze them.

K eywords: Therm odynam ics, Statistical M echanics, Integrable System s, N onlinear D ynam ics, P robability, A rrow of T in e

W ord count (including notes): 9,547

# 1 Introduction

Part I of this essay discussed the earlier sim ilarity transform ation approach to nonequilibrium statistical mechanics of Ilya Prigogine and his cow orkers. This approach, along with that of subdynam ics, is perhaps som ewhat fam iliar as it has received some attention in philosophical literature and was the subject of Prigogine's well-known book, From Being to Becom ing: Time & Complexity in the Physical Sciences (1980). Part II of this essay focuses on their more recent and less fam iliar work on nonequilibrium statistical mechanics in rigged H ilbert spaces.

It has been argued that no current approaches to microscopic dynam ics can explain or derive the second law of therm odynam ics, since it is both necessary and su cient for the derivation of the second law from microscopic dynam ics that the dynam ics be exact (e.g. Mackey 1992, pp. 98-100; 2002).<sup>1</sup> A lthough it

 $<sup>^1\</sup>text{A}$  dynam ics on a state space ~ with a transfer operator P  $_{t}$  is exact if and only if lim  $_{t!}$   $_1$ 

can be shown that the coarse-grained projection operator arising from the earlier B russels-A ustin approach yields an exact dynam ics, whether their sim ilarity transform ation yields exact dynam ics is unknown (Antoniou and G ustafson 1993; Antoniou, G ustafson and Suchanecki 1998, p. 119). Nevertheless, one of the crucial claim s of the earlier approach was that trajectory descriptions at the m icroscopic level and probabilistic descriptions at the m acroscopic level of therm odynam ic behavior are related via a transform ation (Part I).

This way of view ing the relationship be trajectory and probabilistic descriptions is deem phasized in their more recent work. So the core point is no longer to derive inceversible therm odynam ic behavior from reversible microscopic descriptions, so much as to argue for the priority of inceversible macroscopic descriptions for a particular class of systems known as large Poincare systems. How ever, the core intuitions of the new approach remain continuous with their earlier work; namely, that inceversibility is fundamentally dynamical in character and that distributions are ontologically fundamental explanatory elements for complex statistical system s.

The B russels-A ustin G roup's recent work develops a method for constructing a complete set of eigenvectors for the model equations describing the thermodynamic approach to equilibrium for Large Poincare systems as well as nonlinear dynamics more generally. This approach reformulates the question of how to relate reversible trajectory and irreversible probabilistic descriptions as follows: How can the trajectory dynamics of a large Poincare system (LPS) yield necessary conditions for the thermodynamics approach to equilibrium and what further mechanisms account for the su cient conditions for such behavior?

Large Poincare system s are de ned and illustrated in x2 using nonintegrable H am iltonians and classical perturbation theory as a way of motivating some of the key physical and m athem atical problem s for such system s. The rigged H ilbert space approach to these system s is outlined in x3, and the corresponding tim e-ordering rule and sem igroup operators governing the dynam ics are introduced. Particular details of the approach are discussed in x4, where an alternative interpretation of Prigogine's treatment of trajectories and their relationship to the dynam ics of distributions is developed. Some remarks on probabilistic vs. determ inistic dynam ics closes the essay (x5).

## 2 Large Poincare System s and Integrability

Toward the end of the 19th century, Poincare was investigating planetary motion, am ong other things. Solving the equations of motion for the solar system is extrem ely di cult because all the planets interact with each other through gravitational forces. One of the questions Poincare pursued was whether there was a suitable way to transform these equations of motion into a system of

 $<sup>\</sup>mathcal{P}_t = e_q \dot{\mathbf{j}}_1 = 0$  for every initial density , where  $e_q$  is the unique stationary density (i.e. equilibrium density),  $P_t$  governs the dynam ics (e.g. Liouville or the Frobenius-Perron operators), and the norm is in the sense of Lebesgue integrable functions. Am ong other properties, exact dynam ics are noninvertible and always yield a unique stationary density.

equations where the gravitational interaction would vanish and one could solve the evolution equations for the angle variables of each planet independently of the others. W hat Poincare showed was that in general such a transform ation was impossible for system s of N mutually interacting bodies. If a canonical transform ation for a system of equations describing a set of interacting particles that carries the equations into a set where the interactions vanish exists, then the system is classified as integrable. This means that the original system of equations can be transformed into one where each particle's angle variable is fully described by an equation that is independent of any other particle's angle variable.

Poincare showed that systems of equations were nonintegrable when they contained resonances between various degrees of freedom. In essence a resonance is a transient m etastable state establishing a narrow, precise frequency gateway through which energy can be e ciently transferred from one element of a physical system to another. Physical exam ples of resonances include transient bound states produced in particle collisions and transient interm ediates in chem ical reactions.

#### 2.1 Integrable System s and C lassical Perturbation T heory

In order to m ake these notions of resonances and nonintegrability m ore precise, consider H am iltonian system s in classical m echanics. W hile m odels w ith com - pletely integrable H am iltonians are rare, they are still very useful in the study of physical system s. For m any system s can be m odeled using H am iltonians of the form

$$H = H_0(\mathcal{J}) + \nabla (\mathcal{J}; \sim), \qquad (1)$$

where H<sub>0</sub> is assumed to be completely integrable,  $\mathcal{J}$  represents the action variables (e.g. generalized momentum vectors), ~ the angle variables (e.g. generalized coordinate vectors) and (assumed 1) is the coupling coe cient roughly describing the strength of the interactions through the potential V. The question of whether or not a H am iltonian system is integrable is equivalent to being able to nd a canonical transformation from the old state space coordinates  $(\mathcal{J}; \sim)$  to the new coordinates  $(\mathcal{I}; \sim)$  corresponding to a transformation operator of the form

$$e^{iF(\tilde{I};~)}$$
 (2)

decoupling all the equations for the angle variables (in essence turning o all the interactions by making zero). When such a transformation can be found, the H am iltonian is said to be completely integrable and I will refer to this type of integrability as complete integrability (to be distinguished from the Brussels-Austin sense of integrability below).

In general one then must proceed using a perturbation method where the strategy is to nd approximate solutions of (1) in terms of H $_{\circ}$ (I) plus small perturbations due to V (I;  $\sim$ ). In the course of standard perturbation analysis

of such a model (e.g. Tabor 1989, 89-108), term s of the form

$$\frac{V_{n_{i};n_{j};n_{k}:::}}{n_{i} + n_{j} + n_{k} + \cdots}$$
(3)

em erge where i, j, and k are integers labeling the particles,  $V_{n_1n_jn_k}$  represents the Fourier transform ed potential, the  $n_1$  indicate the (discrete) degrees of freedom of the particles in the Fourier expansion, and the  $_1$  can be negative and are often interpreted as generalized frequencies. C learly terms like (3) increase without bounds when the denom inator approaches zero. The denom inator being zero represents a resonance. It is the presence of a su cient num ber of these resonances that prevents us from using the standard canonical transform ation techniques to turn the model into a completely integrable system of equations. For an N-body problem, the resonance condition takes the form that the - nite sum  $n_{i-1} + n_{j-1} + n_{k-k} + \dots + n_{N-N} = 0$ . In general there are several combinations of  $n_1$ 's satisfying this condition.

#### 2.2 Large Poincare System s

First consider an integrable H am iltonian for a system with two degrees of freedom. The state space trajectories will then be conned to the surfaces of nested tori, where each surface corresponds to a di erent combination of the values of the two constants of the motion. Now add perturbations V to this H am iltonian where 1. If the perturbations leave the H am iltonian integrable, then the model dynam ics are not appreciably a ected. In contrast, if the perturbations render the H am iltonian nonintegrable (e.g. resonance phenomena), then these periodic orbits will be disrupted because such perturbations are as physically important as the unperturbed orbits of the integrable part of the model, due to the transfer of energy involved. The KAM theorem speci es the conditions under which tori associated with quasi-periodic trajectories survive and constitute the majority of motions realized in state space, so that most regions in state space for nonintegrable models close to integrable models show stable nonperiodic orbits (e.g. H ilbom 1994, 337-9).

There are two types of xed points for the state space trajectories in H am iltonians of the form (1): elliptic and hyperbolic (saddle points). Elliptic xed points correspond to stable periodic orbits which are disrupted by resonances. Hyperbolic xed points present complex behavior: trajectories exhibiting sensitive dependence on initial conditions and which wander erratically over large regions of state space. These structures also exhibit self-sim ilarity. The chaotic behavior in H am iltonian system s is sim ilar to chaotic behavior in dissipative system s. However, since H am iltonian system s do not contract to some xed point as do dissipative system s, orbits near hyperbolic xed points will become unstable leading to exponentially diverging trajectories. It should be pointed out that stable and chaotic orbits can coexist sim ultaneously in state space.

Large Poincare system s are of interest to Prigogine and cow orkers. Consider

a typical SM Ham iltonian of the form

$$H (p;q) = \frac{X^{N}}{\sum_{i=1}^{M} \frac{p'_{i}^{2}}{2m_{i}} + \sum_{j>i}^{N} V (jq_{i} q_{j}), \qquad (4)$$

where q and p are N -component vectors representing generalized coordinates and m om enta respectively, and the system is in a large box with volum e L<sup>3</sup>. The B russels-A ustin group is interested in \large" system s, m eaning they work in the lim it L<sup>3</sup>! 1 (the number of particles N m ay be nite or in nite). A LPS is obtained when the system is large and the number of degrees of freedom of the system tends to in nity. An example of a LPS with a nite number of particles would be a nite number of charges interacting with an electrom agnetic eld, while an example with an in nite number of particles would be the therm odynam ic lim it (L<sup>3</sup>! 1, N ! 1, N = L<sup>3</sup> nite). Such system s possess \continuous sets of resonances". By continuous sets of resonances, the B russels-A ustin G roup m eans that in the Fourier transform ed representation, the eigenfrequencies are continuous functions of the wave vector k, so that the sum m ation operations over term s like (3) m ust be replaced by integrals and the denom inators of such term s can be arbitrarily close to zero.

The resonance condition for a continuous set of resonances for a LPS in the context of perturbation theory takes the form

Ζ

$$b dbd = 0, (5)$$

where b (representing degrees of freedom ) and are continuous functions dened over the real num bers. Under condition (5) m otion will not even be quasi-

periodic so that variables have a continuous spectrum <sup>2</sup> No canonical transform ation exists that can turn these LPS m odels into com pletely integrablem odels (P rigogine et al. 1991, pp. 6–7). Such m odels exhibit the type of random ness associated with m ixing, K – ow s and B emoulli system s, but are usually interpreted as determ inistic.<sup>3</sup>

As an example of a LPS, imagine a gas containing an in nite number of particles continually undergoing collisions, where the collision processes never cease. A more realistic example is an electrom agnetic oscillator with frequency  $!_{osc}$  interacting with an electrom agnetic eld. The eld has an in nite number of degrees of freedom and the frequency  $!_k$  of the eld varies continuously with k, giving rise to an in nite number of resonances. C ontinuous resonances like those

 $<sup>^{2}</sup>$ A s K oopm an and von N eum ann rst pointed out, for dynam ical system s with continuous spectra, the states of m otion corresponding to any set become m ore and m ore spread out into an am orphous everywhere dense chaos. Periodic orbits, and such like, appear only as very special possibilities of negligible probability (K oopm an and von N eum ann 1932, p. 261). This is generally acknow ledged to be the rst reference to the term \chaos" in the context of dynam ics.

<sup>&</sup>lt;sup>3</sup>The Baker's transform ation is a favorite m odel of a determ inistic random system for the Brussels-Austin G roup (Part I). The equations are reversible, determ inistic and conservative, yet the mapping turns out to have the Bernoulli property (random ness of a coin toss).

in LPS are involved in fundam entalphenom ena such as absorption and em ission of light, decay of unstable particles and the scattering of electrom agnetic waves o of uids or other form s of m atter, and are found in both classical mechanics (CM) and quantum mechanics (QM).

The rigged H ilbert space (RHS) approach of the P rigogine school is a m ethod for solving the equations of a LPS (both CM and QM) consisting in constructing a complete set of eigenvalues and eigenvectors for the Liouville operator acting on distribution functions <sup>4</sup> The construction of such eigenvalues and eigenfunctions is what P rigogine and colleagues call the 'generalized problem of integration' (P rigogine et al. 1991, p. 4). To be clear about term inology, nding a transform ation that decouples the H am iltonian in (1) is what is required to show that the system is completely integrable in the sense described earlier. C onstructing the complete set of eigenvalues and eigenvectors for a set of equations derived from (1) is what P rigogine and colleagues refer to as 'integrating' or solving the equations of m otion. A lthough initially m otivated in the context of perturbation theory (as sketched here), the rigged H ilbert space approach is m ore general in nature and applicable to any LPS (e.g. m ost systems in SM, system s involving interacting elds).

# 3 M athem atical D etails of the R igged H ilbert Space A pproach

There are three key elements in the Brussels-Austin method to solving LPS equations. First, they utilize distribution functions to describe the dynamics. Second, they adopt extended spaces such as RHS as a mathematical framework for solving the equations. Third, they introduce an \appropriate" time ordering of the dynamical states of the system.

#### 3.1 The Need for Distributions

W hen solutions of the generalized integration problem sketched at the end of x2.2 exist, they reduce to classical trajectories for most CM systems and to state vectors form ost QM systems. In the context of a LPS, how ever, P rigogine and colleagues argue solutions are not reducible beyond distributions for CM systems. E xam ples include systems in kinetic theory, radiation dam ping and interacting elds. One important feature of such physical contexts is that they are characterized by persistent interactions. A coording to P etrosky and P rigogine, a system 's interactions are persistent if there are no asymptotic states such that the interactions nally cease (1997, pp. 33 and 35). For example in kinetic theory, the m olecules of a gas are in constant interaction with one another because they are undergoing continuous collisions. This physical situation should

<sup>&</sup>lt;sup>4</sup>These distribution functions may be understood in terms of the probability density  $(q_1;q_2;q_3;...;p_1;p_2;p_3:..;t)$  of nding a set of molecules (say) with coordinates  $q_1;q_2;q_3;...;$  and m omenta  $p_1;p_2;p_3:..:$  at time t on the relevant energy surface and are analogous to the m icrocanonical distribution.

be contrasted with the idealized case of a single neutral particle scattering o a xed target. In the latter situation, there is a transitory interaction because the particle undergoes an interaction only in a nite region near the target over a very short time interval, while the particle spends the majority of its life in the so-called asymptotic in and out states free of any interactions with the target. Since interactions never cease for system s with persistent interactions, the model equations typically will not be completely integrable.

The presence of persistent interactions is one of the features giving rise to the continuous set of resonances in a LPS. In a gas containing a large num – ber of particles, these resonances allow for energy to be transferred and leveled throughout the system. Through persistent interactions and the resulting resonances, the particles will bose energy and any ordered patterns are destroyed through di usion (see x4.2 below).

A further consequence is that the physical dynam ics are no longer localized, but are spread throughout the space occupied by the LPS.For the gas exam ple, these nonlocal dynamics will take the form of correlations as described in x4.2below. In addition if the number of particles is large enough, then the degrees of freedom for such a gas of particles will have a continuous spectrum qualifying it as a LPS. This implies that we should expect the dynam ical description of such systems to be in terms of distributions of particles rather than in terms of individual particles, because the e ects of long-range and higher-order correlations due to such interactions become at least as in portant as the trajectory dynam ics. The particles remain coupled to one another through their interactions resulting in collective e ects (x4.2 below). This type of long-range coupling at least in plies that the global or collective dynamics of the system cannot be accurately represented by trajectory dynamics alone (see x4.3 below). As a consequence, Prigogine and colleagues believe we must view irreversibility as a property of a system that emerges at the global level which is not derivable from the trajectory description, meaning that distributions are the natural elements for representing statistical phenom ena rather than trajectories.<sup>5</sup>

#### 3.2 The Need for RHS

A RHS is an extended m athem atical space rst introduced by the Russian m athem atician G el'fand and his collaborators (G el'fand and V ilenkin 1964).<sup>6</sup> B rie y a RHS can be understood in the following way. Let be an abstract linear scalar product space and complete it with respect to two topologies. The rst topology is the standard H ilbert space (HS) topology  $_{\rm H}$ 

$$hj = \frac{p(h;h)}{(h;h)},$$
(6)

 $<sup>^{5}</sup>$ To avoid a simple confusion (e.g. B ricm ont 1995, pp. 165-6), note that singular distributions such as delta functions are not used to represent probability distributions in the rigged H ilbert space approach.

<sup>&</sup>lt;sup>6</sup> In more recent work Petrosky and Prigogine (1997) have explored rigging \Liouville space" (the space of density functions or density operators (for dynam ics. Ordonez (1998) has demonstrated that these Liouville spaces can be rigged as a Gel'fand triplet, yielding sem i-group operators and generalized eigenvectors.

where h is an element of resulting a HSH. The second topology is dened by a countable set of norm s

$$j_n j = p_{(; ; )_n}, n = 0; 1; 2:...$$
 (7)

where is also an element of and the scalar product in (7) is given by

I

$$(; {}^{0})_{n} = (; (+1)^{n} {}^{0}); n = 0; 1; 2:::$$
 (8)

and

where is the Nelson operator =  $X_i^2$  (Nelson 1959, 587). The i are the generators of an enveloping algebra of observables for the system in question and they form a basis for a Lie algebra (Nelson 1959; Bohm et al. 1999). For example if we are modeling the harm onic oscillator, the  $X_i$  could be the raising and low ering operators (Bohm 1978, 7-9). Furtherm ore if the operator + 1 is a nuclear operator then this ensures that is a nuclear space (Treves 1967, 509-34; Bohm 1967, 276-7). An operator A is nuclear if it is linear, essentially self-ad joint (Rom an 1975, pp. 540-3) and its inverse is H ibert-Schm idt. The operator A<sup>-1</sup> is H ibert Schm idt if A<sup>-1</sup> =  $X_iP_i$ , where the  $P_i$  are mutually orthogonal projection operators on a nite dimensional vector space and  $a_i^2 < 1$ ,  $a_i$  denoting the eigenvalues of A<sup>-1</sup> (Bohm 1967, 273-6). Notice that the norm (6) is a special case of (7) where n = 0.7

We obtain a Gelfand triplet if we complete with respect to to obtain and with respect to  $_{\rm H}$  to obtain H. In addition we consider the dual spaces of continuous linear functionals and H respectively. Since H is self dual, we obtain

where is characterized by the induced topology . The meaning of the symbol in relation (10) is that every space to the left of a is a subspace of every space to the right of and every space to the left of is dense in the space to the right of with respect to the topology of the space to the right of (see G el'fand and V ilenkin 1964 for more details).

For the B russels-A ustin G roup, the chief reason to work in a RHS is the ability to naturally model unstable physical phenomena such as decay, scattering and the irreversible approach to equilibrium which is lacking in HS (e.g., B ishop 2003a). These kinds of time-dependent processes require complex eigenvalues

<sup>&</sup>lt;sup>7</sup> There are m any di erent inequivalent irreducible representations of an enveloping algebra of a group characterizing a physical system (e.g. the rotation group has an inequivalent irreducible representation for each value of j). They can be combined in m any ways by taking direct products describing combinations of physical system s. These representations are characterized by the values of the invariant or C asim ir operators of the group. So although the N elson operator fully determ ines the topology of , there is freedom in choosing the enveloping algebra describing elementary physical system s. Further restrictions on the choice of function space for a realization of are due to the particular characteristics of the physical system being modeled. This is analogous to the situation for W -algebras in the algebraic approach to Q M (P rim as 1981 pp. 161-249; A m ann and A tm anspacher 1999).

and generalized eigenfunctions (G el'fand and Shilov 1967). Such m athem atical quantities are not well-de ned in a HS, but are given rigorous justi cation in a suitable RHS. In particular the Liouville operator, which characterizes a LPS's approach to equilibrium, does not have a complete set of eigenvalues and eigenfunctions in a HS.Recently the Brussels-Austin Group has dem onstrated that a com plete set of eigenvalues and eigenvectors for this in portant operator can be de ned and calculated for several chaotic models in extended spaces (Antoniou and Tasaki 1992 and 1993; Hasegawa and Shapir 1992; Hasegawa and Driebe 1993). An additional motivation for switching to a RHS is that the equations of motion de ned on a HS are time-symmetric. Time-asymmetric equations may be de ned and solved in a RHS making the latter type of space a natural choice for modeling intrinsic irreversible processes (irreversibility without explicit reference to an environm ent; see Part I). Intrinsic irreversibility is of prine interest to the Brussels-Austin G roup because these types of irreversible processes are related to intrinsic arrows of time in physics (i.e. arrows of time which are independent of hum an intervention or approximation).

#### 3.3 Sem igroup Operators in RHS and Irreversibility

O ne of the important features of RHS is that evolution operators are offen elements of sem igroups rather than groups, so that inversible behavior can be appropriately modeled. The case of simple scattering is a good example for illustrating the concepts. There is a preparation apparatus which prepares particles in a particular state (energy, angular momentum, etc.). The particles are emitted at a target (assumed to be xed in this analysis). The free particle Ham iltonian in (1) is H<sub>0</sub> while the potential in the interaction region surrounding the scattering center is given by V. A fler the interaction with the target, the detector registers the particle measuring quantities such as the angle of scattering relative to the initial direction of the particle as emitted from the accelerator or the energy of the particle after the scattering event.

Each interaction involves a resonance which can be described as

$$E > = 1 + \frac{1}{E + i''} V E > ,$$
 (11)

a Lippm ann-Schwinger-type equation for the evolution of the energy eigenstates as they pass through the scattering region. W henever the operator on the right hand side of (11) applied to the energy eigenstate  $\frac{1}{2}$  > goes to in nity, we have a resonance. A coording to the Brussels-Austin G roup, if, given a su ciently large number of interacting particles, the number of resonances in a system is su ciently large, then the system will evolve from a highly ordered state to a completely random ized or equilibrium state. This evolution is intrinsically irreversible, due to the internal dynam ics of the system .

The intrinsic irreversibility of LPS m odels must be described by sem igroups. This necessitates leaving the HS fram ework and working in a broader mathematical space such as a RHS which Antoniou and Prigogine (1993) adopt in their analysis of the Friedrich's model for scattering. In the Gel'fand triplet H , is the space of particle distribution functions. Furtherm ore Antoniou and Prigogine adopt the following time ordering condition: any excitations or preparations are to be interpreted as events taking place before t = 0 while any de-excitations or detections are to be interpreted as events taking place after t = 0 (1993, pp. 445 and 455).

At the point in the analysis of the scattering experiment where choices have to be made regarding how to interpret the directions of integration for the analytic functions involved in the upper and lower complex half-planes, they choose the following interpretations (1993, pp. 454-5): excitations are identied as taking place before t = 0 (taken to be represented as extensions from the lower to the upper half-plane), while de-excitations are identied as taking place after t = 0 (taken to be represented as extensions from the lower half-plane). So the time-ordering rule is applied with respect to the choice of how to deform the contours in the complex plane with respect to the choice of direction of integration along the contours. Proceeding in this fashion Antoniou and Prigogine derive concrete realizations for the space involving H ardy class function spaces (1993 pp. 457-9; see also B ishop 2003a and 2003b).

Antoniou and Prigogine discuss two sem igroups of evolution operators. The rst is  $U^{y}(t) = e^{iH t}$ , initially de ned on H for 1 < t < 1, extended to It is continuous and com plete in the topology of , valid fort 0 and they identify its temporal direction as carrying states into the forward direction of time. This operator describes evolution reaching equilibrium in the future. The second operator is U<sup>Y</sup> (t) extended to , continuous and com plete in the topol-, but valid for t  $0^8$ . They identify the tem poral direction of this latter oqy operator as carrying states into the backward direction of time ( tincreasing), so this operator describes evolution reaching equilibrium in the past. Since no physical system s are ever observed evolving to equilibrium from the future into the past, they select  $U^{y}(t)$  extended to for t 0 as the physically relevant sem igroup of evolution operators for modeling statistical mechanical system s. This selection is taken to be an expression of the second law of therm odynam ics based on our em pirical observations (Antoniou and Prigogine 1993, p. 461).

The approach sketched in this section for the case of transient scattering can be extended to the case where the interactions are continuous and persistent, yielding sim ilar results (Petrosky and Prigogine 1996 and 1997).

# 4 Discussion of the RHS Approach

The Brussels-Austin Group's RHS approach has yielded solutions (mostly numerical) to nonequilibrium statistical mechanical system equations. Based on these solutions and the insights gained from the new approach, Prigogine and cow orkers make a number of important claims needing detailed discussion.

 $<sup>^8\,{\</sup>rm T}$  he requirem ents of continuity and com pleteness force the unitary group extended to to be restricted to the separate time ranges t ~0 and t ~0 (Bohm and Gadella 1989, pp. 35-119).

#### 4.1 Therm odynam ic Arrow of Time

One of the claim ed virtues of the approach is that it provides an explanation for the therm odynam ic arrow of time (the law of increasing entropy de ned entropy close to equilibrium). This has been one of the central goals of Prigogine since he began his work in SM. One feature that both the earlier similarity transform ation approach (discussed in Part I) and the RHS approach share in this quest is a kind of vacillation between seeking an explanation of the therm odynam ic arrow in the dynam ics of the physical system, and taking the empirically observed direction of the arrow as a fundam ental principle.

In the RHS approach, the types of mechanism s to which the Brussels-Austin G roup appeals for explaining the therm odynam ic arrow are di usion, the grow th of correlations and collective e ects, all of which are generated by Poincare resonances (Antoniou and Prigogine 1993; Petrosky and Prigogine 1996 and 1997). The extension of the description of a LPS with their Poincare resonances, persistent interactions and chaotic dynam ics to G el'fand triplet spaces allow s the eigenvector equations to be solved. In the course of analyzing these solutions, characteristically there are two sem igroups that emerge as sketched in x3.3. At this point in the analysis, one sem igroup is selected because it represents system s approaching equilibrium in the tem poral direction of the future, while the other sem igroup is disregarded because it describes system s approaching equilibrium in the temporal direction of the past which is never observed and, therefore, deem ed to be unphysical (Antoniou and Prigogine 1993, p. 461; Petrosky and Prigogine 1996, p. 453 and 1997, p. 13). By making this latter appeal to observations, the B russels-A ustin G roup is appealing to the very facts they seek to explain via the dynam ics of the physical system .

The model equations alone do not uniquely determ ine which sem igroup is the appropriate one, so some kind of appeal to physical considerations is needed. A s discussed in x3.3 above, the B nussels-A ustin G roup does make an appeal to a criterion for choosing a tem poral ordering: any excitations are to be interpreted as events taking place before t = 0 while any de-excitations are to be interpreted as events taking place after t = 0. W hile there is a clear ordering of time from excitation to de-excitation, the criterion invoked still ultimately rests upon our observations that a system is excited before it undergoes de-excitation. The physical reason why the therm odynam ic arrow runs from the past tow ard the future is still undiscovered in the RHS approach, though the approach gives us the mathematical tools to explore and describe the arrow precisely.

#### 4.2 Correlation D ynam ics

The RHS approach highlights the role of nonlocal and collective e ects due to long-range correlations that introduce new dynamics in the probabilistic description that are typically ignored in the trajectory description of a LPS.<sup>9</sup> The

 $<sup>^9{\</sup>rm P}\,{\rm rigogine}$  (1962, 138-95) introduced a simplied version of correlation dynam ics and G eorge (1973a) developed the idea in the direction indicated in this section.

term \collective e ects" is used to describe the behavior of an aggregate of particles coupled together in some fashion that is distinct from the behavior of individual particles. Collective e ects can arise from long-range forces such as electrom agnetism, gravity or from spatial correlations caused by interactions.

Spatial correlations play an important role in the tem poral ordering of the dynam ics of SM system s. In atom ic orm olecular gases, collective e ects are due to collisions. Consider the idealized textbook situation, where we start with an isolated gas of N particles in a volum e V that have yet to interact with one another. If the initial distribution of the particles is hom ogeneous and isotropic, then the particles are equally likely to be at any point r in V<sup>10</sup>. This result holds for each individual particle under the condition that the positions of the other particles are arbitrary. In a typical gas or liquid, this latter condition is not ful led in general, however. Consider two particles at a time in our gas. G iven the position of one particle, di erent positions of the second particle are not equally likely to obtain; namely, the second particle cannot occupy the position of the rst particle. Due to interparticle interactions and the symmetry properties of the state vectors, di erent values of the relative position  $(\mathbf{r}_2$ 瑫) between our two test particles in the entire gas do not appear with equal likelihood. This feature is known as a spatial correlation between the simultaneous positions  $\mathbf{r}_1$  and  $\mathbf{r}_2$  of the two particles.

In a plasm a, for exam ple, where the gas is composed of charged particles, spatial correlations are the tendencies of unlike charges to cluster together and the tendencies of like charges to repel each other. The simultaneous positions of the particles in the plasm a are not all equally likely. It turns out that there is a sim ple relationship between the spatial integral of the correlation function representing spatial correlation and the mean square uctuation of the density of the gas particles (P athria 1972, 447-50), meaning the spatial distribution of the particles is in uenced by the presence of such correlations. In addition these correlations are directly dependent on the density of particles in the gas. A s the density decreases, such collective e ects disappear because the mean free path of the particle, a measure of the likelihood of a collision during a given distance traveled, become so comparable to V. This means collision events will be very rare and correlations will be kept to a minimum when the mean free path is large.

C ollisions are frequent in dense gases and the spatial correlations induced by collisions couple each particle with m any other particles in the gas. It is this coupling due to correlations that leads to collective behavior responsible for gas particles being collected into coherent structures rather than being uniform ly spread throughout the volum e. E xam ples would be turbulence and shock waves.

To see how these correlations develop, start with the particles in the gas before they have interacted with each other. As they begin colliding, the rst interactions set up binary correlations between particles. As the interactions persist, temary correlations begin to appear. The process will continue by es-

 $<sup>^{10}\,\</sup>text{O}$  f course, in this idealized example the assumption of equiprobability of states is reasonable. In a LPS, by contrast, interactions are persistent, so this assumption cannot be made.

tablishing quaternary correlations and so on through N -ary correlations as more and more particles become involved others through collisions. The progression from lower order correlations (which appear rst) to higher order correlations (which appear later) corresponds to a natural tem poral ordering for the evolution of the states of the gas. C orrelations and other collective e ects can rival or exceed the role of individual particle trajectories and be masked by a dynam ical description that treats trajectories as its basic explanatory element.

For example the electrom agnetic force is a long-range force. It is the dominant force in many situations in a plasma, so the behavior of a plasma is not reducible to the dynam ics of the trajectories of the individual particles alone. In the case of a plasma, the energy of the plasma is a ected by the presence of correlations, such that one of the di erences between the energy of a plasm a and that of an ideal gas (non interacting particles) is given by a correction term due to correlation e ects (K rall and Trivelpiece 1986, 63-5). Not only do these e ects interact with the electrom agnetic elds of the plasma itself, but they also generate new electrom agnetic elds that react back on the plasma leading to very com plex dynam ics.

A m ong the physical mechanisms playing a role in LPS, correlations appear to play a crucial role in irreversibility. A s was apparent in the earlier sim ilarity transform ation approach, the progression of correlations suggests a natural direction for the therm odynam ic arrow (G eorge 1973a). But this is not sim ply another way of saying that entropy increases for such systems because in an open system the order of correlations may continue to grow while the measure of disorder in the system may remain constant or decrease. So correlations are not the complete explanation for the therm odynam ic arrow of tim e.

Long-range correlations are another e ect in the dynam ics of correlations that become apparent in RHS (discussed in its earliest form in Prigogine 1962 and George 1973a). As gas particles begin to interact, correlations develop am ong the particles due to interactions (recall that in a LPS these interactions are associated with resonances). A long with the growing order of correlations, long-range correlations develop as particles interact with one another and then separate over long distances while carrying the \m em ory" of their prior interactions (correlations) with them to other parts of the gas. Over short time scales, the growing order of correlations appears to be the m ore dom inant of the two e ects. As tim e goes on, the long-range correlations due to resonances are built up so that collective e ects become in uential. These long-range correlations are associated with nonequilibrium m odes of energy transfer (P etrosky and P rigogine 1996, p. 468).

O ver longer tim e-scales, another very interesting phenom enon occurs. E quilibrium short-range binary correlations rem ain nite, but nonzero around each particle. In turn temary nonequilibrium correlations are built up am ong particles in a sm all region. These correlations di use throughout the system, leaving the equilibrium correlations, while quartinary nonequilibrium correlations are built up am ong the local particles. These correlations di use throughout the system while quintinary nonequilibrium correlations build up and so forth. A s tim e continues the variously ordered nonequilibrium correlations can propagate over large distances due to di usion so that the corresponding inform ation is transferred globally among the particles of the gas. The end result is a \sea" of multiple incoherent correlations (Petrosky and Prigogine 1996, p. 468). This effect provides a natural tem poral direction for the ow of entropy and is revealed in the types of com plex spectral representations of the statistical evolution operators m ade possible by working in an RHS.

In this sense one m ight argue that as the order of correlations increases, as long-range correlations grow and as higher-order nonequilibrium correlations propagate throughout the gas, the e ects of individual trajectories on the global dynam ics of the gas become eless in portant relative to the elects of the dynam ics of correlations. This does not mean that particles lack trajectories and positions in state space as these types of interaction events are parasitic on these concepts (e.g. mean free path between collisions). In my view correlations and collective elects make the signi cant contributions to the global dynam ics while the elects of trajectories play a role only locally (see below).<sup>11</sup>

O nem ight object that the dynam ics of correlations can somehow be reversed even though the probability of the right kinds of reversals to run the whole evolution backwards (like a lm in reverse) is extremely small. If true, then the situation is still the same as in standard thermodynamics where the increase in entropy in systems is viewed as being reversible though the probability is vanishingly small.

The B russels-A ustin response to such an objection for open system shas been given in x4 of Part I. For closed systems they have shown that as the dynam ics of correlations continue, an \entropy barrier" against inversion develops. This barrier can be de ned as the value of the H -function {a therm odynam ic function related to the entropy, which does not require coarse graining or the invocation of an environm ent in the B russels-A ustin approach {after such an inversion m inus its value before such an inversion. This di erence increases exponentially with time, so the longer the LPS evolves, the higher the barrier to inversion. E sentially this means that the energy requirem ents to invert the system of particles increases very rapidly with time. As the model approaches equilibrium, this energy barrier diverges, hence, there is no physical way of \going back" in the anti-therm odynam ic direction (P etrosky and P rigogine 1996, pp. 468-9 and 494-5).

#### 4.3 \C ollapse of Trajectories"

In the sim ilarity transform ation approach (Part I), Prigogine and collaborators put forward several arguments to the e ect that smooth (i.e., everywhere di erentiable), determ inistic trajectories do not exist for unstable statistical mechanical systems. These arguments were fundamentally awed in similar ways in that epistem ological claims were treated as ontological claims. In the new

 $<sup>^{11}\</sup>text{O}$  f course I have used idealized examples in this section in the sense that we imagined starting with a gas of noninteracting particles and then \turning on" the interactions. Recall that interactions are persistent in a LPS so there is never a time in such system s when the m icroscopic dynam ics can be characterized by sm ooth, sm ooth trajectories.

approach, this bias against such sm ooth trajectories and the dynam ics derivable from trajectories resurfaces in a di erent form that clari es the B russels-A ustin attitude tow and trajectories.

It is well known that in the traditional description, the trajectory of a point particle free of any external forces can be represented m athem atically as a superposition of \plane waves" by taking the position of the particle and applying a Fourier transform from (q;p) space to (k;p) space. In this latter space, a trajectory is a coherent superposition of plane waves and this superposition is modeled by a D irac delta function. For a particle undergoing free m otion, this distribution function is a solution to the equation of m otion, has unchanging width and is everywhere di erentiable throughout its determ inistic evolution (\sm ooth" trajectory).

For a nite number of particles with normalizable distributions, the trajectory description in (k;p) space and the B russels-A ustin probabilistic description agree.<sup>12</sup> In the therm odynam ic lim it, how ever, P rigogine and cow orkers argue that resonances destroy sm ooth trajectories in the follow ing way. In the therm odynam ic lim it, the D irac delta function describing the trajectories of particles at t = 0, once evolution begins, im mediately begins spreading throughout a subspace of (k;p) space under the action of resonances, though maintaining a delta function singularity<sup>13</sup> (Petrosky and Prigogine 1996, pp. 479-481 and 1997, pp. 35-37). The trajectories are no longer representable as delta functions, but by broader kinds of distribution functions. Petrosky and Prigogine unfortunately described this phenom enon as the \collapse of trajectories", but all they really mean is that a di erent notion of trajectory is required in a LPS.

In (q;p) space, this im plies that there are no longer any smooth (everywhere di erentiable) trajectories, but, rather, trajectories exhibiting B rownian motion. A simple way to see this is to return to our idealized gas example. As before, assume initially that the particles have not interacted with each other. Prior to any collisions, the motion of the particles can be characterized by smooth trajectories. As they begin interacting, the particle trajectories become piece-wise continuous as instantaneous discontinuities arise associated with each collision. C ontinuous interactions of this type would then prevent trajectories from being everywhere di erentiable, resulting in particles exhibiting B rownian trajectories rather than smooth ones, but this in now ay im plies that there are no trajectories w hatsoever.

 $<sup>^{12}\,\</sup>rm Som$  e critics, such as B ricm ont (1995, pp. 165 and 175), have overlooked the way in which the RHS approach reduces to standard SM approaches for sm all numbers of particles when LPS conditions are not fullled.

<sup>&</sup>lt;sup>13</sup>The signi cance of the delta function singularity appears to be more mathematical than physical. Mathematically it means that so-called reduced distribution functions{where the distribution function refers to a subset s of the total number of particles in the system {exists in the thermodynamic limit, but such distribution functions almost always exist for molecules under most realistic forces. Reduced distributions were introduced into nonequilibrium contexts by (Brout and Prigogine 1956; Prigogine and Balescu 1959).

Consider the special case of a single sm ooth trajectory represented as

$$(p;q) = \begin{cases} Y^{i} \\ (q q^{0}) (p p^{0}) \\ \vdots \end{cases}$$
(12)

in a LPS model where the superscript 0 indicates the contribution from the unperturbed Ham iltonian. To storder the time evolution of the momentum for the component i = 1 is giving by

$$p_{1}(t) = p_{1}^{0} + \lim_{i=1}^{t} - \sum_{k=n=2}^{X \times X^{N}} (K) \frac{V_{k}}{K \times Y^{0}} = e^{iK \times (\frac{1}{Y} \times v_{n}^{0})t} + 1 e^{iK \times (\frac{1}{Y} \times v_{n}^{0})t}$$
(13)

where is the volume,  $\tilde{x}$  is the wave vector,  $v_1$  is the velocity vector of particle 1,  $v_n$  is the velocity vector of particle n, and " is an in nitesim alpositive constant. The rst term represents the contribution from the unperturbed H am iltonian and the second term represents contributions from the interactions. If N is nite, (13) becomes

$$p_{1}(t) = p_{1}^{0} + \frac{\chi^{N-L}}{n-2} d\tilde{k} \frac{V_{k}}{\tilde{k}} \frac{V_{k}}{\tilde{k}} \chi^{0} + O(2^{0}), \qquad (14)$$

in the limit t ! 1 because the pole at  $\tilde{k}$  (p + q) = i'' vanishes as ! 1, the LPS condition. A coording to (14) the value of the momentum to rst order asymptotically approaches a constant and the time dependence drops out. Note that in the limit  $\dot{g}_{1}^{0}$  (q, j)! 1, the interactions from particles n remains nite even if such interactions are short-ranged due to resonances, so that longrange correlations are built up. In the therm odynam ic limit, (13) generally diverges and Petrosky and Prigogine conclude that point distributions such as (12) representing trajectories are not physically admissible and, therefore, sm ooth trajectories are inconsistent with the therm odynam ic limit in a LPS (1996, p. 480). Only singular nonlocal distributions appear to be consistent with the therm odynam ic limit and such distributions lie outside of HS (Petrosky and Prigogine 1996, pp. 479-81).

These results are related to the nonlocal nature of the collective e ects of the entire distribution described in x4.2 above. If any arbitrary nite number of particles were selected within the system and treated in isolation, all nonlocal di usion and correlation e ects become negligible and we are left with the standard description and results in terms of trajectories (how ever, these trajectories would not necessarily be everywhere di erentiable).

In more realistic situations, the nonexistence of smooth trajectories leads directly to the Brussels-Austin claim that a LPS exhibits behavior that cannot be derived from trajectory dynamics. Such e ects include the breaking of time symmetry (i.e., the appearance of sem igroups of operators governing the evolution instead of groups), di usion and nonlocal correlations. Prigogine and cow orkers refer to these e ects as \non-New tonian" to emphasize the fact that the trajectory description is inadequate to account for them. The existence of collision operators such as the Fokker-Planck operator is only a necessary condition for inteversibility and other \non-N ewtonian" e ects. Particular types of distributions (nam ely singular distributions) must also be present in order to have su cient conditions for such behavior. The class of singular distribution functions is quite broad and applicable to m any ordinary situations in SM (the canonical distribution is an example; see also Prigogine 1962 and 1997). Petrosky and Prigogine have carried out algebraic and computer m odeling to dem onstrate that the trajectory and distribution descriptions yield di erent results for LPS (e.g. 1993, 1994 and 1996).

Ibelieve the appropriate way to understand this new approach with its \non-New tonian" e ects is to agree with them that distribution descriptions cannot be reduced to point-wise descriptions. However, both descriptions should be viewed as valid within their domains. The trajectory description is valid for local regions of a LPS, where there are relatively few particles, so that trajectory dynam ics is the dom inant feature (the trajectories may be either sm ooth and exact, or exhibit random walks). Interactions take place among particles at this local level and to the extent that we can ignore higher-order and long-range correlations, trajectory and distribution descriptions agree in their account of physical behavior as was noted earlier.

W here my interpretation of the Brussels-Austin work diers from their own is when the conditions for a LPS are met (large num ber of particles, continuous frequencies, etc.). I agree that in exam ining the global evolution of LPS, higherorder correlations and collective e ects due to long-range, persistent interactions are the dom inant features, which are not reducible to trajectory dynam ics alone. Trajectories are not irrelevant, how ever, because such features as correlations and collective e ects presuppose particle positions and trajectories. For example, collective e ects in ordinary gases do not discon m the existence of trajectories, though the elects of correlations can rival or exceed the elects of individual particle trajectories and be masked by a dynamical description that treats trajectories as the sole explanatory element. Note that (14) does not im ply sm ooth point trajectories are im m ediately expunded from a LPS. Physically sm ooth trajectories are converted into random walks due to the persistent interactions and the long-range higher-order correlations that di use throughout the system over time. As described above, resonances, collisions and correlations are closely related to long-range correlations and collective e ects, behavioral features of unstable systems for which the trajectory description alone cannot adequately account. For LPS m odels the whole ism ore than the sum of its parts. Particle trajectories are necessary for global distributions to exist, but are insu cient for determ ining how such global distributions evolve in time. The therm odynam ic paradox m ight be dissolved because (1) the tim e-sym m etric behavior of the trajectory dynamics contributes nothing more to the global evolution of the SM system than the necessary conditions for the existence of such a system and (2) in a LPS trajectories exhibit B rownian motion and correlation dynam ics dom inate the macroscopic dynam ics. Therm odynam ic behavior is, then, an emergent global phenom enon possessing a tem poral direction.

M y interpretation suggests a way to reduce the tension in their view between operationalism with respect to trajectories and realism with respect to distributions (see Part I), where the Brownian trajectories of the system give the necessary conditions for the existence of the distribution , but not su cient conditions for its evolution. In my judgement the new approach the Brussels-Austin G roup has been exploring illum inates some of the underlying physical mechanisms of therm odynam ic behavior. Focusing on the growth and dynam ics of correlations and collective e ects are in portant physical insights which have advanced our understanding of therm odynam ics processes. And by employing extended mathematical structures such as RHS, they have developed powerful tools for describing such processes which will doubtless lead to further insights.

As a last comment, I should point out that this R H S approach does not represent a kind of coarse-graining approach, at least as norm ally understood. Em phasis shifts away from trajectories because they are only a part of the story of the behavior of a LPS (coarse-grained accounts typically assume that trajectory dynam ics is the whole story, but that complete descriptions at the trajectory level are computationally intractable). And, as in the sim ilarity transform ation approach, the RHS approach distinguishes between manifolds of stable and unstable motions (in contrast to typical coarse-grained accounts). Furtherm ore, if the global behavior of a LPS is not only emergent, but also constrains the m otion of individual particles (say by restricting the m odes of energy transfer), then an appropriate m athem atical description should be able to describe this kind of feedback between levels in a system. The RHS approach can describe such feedback e ects, whereas coarse-grained accounts cannot because they deal with only one level of a given system . Finally, whether trajectories that are not everywhere continuous nor everywhere di erentiable are determ inistic or not is an open question in the RHS approach, as I discuss in the next section (coarsegrained accounts typically assum e trajectories are determ inistic, though usually no explicit assumptions are made regarding the trajectories' continuity and differentiablility).

## 5 Possibility Rather than Certainty?

P rigogine's provocatively titled book, The End of Certainty (1997), sum s up one of arguably the most important and far reaching consequences of the Brussels-Austin Group's work: Namely, that the certainty of the determ inistic, timesymmetric trajectory description is not applicable to the global dynamics of a LPS. Instead only a statistical description of probability densities remains. In conventional CM and SM models, particle positions and trajectories are treated as the fundamental ontological entities determining the dynamical evolution of the system. In the Brussels-Austin view this is no longer the case for LPS models. The fundamental ontological feature for these models are the probability distributions, i.e., the large-scale arrangements of the particles them selves. To reform ulate the laws of classical dynamics along the statistical lines suggested by Prigogine and co-w orkers leads to the conclusion that such laws now express \possibilities" and no m ore \certainties"' (Petrosky and Prigogine 1997, p. 1).

W here there are relatively few numbers of particles, the B nussels-A ustin G roup's approach to dynam ics reduces to the standard results of CM, so the trajectory picture with its determ inistic and tim e-reversible character is preserved as a limiting case. In non-LPS cases, the RHS approach recovers the usual results of SM (e.g. Fokker-P lanck equation, Boltzm ann equation, non-M arkovian m aster equations). It is in cases where the LPS criteria apply that probability become es the fundam ental notion, irreducible to the trajectory description. System s m ust be treated as wholes. If any subset of the total num ber of particles N is treated by itself all the \non-N ew tonian" e ects disappear and the conventional descriptions are recovered. It is in this sense that P rigogine believes, W hat is now emerging is an \interm ediate" description that lies som ew here between the two alienating in ages of a determ inistic world and an arbitrary world of pure chance...[T] he new laws of nature dealw ith the possibility of events, but do not reduce these events to deductible, predictable consequences' (P rigogine 1997, p. 189).

The nature of this possibility supposedly represents a new conception which remains to be claried, however. It is clearly not the kind of irreducible indeterminism described in von Neumann collapse, where some sort of collapse from multiple possibilities to a single actuality is envisioned. As Prigogine and colleagues describe it, their probabilistic form ulation of physics is also to be distinguished from the type of chaotic dynamics, where the underlying dynamics is deterministic, but the outcomes of the system are not predictable. The latter is epistemically indeterminable but not ontically indeterministic.<sup>14</sup> Instead the dynamics envisioned by Prigogine and his colleagues involve an interplay between unitary reversible processes and irreversible processes. The LPS are important examples of dynamical systems which show this kind of interplay and are, therefore, intrinsically probabilistic.

But the relationship of this probabilistic evolution to determ inistic dynam ics remains unclear and requires attention because under some conditions the dynam ics of probability distributions can be \embedded" into completely determ inistic dynam ics and M arkov processes can almost always be \embedded" into determ inistic K olm ogorov processes (Antoniou and G ustafson 1993; G ustafson 1997, pp. 55-76). This leaves open the possibility that there is no signi cant fundamental di erence between this new conception of probabilistic evolution and the conventional conception of determ inistic evolution, or so one could plausibly arque.<sup>15</sup>

Though more needs to be said regarding the notion of probabilistic dynamics

 $<sup>^{14}</sup>$ Understanding what it means for a system or a description to be ontically indeterm inistic is by no means straightforward (e.g. B ishop 2002).

<sup>&</sup>lt;sup>15</sup> I should point out that although there m ay exist theorem s show ing that given any M arkov process, that process can be embedded in a larger determ inistic K olm ogorov process, the general result does not necessarily m ean that the given M arkov process is determ inistic. W hether or not a given M arkov process is determ inistic or not is an ontological rather than a m athem atical question. It should also be clear, how ever, that sim ply characterizing the probability densities via K olm ogorov m easures is insu cient because this cannot settle the ontological nature of the probability.

they are working out, it must be internally generated by the dynam ics of the system (e.g. via correlation dynam ics) rather than im posed from the outside by observers, m easuring apparatuses or the environm ent. I do not take it that this need for m ore clari cation is a serious weakness of their program. On the contrary, I look at the situation as analogous to the early days of quantum theory where m any concepts (indeterm inacy being one of them) were very hazy at the start inviting serious re ection and exploration.

The RHS form alism gives us a uni ed description of dynam ics and them odynam ics within a statistical fram ew ork and a consistent, rigorous description of irreversible processes. The m athem atical developm ents are indeed in pressive, including new results regarding the theory of com plex spectral representations of operators. Furtherm ore this fram ew ork is pow erful enough to allow a unication between CM and QM (Prigogine et al. 1991; Petrosky, Prigogine and Tasaki1991; Petrosky and Prigogine 1994). How ever, the prom ise of the recent Brussels-Austin work must be balanced against two im portant open questions: (1) W hat is the physical and m athem atical status of the past-directed t 0 sem igroup (x3.3) and (2) W hat is the precise nature of the probability lying at the heart of an LPS? A nsw ering these two questions holds the key to their being able to o er an explanation for the therm odynam ic arrow of tim e and for their developing a notion of indeterm inism that is di erent in kind from that discussed in conventionalQM developm ents that would be truly revolutionary.

As things stand, the Brussels-Austin G roup has given us a powerful descriptive tool for irreversible processes, and nonlinear dynam ics m ore generally, but they have not given us an explanation for the origination of the irreversibility we observe in our world. O nem ight object that the RHS approach is ultimately only of mathematical interest since there is nothing philosophically interesting given the current state of the above open questions. This response is too quick, how ever. These open questions can also be viewed as opportunities for exploration of the underlying concepts of the approach in order to attempt to answer these questions. For example, by adopting a di erent arrow of time in the context of scattering in a RHS formulation of QM , one can show that the 0 sem igroup is also future oriented (this time arrow is, however, highly t operational in character and not generally applicable outside of laboratory contexts; for discussion, see B ishop 2003a and 2003b). So interesting conceptual questions are raised by the Brussels-Austin work. Besides, even if questions (1) and (2) should ultim ately be answered in a way that closes o this avenue for nonequilibrium SM, that information is also valuable to philosophers.

A cknow ledge ents {This essay was considerably strengthened from discussions with and comments by Ioannis Antoniou, Harald Atmanspacher, Jerem y Butter eld, Dean Driebe, Fred Kronz, Tom io Petrosky, MichaelRedhead, Michael Silberstein and num erous anonym ous referees. I take full responsibility for remaining weaknesses.

#### References

Amann, A. and Atmanspacher, H. (1999) C - and W - Algebras of Ob-

servables, Their Interpretations, and the Problem of Measurem ent', in H.Atmanspacher, A.Amann and H.Muller-Herold (eds), On Quanta, Mind, and Matter: Hans Primas in Context (Dordrecht, The Netherlands: Kluwer Academ ic Publishers).

Antoniou, I. and Gustafson, K. (1993) From Probabilistic Descriptions to Determ inistic Dynam ics", Physica A 197:153-66.

Antoniou, I., Gustafson, K. and Suchanecki, Z. (1998) From Stochastic Sem igroups to Chaotic D ynam ics', in A.Bohm, H-D.Doebner and P.K ielanowski (eds.), Irreversibility and Causality: Sem igroups and Rigged H ibert Spaces (Berlin: Springer-Verlag).

Antoniou, I. and Prigogine, I. (1993) Intrinsic Irreversibility and Integrability of Dynam ics', Physica A 192:443-464.

Antoniou, I. and Tasaki, S. (1992) Generalized Spectral Decomposition of the -adic Baker's Transform ation and Intrinsic Irreversibility' Physica A 190: 303-329.

\_\_\_\_\_ (1993) SpectralD ecom position of the RenyiM ap', Journal of Physics A 26: 73-94.

Bishop, R. (2002) Determ inistic and Indeterm inistic Descriptions', in H. Atmanspacher and R. Bishop (eds.), Between Chance and Choice: Interdisciplinary Perspectives on Determ inism (Thorverton: Imprint A cadem ic), in press.

\_\_\_\_\_ (2003a) The Arrow of Time in Rigged Hilbert Space Quantum Mechanics', International Journal of Theoretical Physics, in press.

\_\_\_\_\_ (2003b) Q uantum T in e A now s, Sem igroups and T in e-R eversal in Scattering', International Journal of Theoretical Physics, accepted.

Bohm, A. (1967) Rigged Hilbert Space and Mathematical Description of Physical Systems', in W. Brittin, A. Barut and M. Guenin (eds), Lectures in Theoretical Physics Vol IX A: Mathematical Methods of Theoretical Physics (New York: Gordon and Breach Science Publishers, Inc), pp. 255–317.

\_\_\_\_ (1978) The Rigged Hilbert Space and Quantum Mechanics (Berlin: Springer-Verlag).

Bohm, A. and Gadella, M. (1989) Dirac Kets, Gamow Vectors, and Gelfand Triplets, Lecture Notes in Physics, vol. 348 (Berlin: Springer-Verlag).

B raunss, G. (1984) On the Construction of State Spaces for Classical Dynam ical Systems with a Time-dependent Ham iltonian Function', Journal of M athem atical Physics 25, 266-70.

Brian ont, J. (1995) Science of Chaos or Chaos in Science?' Physicalia Magazine 17: 159-208.

Brout, R. and Prigogine, I. (1956) Statistical Mechanics of Irreversible Processes, Part VII: A General Theory of Weakly Coupled System s', Physica 22: 621-36.

G el'fand, I. and Shilov, G. (1967) G eneralized Functions Volum e 3: Theory of D i erential Equations, M einhard E. M ayertr. (New York: A cadem ic Press).

Gel'fand, I. and Vilenkin, N. (1964) Generalized Functions Volume 4: Applications of Harmonic Analysis, Amiel Feinstein tr. (New York: Academic Press).

George, C. (1973a) Subdynam ics and Correlations', Physica 65, 277-302.

(1973) Lectures in Statistical Physics II, Lecture Notes in Physics (Berlin: Springer-Verlag).

Gustafson, K. (1997) Lectures on Computational Fluid Dynamics, Mathematical Physics, and Linear Algebra (Singapore: World Scientic).

Hasegawa, H. and Driebe, D. (1993) Spectral Determ ination and Physical C onditions for a class of chaotic Piecew ise-Linear M aps' Physics Letters A 176: 193-201.

Haægawa, H. and Shapir, W. (1992) Unitarity and Irreversibility in Chaotic System s', Physical Review A 46: 7401-7423.

Hilborn, R. (1994) Chaos and Nonlinear Dynamics: An Introduction for Scientists and Engineers (Oxford: Oxford: Oxford University Press).

K rall, N . and Trivelpiece, A . (1986) Principles of Plasm a Physics (San Francisco: San Francisco Press).

Koopm an, B. and von Neum ann, J. (1932), Dynam ical System s of Continuous Spectra, Proceedings of the National Academ y of Sciences 16: 255-261.

M ackey, M. (1992) T in e's Arrow: The O rigins of Therm odynam ic Behavior (Berlin: Springer-Verlag).

\_\_\_\_\_ (2002) M icroscopic D ynam ics and the Second Law of Thermodynam ics', in C.M ugnai, A.R anfagni and L.Schulm an (eds.), T im e's Arrows, Q uantum M easurem ent and Superlum inal Behavior (R om e: Consiglio N azionale D elle R icerche), pp. 49-65.

O rdonez, A. (1998) R igged H ilbert Spaces A sociated with M isra-P rigogine-C ourbage T heory of Irreversibility', Physica A 252: 362-376.

Pathria, R. (1972) Statistical Mechanics (Oxford: Pergam on Press).

Petrosky, T. and Prigogine, I. (1993) Poincare Resonances and the Limits of Trajectory D ynamics', Proceedings of the National Academ y of Sciences USA 90: 9393-7.

\_\_\_\_\_ (1994) C om plex SpectralR epresentation and T im e-Sym m etry B reaking', Chaos, Solitons & Fractals 4: 311-359.

\_\_\_\_\_ (1996) Poincare Resonances and the Extension of Classical Dynam ics', Chaos, Solitons & Fractals 7:441-497.

(1997) The Extension of Classical Dynam ics for Unstable Ham iltonian System s', Computers & Mathematics with Applications 34:1-44.

Petrosky, T., Prigogine, I. and Tasaki, S. (1991) Quantum Theory of Non-Integrable System s', Physica A 173: 175-242.

Philippot, J. (1961) Initial Conditions in the Theory of Irreversible Processes', Physica 27: 490-6.

Prigogine, I. (1962) Non-Equilibrium Statistical Mechanics (New York: John W iley & Sons).

\_\_\_\_\_ (1980) From Being to Becom ing: Time & Complexity in the Physical Sciences (New York: W.H.Freeman).

\_\_\_\_\_ (1997) The End of Certainty: Time, Chaos, and the New Laws of Nature (New York: The Free Press).

Prigogine, I. and Balescu, R. (1959) Trreversible Processes in Gases I.: The Diagram Technique' Physica 25: 281–301.

P rigogine, I. and Petrosky, T. (1999) LawsofN ature, Probability and Time Symmetry Breaking' in Antoniou and Lumer (ed) Generalized Functions, Operator Theory, and Dynamical Systems (Boca Raton, FL: Chapman & Hall/CRC Press).

Prigogine, I., Petrosky, T., Hasegawa, H And Tasaki, S. (1991) Integrability and Chaos in Classical and Quantum Mechanics', Chaos, Solitons & Fractals 1:3-24.

Prim as, H. (1990) M athem atical and Philosophical Questions in the Theory of Open and M acroscopic Quantum Systems', in A.M iller (ed), Sixty-Two Years of Uncertainty: Historical, Philosophical, and Physical Inquiries into the Foundations of Quantum M echanics (New York: Plenum Press).

Roman, P. (1975) Som e M odern M athem atica for Physicists and O ther O utsiders: An Introduction to A bebra, Topology, and Functional Analysis Volum e 2 (New York: Pergam on Press).

Tabor, M . (1989) Chaos and Integrability in Nonlinear D ynam ics (New York: John W iley & Sons).

Treves, F. (1967) Topological Vector Spaces, D is tributions and K emels (New York: A cadem ic P ress).