## arXiv:physics/0403102v1 [physics.optics] 22 Mar 2004

## Entanglement and inequality of forward and reversed processes as a physical base of nonlinear phenomena in optics.

V.A.Kuz'menko<sup>1</sup>

Troitsk Institute for Innovation and Fusion Research, Troitsk, Moscow region, 142190, Russian Federation.

## Abstract

Connection between the concept of entanglement and origin of nonlinear phenomena in optics is discussed.

PACS number: 03.65.-w, 42.50Gy, 42.50.Hz

The concept of entanglement plays an important role in quantum mechanics and usually it is connected with so-called nonlocality of quantum mechanical interactions in space [1, 2]. Many years it is extensively studied due to fundamental interest and also for possible practical applications in the fields of communication and quantum computation. Substantially less attention is devoted to the so-called *entanglement in time* [3-7].

The concept of entanglement in time can also have other sense as a memory of quantum system about the *initial state*. Does such memory possible? The experiments clearly show that such memory of quantum system about the initial state really exists and manifests itself through inequality of forward and reversed processes in optics [8-10]. Efficiency of the reversed process may be much greater, than the efficiency of forward one.

In the recent experimental works [8, 9] the forward (splitting) and reversed (mixing) processes with a photons were studied (Fig.1b). On the first stage the narrowband (0.04 nm) radiation of nanosecond laser was transformed through down-conversion in the nonlinear crystal into two intense broadband beams (each spectral width ~ 100nm). On the second stage this two broadband beams were mixed in the sum frequency generator [9] or in the process of two photon excitation of rubidium atoms [8]. The mixing of entangled photons leads to regeneration of initial narrowband radiation and is the example of reversed process into the *initial state*. In contrast, the mixing of non-entangled photons should give broadband radiation and is the example of again only forward process. Both experiments show the same result: the efficiency of reversed process is much greater, than the efficiency of forward process. In these experiments the entangled photons demonstrate the *memory* about the *initial state*.

In the next case a polyatomic molecules demonstrate such memory [10]. Excitation and deexcitation of  $SF_6$  molecules by  $CO_2$ -laser radiation was studied in this work (Fig.1a). The forward process (photon absorption) in this case has extremely high spectral width ~ 150 GHz (the so-called wide component of line [11]). In contrast, the reversed process (stimulated emission) has very

<sup>&</sup>lt;sup>1</sup>Electronic address: kuzmenko@triniti.ru

small spectral width. The difference in spectral width of forward and reversed processes exceeds five orders of magnitude. Accordingly, the cross-section of the reversed process turns out to be in several orders of magnitude greater, than the cross-section of forward process [12].

The discussed experiments [8-10] are the direct experimental proofs of inequality of forward and reversed processes. Furthermore, we have enormous quantity of indirect proofs. The nonlinear optics, as a whole, looks like as a large accumulation of indirect evidences of such inequality. Nonlinear phenomena in optics usually are described by mathematical models which are based on the Bloch equations. Such mathematical models usually give good description of nonlinear effects but, in fact, its do not have any clear physical sense. The concepts of coherent states, wavepackets and their interference are used for explanation of physical nature of nonlinear phenomena. At the same time the question about the existence of coherent states is debated in literature [13]. Some theorists argue that the inability to measure the absolute phase of an electromagnetic field prohibits the representation of a lasers output as a quantum optical coherent state [14, 15]. In fact, it means, that using the concept of coherent states for explanation of nonlinear phenomena does not have physical sense.

Alternative explanation of origin of nonlinear phenomena in optics is based on the concept of inequality of forward and reversed processes [16]. It gives really simple and clear interpretation of origin of nonlinear phenomena. A terminal stage of nonlinear processes (there are usually a multiple step photon mixing processes) is returning the system into the initial state. This process proceeds with extremely high efficiency. Because of a photon has spin, it is impossible to return exactly into the initial state with odd number of photons. That is why in the gas and liquid phase the observed nonlinear effects are based only on schemes with even number of photons (four, six and so on photon mixing). In a solid state with rigid lattice the rotational motion of atoms and molecules may be completely suppressed and the role of a photon's spin vanishes. In this case the nonlinear effects with odd number of photons become possible (three photon mixing).

The concept of *initial state* of quantum system should include also the orientation of molecule in space and even the phase of vibrational motion of atoms. Thanks to this fact the experimental study of rotational and vibrational motions of molecules in the real time become possible [17, 18].


So, we need to originate a new mathematical model (alternative to the Bloch equations) for description the dynamics of optical transitions. Such model should be based on the concept of inequality of forward and reversed processes.

In conclusion, we point out that the study of *entanglement in time* does not less important task, than the study of *entanglement in space*.

## References

[1] A.Zeilinger, Rev.Mod.Phys. **71**, S288 (1999).

- [2] C.Brucner, M.Zukowski and A.Zeilinger, E-print, quant-ph/0106119.
- [3] C.Brukner, S.Taylor, S.Cheung and V.Vedral, E-print, quant-ph/0402127.
- [4] A.C.Elitzur, S.Dolev and A.Zeilinger, E-print, quant-ph/0205182.
- [5] C.F.Boyle and R.L.Schafir, E-print, quant-ph/0107098.
- [6] A.Suarez, E-print, quant-ph/ 9801061.
- [7] K.Hess and W.Philipp, E-print, quant-ph/0211117.
- [8] B.Dayan, A.Pe'er, A.A.Friesem and Y.Silberberg, E-print, quant-ph/0401088.
- [9] B.Dayan, A.Pe'er, A.A.Friesem and Y.Silberberg, E-print, quant-ph/0302038.
- [10] C.Liedenbaum, S.Stolte and J.Reuss, Chem. Phys. 122, 443 (1988).
- [11] V.A.Kuz'menko, E-print, physics/0204003.
- [12] V.A.Kuz'menko, E-print, physics/0306148.
- [13] K.Nemoto and S.L.Braunstein, E-print, quant-ph/0312108.
- [14] K.Molmer, Phys.Rev.A 55, 3195 (1997).
- [15] T.Rudolph and B.C.Sanders, Phys.Rev.Lett. 87, 077903 (2001).
- [16] V.A.Kuz'menko, E-print, physics/0401051.
- [17] P.M.Felker, J.Phys.Chem., 96, 7844 (1992).
- [18] A.Zewail, Laser Physics, 5, 417 (1995).



**Fig.1 a)** Principle scheme of forward (absorption) and reversed (stimulated emission) processes in a molecules. **b)** Principle scheme of forward (splitting) and reversed (mixing) processes with a photons. 1- down-converting crystal, 2- sum frequency generation crystal or rubidium cell, 3- spectrometer, 4- photomultiplier.