A regularized inverted perturbation approach method: potential energy curve of the $4^{1}\Sigma_{u}^{+}$ state in Na₂

A. Grochola and P. Kowalczyk^{*}

Institute of Experimental Physics, Warsaw University, ul. Hoża 69, 00-681 Warsaw, Poland

W. Jastrzebski

Institute of Physics, Polish Academy of Sciences, Al.Lotników 32/46, 02-668 Warsaw, Poland

A. Pashov

Institute for Scientific Research in Telecommunications, ul. Hajdushka poliana 8, 1612 Sofia, Bulgaria

(Dated: July 19, 2018)

Abstract

We describe a modification of the inverted perturbation approach method allowing to construct physically sensible potential energy curves for electronic states of diatomic molecules even when some parts of the potential are not adequately characterized by the experimental data. The method is based on a simple regularization procedure, imposing an additional constraint on the constructed potential curve. In the present work it is applied to the double minimum $4^{1}\Sigma_{u}^{+}$ state of Na₂, observed experimentally by polarization labeling spectroscopy technique.

PACS numbers: 31.50.Df, 33.20.Kf, 33.20.Vq

^{*}Corresponding author; Electronic address: pfkowal@fuw.edu.pl

I. INTRODUCTION

Perhaps the most challenging among the electronic states of diatomic molecules are those characterized by exotic shapes of the potential energy curves, which may exhibit barriers towards dissociation, unusual bends or shelves as well as multiple minima. Most of these features reflect interactions between the initially regular diabatic molecular states which, if belonging to the same irreducible representation of the symmetry group of the molecule, in the adiabatic approach cannot cross each other. Except of satisfying a natural, fundamental interest, investigation of such states can bring an immediate, twofold profit. First, the complicated shape of the exotic curves is very sensitive to relative positions of the original diabatic potentials. Therefore experimentally determined curves provide particularly rigorous tests of the quality of theoretical calculations. Second, it happens frequently that exotic potential wells are unusually wide. Such a well can be employed then as an intermediate state to access the long range region of other states situated below or above it.

The unusual electronic states require special methods of analysis and reduction of the experimental data to potential energy curves. In the simplest instance of regular Morselike potentials, the standard Rydberg-Klein-Rees (RKR) algorithm [1], based on the Bohr-Sommerfeld quantization of the phase integral for the vibrational motion, is used. This algorithm can also be generalized to treat some "irregular" curves [2, 3], but evidently fails in the case of double minimum potentials. As an alternative the variational inverted perturbation approach (IPA) has been developed [4], providing fully quantum mechanical determination of potential energy curves directly from the experimental data. In this procedure, one starts with some estimated approximate potential curve and iteratively seeks corrections to it until the quantum mechanical eigenenergies calculated from the improved curve agree with the experimental term values in the least squares approximation (LSA) sense. Several routines have been proposed for realization of this idea, differing mainly in mathematical representation of the potential [5, 6, 7, 8, 9, 10]. In most of them analytical expressions of various forms [5, 7, 8, 9, 10] allow to accurately describe potential energy curves with few fitting parameters, but up to now they have been designed to deal only with "regular" potentials with a Morse-like shape.

The pointwise representation of the potential, proposed within the framework of the IPA method by Pashov *et al.* [6], is more general. As an essentially model-free approach,

it imposes no limitation on the shape of the fitted curve and therefore can be applied to both "regular" and "irregular" potential energy curves [11, 12, 13, 14, 15, 16]. However, the pointwise approach has also disadvantages following from its model-free nature. The method works well for the parts of the potentials covered by abundant experimental data (i.e. energies of rovibrational levels), but when the experimentally determined energy levels become sparse, the inversion problem starts to be highly ill-conditioned and the procedure may become unstable, producing irregularities in a form of unphysical wiggles on the constructed potential curve. In the original algorithm [6] the problem was partially solved by using the singular value decomposition (SVD) technique [17] instead of the standard LSA for fitting the molecular potential. Additional smoothing of the potential resulted from using a cubic spline function for interpolation between the points defining the potential energy curve; increasing the grid spacing could reduce the undesirable flexibility of the constructed out to be nontrivial and even tricky.

We have recently encountered such a problem when investigating the $4^{1}\Sigma_{u}^{+}$ state in Na₂ molecule. To the best of our knowledge this state has escaped experimental characterization up to now. The ${}^{1}\Sigma^{+}$ states in alkali dimers are expected to show an intricate system of avoided crossings between different Rydberg and valence states at short internuclear distances and also ion-pair states (Na⁺Na⁻) at long internuclear distances. In particular, for the $4^{1}\Sigma_{u}^{+}$ state under consideration the theoretical calculations predict a double minimum potential, with a deep inner well and a shallow but broad outer one, in addition to a clearly visible inflection in the right-hand wall of the inner well. Our experimental data related to the $4^{1}\Sigma_{u}^{+}$ state contained 277 energies of levels located in the inner well; they allowed us to find the detailed shape of this well. On the other hand, only 55 levels have been observed in the region above the potential barrier, whereas the outer potential well was totally inaccessible in the present experiment. Therefore determination of the outer well turned out to be an ill-posed problem. We have shown, however, that a physically sensible solution can be found by application of a simple but effective regularization procedure, based on an additional constraint imposed on a constructed potential. As a result, a smooth potential energy curve corresponding to a broad range of internuclear distances (2.4 to 20.0 A) has been generated, reproducing energies of all experimental rovibronic levels to within the experimental accuracy and displaying the expected physical behavior.

In Section II we present the experimental setup and techniques used. Section III contains a brief description of the recorded spectra. The basic concepts of the inverted perturbation approach as well as the proposed modification of the algorithm are discussed in Section IV and their application to deduction of the potential energy curve for the $4^{1}\Sigma_{u}^{+}$ state is shown in Section V. Finally, Section VI summarizes our results.

II. EXPERIMENTAL

To study the $4^{1}\Sigma_{u}^{+} \leftarrow X^{1}\Sigma_{g}^{+}$ system in Na₂, we employed the V-type optical-optical double resonance polarization labeling spectroscopy technique with two independent pump and probe light beams. In our version of the method the probe beam had a fixed frequency and excited a few assigned molecular transitions, whereas the pump beam was tuned over an investigated spectrum. The experimental apparatus and method have been described in detail elsewhere [18, 19] and therefore only the essential features will be presented here. Na₂ molecules were produced in a heatpipe oven, operating at around 750 K, with 4 Torr of helium as a buffer gas. The pump and probe light beams were superimposed collinearly in the sodium vapor zone. As a probe light we employed one of the five blue-green lines of a linearly polarized multimode Ar⁺ laser (Carl Zeiss ILM 120, $\lambda = 476.5, 488.0, 496.5, 501.7$ and 514.5 nm), with a typical power ranging from 20 to 150 mW. The probe laser light excited several known transitions in the $B^1\Pi_u \leftarrow X^1\Sigma_g^+$ system of sodium dimer [20], thus labeling the involved rovibrational levels in the ground electronic state. A parametric oscillator/amplifier system (Sunlite EX, Continuum) provided with a frequency doubler (FX-1) and pumped with the third harmonic of an injection seeded Nd:YAG laser (Powerlite 8000) served as a source of the pump light. The system produced UV radiation with a typical energy 3 mJ per pulse and a spectral width of 0.16 cm^{-1} . The tunable pump light excited sodium molecules from the ground $X^1\Sigma_g^+$ state to the $4^1\Sigma_u^+$ state studied in this experiment. The frequency of the pump beam was calibrated against the optogalvanic spectrum of argon in a hollow cathode discharge tube and the transmission fringes of a Fabry-Pérot interferometer 0.5 cm long. The uncertainty in determining the line centers for strong molecular lines is estimated to 0.05 cm^{-1} .

Crossed polarizers were placed at both sides of the heatpipe oven in the path of the probe beam. At the frequencies at which transitions induced by the pump beam shared the same



FIG. 1: Part of the polarization spectrum of Na₂ obtained with the 501.7 nm line of the Ar⁺ laser as the probe and circularly polarized pump light. The assigned v' progressions correspond to transitions $4^{1}\Sigma_{\rm u}^{+}(v', J' = J'' \pm 1) \leftarrow X^{1}\Sigma_{\rm g}^{+}(v'' = 2, J'' = 43)$. Note that because of near resonance of vibrational levels in the inner and outer potential wells, in two observed doublets v' numbers differ by one for P and R lines. A systematic disparity of intensities of P and R lines for v' = 52to 59 is presently not understood. Lines denoted by stars correspond to $D^{1}\Pi_{\rm u} \leftarrow X^{1}\Sigma_{\rm g}^{+}$ transition in Na₂, overlapping the investigated band system.

lower levels with the probe transitions, the probe light passed through the analyzer. The signal was recorded with a photomultiplier and processed with a computer. For measurements of polarization spectra labeled via P or R transitions, the pump beam was circularly polarized; for polarization spectra labeled via Q transitions, the linear polarization of the pump beam was chosen [21].

III. RESULTS

We recorded the polarization spectrum of the $4^{1}\Sigma_{u}^{+} \leftarrow X^{1}\Sigma_{g}^{+}$ system of Na₂ in a range between 32100 and 35100 cm⁻¹. Fig. 1 shows a particularly interesting fragment of the spectrum where an abrupt change of vibrational spacing in the observed v' progression corresponds to transition from the inner well to a region above the potential barrier. The analyzed spectral lines provided information about 332 rovibrational levels in the $4^{1}\Sigma_{u}^{+}$ state. The data field is illustrated in Fig. 2. The highest vibrational level identified by us corresponds to v' = 77 (vide infra) whereas the rotational quantum numbers J' are spread between 12 and 56. Assignment of v' = 0 level was based on an assumption that this was the lowest level observed. After Franck-Condon factors for transitions from the ground $X^{1}\Sigma_{g}^{+}$ state were calculated, comparison of them with the measured relative strengths of the spectral lines confirmed the assignment. It must be noted that the outer well in $4^{1}\Sigma_{u}^{+}$ was inaccessible in the present experiment because of negligible overlap between vibrational wave functions located there and those corresponding to the ground state levels. The measured wave numbers of lines have been converted to energies of $4^{1}\Sigma_{u}^{+}$ state levels referred to the bottom of the $X^1\Sigma_g^+$ state potential well, using the ground state molecular constants of Kusch and Hessel [20]. As they reproduce energies of the rovibrational levels in the ground state with an accuracy exceeding the precision of our measurements, no additional errors were introduced into our analysis of the $4^{1}\Sigma_{u}^{+}$ state.

IV. THE IPA ALGORITHM USING A POINTWISE POTENTIAL AND ITS MODIFICATION

The main idea of the IPA is to start with an approximate potential energy curve $U_0(R, \boldsymbol{a})$ with eigenenergies E_i^0 and eigenfunctions Ψ_i^0 . The fitting parameters $\boldsymbol{a} = (a_1, a_2, ..., a_M)$ are used in order to modify $U_0(R, \boldsymbol{a})$ in such a way that its new eigenenergies E_i agree with the experimental ones E_i^{exp} in the LSA sense [4]. If the approximate potential is close to the "true" one, we can assume a linear dependence of the eigenenergies on small changes of the parameters a_j

$$E_i = E_i^0 + \sum_j \frac{\partial E_i}{\partial a_j} \Delta a_j ; \qquad (1)$$



FIG. 2: Range of rovibrational levels used in the analysis of the $4^{1}\Sigma_{u}^{+}$ state in Na₂.

 E_i^0 and $\frac{\partial E_i}{\partial a_j}$ denote the eigenvalue and its first derivatives with respect to a_j for the approximate potential, i.e. at $\boldsymbol{a} = \boldsymbol{a}^0$. Application of the Hellman-Feynman theorem leads to

$$\frac{\partial E_i}{\partial a_j} = <\Psi_i^0 | \frac{\partial U(R, \boldsymbol{a})}{\partial a_j} | \Psi_i^0 > .$$
⁽²⁾

In order to parametrize the molecular potential we use an equidistant grid of points (R_k, u_k) connected with a cubic spline function [6]. Values of the potential at $R = R_k$, that is u_k , are fitting parameters. In this case it is possible to show [6] that

$$U(R, \boldsymbol{u}) = \sum_{k} S_k(R) u_k , \qquad (3)$$

where $S_k(R)$ are known functions of R. Since

$$\frac{\partial U(R, \boldsymbol{u})}{\partial u_j} = S_j(R) , \qquad (4)$$

Eq. 1 can be rewritten as

$$E_{i} = E_{i}^{0} + \sum_{j} \langle \Psi_{i}^{0} | S_{j}(R) | \Psi_{i}^{0} \rangle \Delta u_{j}.$$
(5)

where Δu_j is the correction to the *j*-th fitting parameter. Replacing E_i with E_i^{exp} provides a set of N linear equations (N denotes the number of experimental levels) with M unknowns (where M is the number of the fitted parameters), which should be solved in the LSA sense.

Usually the whole spectrum of eigenvalues is not available and therefore some parts of the potential cannot be reliably retrieved from the experimental data. In case of the pointwise representation of the potential this means that the calculated term energies of the experimentally observed levels are not sensitive to some of the u_k . The inversion problem becomes ill-conditioned then and its solution may be unstable to data perturbation and lacking a physical sense, particularly in terms of a desired smoothness of the constructed potential. The ill-conditioning of the problem does not imply that a meaningful approximate solution cannot be found, but an extra care has to be taken to obtain a physically acceptable result. The singular value decomposition (SVD) method [17] chosen by us for solving the initial system of linear equations offers a partial remedy. Also a sensible choice of a sparse grid for the pointwise potential in the regions badly characterized by the experimental data can flatten to some extent the undesirable wiggles on it (see [22]). Here we propose a simpler, more general and more effective solution, based on an additional constraint imposed on the constructed potential.

Generally, the usual way of obtaining a smooth fitted function $f(R, \mathbf{a})$ is to add to the merit function $\chi^2(\mathbf{a})$ a regularizing functional $H(\mathbf{a})$ which is responsible for the smoothness of the solution and to minimize the sum

minimize:
$$\chi^2(\boldsymbol{a}) + \lambda^2 H(\boldsymbol{a})$$
 (6)

where λ is a parameter used to tune the degree of regularization. A possible form of H(a)

is (see e.g. [17], Chapter 18):

$$H(\boldsymbol{a}) = \int_{R_1}^{R_2} (f''(R, \boldsymbol{a}))^2 dR$$
(7)

Here, by minimizing the integral of the square of the second derivative, one sets the additional condition that the fitted function should be as close to a straight line as possible between R_1 and R_2 . Since in our version of the IPA method the fitted potential is defined as a cubic spline function drawn between given grid points (R_k, u_k) , the second derivative $U''(R, \mathbf{u})$ between these points is a linear function of R. Therefore we define a somewhat simplified form of the regularizing functional as

$$H(\boldsymbol{u}) = \sum_{j} (U''(R_j, \boldsymbol{u}))^2$$
(8)

where $U''(R_j, \boldsymbol{u}) = U''_j$ denotes the second derivative of the potential in a grid point R_j .

Similarly to Eq. (3), in Ref. [6] it was shown that

$$U_j'' = \sum_i L_{ji} u_i. \tag{9}$$

where L_{ji} are known coefficients. Therefore in the present case the regularization condition is reduced to supplementing the system of linear equations (5) by a set of M equations:

$$\sum_{i} \lambda L_{ji} (u_i^0 + \Delta u_i) = 0 \tag{10}$$

Here u_i^0 is the initial value of the *i*-th fitting parameter. Each equation requires that the respective second derivative U_j'' equals zero and the summation is performed over all fitted parameters. When $\lambda = 0$, the system (5) remains unchanged, i.e. no regularization is imposed. Increasing λ results in flattening of the potential in competition with the initial condition defined by (5). In principle the value of λ in (10) can depend on *j*, varying smoothly from zero for the parts of the potential with abundant experimental data to some large value for the regions badly characterized by the experiment; the other sensible possibility is to make the change stepwise and this approach has been adopted in the present work.

V. POTENTIAL CURVE OF THE $4^{1}\Sigma_{u}^{+}$ STATE

With the body of data described in Section III we applied the IPA technique to determine the potential energy curve of the $4^{1}\Sigma_{u}^{+}$ state. Initially we limited our analysis to the inner well of the molecular potential, containing a majority of levels observed in the present experiment. Transitions to rovibrational levels located there formed clear vibrational progressions of spectral lines. However, we observed that distances between subsequent P, R doublets did not decrease in a systematic way even for the first few doublets in each progression, as expected for a well-behaved molecular potential. This observation supported theoretical prediction of substantial deformation of the inner well. In consequence, levels situated in the inner well could not be characterized by a set of molecular constants and a standard RKR procedure could not be used for determination of the inner well. Under these circumstances we rather applied a variant of the RKR method first proposed by Stwalley [2], which allows to construct an approximate potential energy curve from vibrational term values G(v) and rotational constants B_v obtained directly from the experimental spectrum, i.e. from a distance of the subsequent P, R doublets and the P-R combination differences measured in a chosen progression, preferably being the longest one.

The potential curve generated in this way was used then as an approximate starting potential for the standard IPA procedure [6] (i.e. without regularization). It provided a refined potential energy curve which reproduced eigenenergies of levels distant from the top of the potential barrier by more than 50 cm⁻¹ (i.e. the levels, for which the presence of the outer well could be neglected) within the experimental accuracy. In the second part of our analysis we used a larger data set including levels close to and above the potential barrier. In this case the existence of both potential wells has to be taken into account and accordingly we extended the region of internuclear distances R, for which the potential was considered, to $R \approx 16$ Å. As a starting potential for the next run of the IPA routine we adopted a hybrid potential consisting of the inner well from the previous step matched smoothly with the theoretical curve calculated by Magnier [23] used to represent the outer well.

An equidistant grid was used for the approximate potential since Eqs. 3 and 9 are derived under this condition. Although a generalization of both equations is possible also for a nonequidistant grid, this would require significant changes in the fitting code [6]. Moreover, it is not clear *a priori* how to find the optimal distribution of the grid points. Hence initially we defined $U_0(R)$ in 110 equidistant points for 2.4 Å $\leq R \leq 15.9$ Å, i.e. a grid dense enough to describe the steepest changes of the potential curve.

After several iterations the r.m.s. deviation of the fit decreased below 0.05 cm⁻¹, that is the generated IPA potential reproduced all the observed levels with an expected accuracy. At closer inspection, however, the potential turned out unsatisfactory: the outer well displayed unphysical irregularities in a form of ripples, resulting from scarcity of experimental data defining the region of large R values. Therefore we refitted the potential including the regularization condition in the form of Eq. 10. During the fit λ was varied between 1.0 and 4.0 for $R > R_{\rm reg}$ and set to zero otherwise. Several values of $R_{\rm reg}$ between 6.5 and 8.0 Å were tested. The value of λ was gradually increased to smooth the outer potential well, until the quality of the fit started to deteriorate. Once a satisfactory shape of the outer well was achieved, we reduced the number of grid points as described in Ref. [6].

The final version of the potential is presented in Table I and in Fig. 3. The last five points were taken from theoretical calculations [23]. Their influence on the energy level positions is negligible and they were added mainly to ensure proper boundary conditions for solving the Schrödinger equation. In order to calculate the value of the potential for arbitrary R, a natural cubic spline should be used [17]. The potential reproduces 325 of totally 332 observed energy levels with a standard deviation of 0.044 cm⁻¹ and a normalized standard deviation of 0.88. Seven levels were excluded from the fit since the deviations for the calculated term energies exceeded 0.15 cm⁻¹. We attribute these discrepancies to perturbations by the neighboring singlet or triplet states [23].

VI. FINAL REMARKS

In the present work we attempted to construct the highly irregular potential energy curve of the $4^{1}\Sigma_{u}^{+}$ state in Na₂ from the experimental data abundant for levels in the inner potential well but scarce for the region around and above the internal barrier. A version of the IPA procedure involving the pointwise representation of the potential and its regularization was used. The inner well has been determined accurately, however problems arose when trying to find the shape of the barrier and the outer potential well. A small number of the measured level energies influencing these parts of the potential suggested that determination of its shape was not possible. In mathematical terms, inversion of the spectroscopic data to a



FIG. 3: The generated potential curve of the $4^{1}\Sigma_{u}^{+}$ state. Zero on the energy scale corresponds to the bottom of the ground $X^{1}\Sigma_{g}^{+}$ potential [20].

potential curve could not provide a unique result. However, from many possible potentials, representing positions of all the experimental levels equally well, we were able to select one with a sensible physical shape. A general procedure, based on the mathematical concept of regularization of the constructed potential curve, has been suggested to deal with such problems. Still, it should be borne in mind that the outer well of the potential reported in Table I is only a plausible smooth solution to the inversion problem and it would be unreasonable to speculate about its accuracy. The only way to improve its reliability is to collect more experimental data related to this part of the potential.

It should be also noted that the uncertainty of the outer potential well affects the vibrational numbering of the rovibrational levels of the $4^{1}\Sigma_{u}^{+}$ state. Small changes of the outer well may cause a change of the numbering of levels which wave functions are significantly nonzero mainly in the inner well. Of course this will change neither their term energies, nor the relative vibrational numbering of the levels belonging to the inner well, i.e. the quantities directly observable in the present experiment.

Acknowledgments

This work has been funded in part by grant No. 2 P03B 063 23 from the Polish Committee for Scientific Research. A.P. acknowledges a support from the Center of Excellence ASPECT (program "Competitive and Sustainable Growth", G6MA-CT-2002-04021).

- H. Lefebvre-Brion and R. W. Field, Perturbations in the Spectra of Diatomic Molecules (Academic Press Press, New York, 1986).
- [2] W. C. Stwalley, J. Chem. Phys. 56, 2485 (1972).
- [3] A. Pashov, I. Jackowska, W. Jastrzębski, and P. Kowalczyk Phys. Rev. A 58, 1048 (1998).
- [4] W. Kosman and J. Hinze J. Mol. Spectrosc. 51, 341 (1974).
- [5] C. R. Vidal and H. Scheingraber J. Mol. Spectrosc. 65, 46 (1977).
- [6] A. Pashov, W. Jastrzębski, and P. Kowalczyk Comp. Phys. Commun. 128, 622 (2000).
- [7] C. Samuelis, E. Tiesinga, T. Laue, M. Elbs, H. Knöckel, and E. Tiemann Phys. Rev. A 63, 012710 (2000).

- [8] J. Y. Seto, R. J. Le Roy, J. Vergés, and C. Amiot J. Chem. Phys. 113, 3067 (2001).
- [9] Y. Huang, Ph. D. Thesis, University of Waterloo, 2001.
- [10] Y. Y. Huang and R. J. Le Roy, J. Chem. Phys. 119, 7398 (2003).
- [11] W. Jastrzebski, W. Jaśniecki, P. Kowalczyk, R. Nadyak, and A. Pashov, Phys. Rev. A 62, 042509 (2000).
- [12] A. Pashov, W. Jastrzębski, and P. Kowalczyk, J. Chem. Phys. 113, 6624 (2000).
- [13] F. Martin, P. Crozet, A. J. Ross, M. Aubert-Frécon, P. Kowalczyk, W. Jastrzębski, and A. Pashov, J. Chem. Phys. 115, 4118 (2001).
- [14] W. Jastrzebski, P. Kowalczyk, R. Nadyak, and A.Pashov, Spectrochim. Acta A 58, 2193 (2002).
- [15] W. Jastrzebski, R. Nadyak, and P. Kowalczyk, Chem. Phys. Lett. 374, 297 (2003).
- [16] A. Grochola, W. Jastrzebski, P. Kowalczyk, A. Ross, and P. Crozet, *Chem. Phys. Lett.* 372, 173 (2003).
- [17] W. H. Press, S. A. Teukolski, W. T. Vetterlingand, and B. P. Flannery, Numerical Recipes in Fortran 77 (Cambridge University Press, Cambridge, 1997).
- [18] W. Jastrzębski and P. Kowalczyk, Phys. Rev. A 51, 1046 (1995).
- [19] A. Grochola, W. Jastrzebski, P. Kortyka, and P. Kowalczyk, J. Mol. Spectrosc. 221, 279 (2003).
- [20] P. Kusch and M. M. Hessel, J. Chem. Phys. 68, 2591 (1978).
- [21] R. Ferber, W. Jastrzębski, and P. Kowalczyk, J. Quant. Spectrosc. Rad. Transfer 58, 53 (1997).
- [22] A. Pashov, W. Jastrzębski, and P. Kowalczyk, J. Phys. B At. Mol. Opt. Phys. 33, L611 (2000).
- [23] S. Magnier, Ph. D. Thesis, Université de Paris-Sud, 1993.

	1	u u	-
R [Å]	$U \ [cm^{-1}]$	${ m R}[{ m \AA}]$	$U \ [cm^{-1}]$
2.4000	40604.123	6.7257	35166.484
2.5239	38549.528	6.9736	35258.930
2.6477	36908.022	7.2216	35327.723
2.7716	35776.823	7.4695	35373.419
2.8954	35026.062	7.7175	35405.959
3.0193	34379.006	7.9650	35425.336
3.1431	33888.597	8.5318	35442.782
3.2670	33549.507	9.0986	35441.533
3.3908	33332.544	9.6654	35434.522
3.5147	33214.236	10.2321	35419.888
3.6385	33174.430	10.7989	35401.202
3.7624	33195.060	11.3657	35380.528
3.9359	33291.770	11.9325	35358.744
4.1119	33428.812	12.4993	35338.612
4.2878	33562.727	13.0661	35326.650
4.4638	33673.469	13.6329	35333.401
4.6397	33769.607	14.1996	35369.969
4.8157	33868.489	14.7664	35446.576
4.9900	33976.789	15.3332	35564.687
5.2380	34148.711	15.9000	35726.822
5.4859	34333.616	16.9000	36005.000
5.7339	34525.107	17.5000	36150.000
5.9818	34715.356	18.0000	36233.000
6.2298	34890.866	19.0000	36277.700
6.4777	35042.921	20.0000	36284.700

TABLE I: The IPA potential energy curve of the $4^{1}\Sigma_{u}^{+}$ state in Na₂.