

# Manipulating the scattering length of a Bose-Einstein condensate in an amplitude - modulated optical lattice.

G.M.Genkin\*.

*Department of Physics and Astronomy, Northwestern University , Evanston , Illinois , 60208 .*

## Abstract

The scattering length in a BEC confined in an amplitude - modulated optical lattice can be manipulated by the modulation strength. Two standing laser waves of main and sideband frequencies of an optical lattice induce a Raman transition; due to resonant Raman driving the effective scattering length depends on the modulation strength and the detuning from resonance and can be tuned to a given value.

\* Electronic address: gena@pluto.phys.nwu.edu.

03.75.Lm, 05.30.Jp.

The recent experimental observation of Bose-Einstein condensation in a dilute gas of ultracold trapped atoms [ 1-3 ] has generated much interest in studying the properties of a Bose-Einstein condensate ( BEC ) and manipulating such coherent matter by external electromagnetic fields. The experimental observation of Bose-Einstein condensation in a dilute gas of ultracold trapped atoms [ 1-3 ] has generated much interest in manipulating such coherent matter by external electromagnetic fields. Recently atoms have been confined in optical potentials created by standing light waves [ 4, 5 ], with the wavelength of the optical potential much smaller than the dimensions of the trap. The unique prospects of this new class of system (the correlated bosons on a lattice ) became evident. The possibility of creating optical lattice in trapped Bose - condensed gases has provided an opportunity to study superfluids in novel situations. The presence of the lattice leads to a variety of solid - state effects. For example, a transition to a Mott insulator becomes possible when a superfluid like a BEC is placed in a periodic potential [4]; the oscillation frequency of the center - of - mass motion of the condensate is reduced [6] as a result of the enhanced effective mass of the atoms tunneling between potential wells. By accelerating the optical lattice [7] Bloch oscillations of the condensate have been observed, and reducing the amplitude of the lattice optical potential leads to a breakdown of these oscillations as a result of Landau - Zener tunneling between bands.

Many physical properties of a dilute gas are determined by the atom - atom interaction. In a dilute and cold gas only binary collisions at low energy are relevant and these collisions are characterized by a single parameter, the  $s$  - wave scattering length. The confinement of a gas in one or more spatial dimensions can strongly modify the collisional properties (two - body scattering in an optical lattice ) of atoms [8 - 12]. In Ref.[12] was shown that in an 3D optical lattice a dilute gas is described by the effective atomic scattering length  $a_{eff}$ , modified and tunable by the lattice parameters,

$$a_{eff} = \frac{\pi \ln 2}{4} a \frac{d}{a + l_*} \frac{\hbar}{m\omega l_0^2},$$

$$l_* = \frac{\ln 2}{4} l_0 \sqrt{D_0}, \quad (1)$$

where  $a$  is the free - space scattering length, the optical lattice potential is determined by a  $V_0$  and lattice spacing  $d$ , and  $l_0$  is the size of the ground state wave function in an individual well of an optical lattice. The wave functions of the relative motion of two particles of mass  $m$  are characterized by the quasimomentum  $q$ , the band index  $s$ , and the energy  $\varepsilon_{sq} = \varepsilon_s - t_s \cos(qd)$ , where  $t_s = \sqrt{D_s} \hbar \omega / \pi$ ,  $D_s \ll 1$  is WKB tunneling exponent between the neighboring wells,  $\omega^2 = V_0 / (md^2)$ ,  $\varepsilon_s$  are the energies of states  $s$  in isolated wells. The effective scattering length  $a_{eff}$  is modified by the optical lattice if the tunneling between the neighboring lattice sites is sufficiently small and the new length scale  $l_*$  contains the tunneling amplitude ( for the band index  $s = 0$  the amplitude  $D_0$  ) and describes the influence of particle states extending over several lattice sites on the two - body scattering in an optical lattice. In Ref. [12] was shown that in the case of attractive interaction, for the free - space length  $a < 0$ , the effective scattering length exhibits resonant behavior and leads to a geometric resonance at  $l_* = |a|$ , and for  $l_* > |a|$  the effective scattering becomes repulsive  $a_{eff} > 0$ .

The goal of this paper is manipulating of the effective scattering length by a time - dependent action of the lattice parameters. We consider a manipulation of the scattering length in a BEC in an amplitude - modulated optical lattice. The such optical lattice can be created by interfering pair of amplitude - modulated beams. In general, there are the two generic types ( a resonant and a nonresonant cases ) of the manipulation. The effect of the modulation is to modulate the optical lattice potential so that it becomes a time - dependent optical lattice potential which the depth of the optical potential is

$$V_0(t) = V_0(1 + \nu \cos \omega_M t), \nu < 1, \quad (2)$$

where  $\nu$  is the modulation strength,  $\omega_M$  is the modulation frequency. If the period of modulation  $2\pi/\omega_M$  is larger the nonequilibrium dynamic time  $\tau_{noneq}$ , then the time behavior of a BEC in an amplitude - modulated optical lattice is quasistatic. The nonequilibrium dynamic time  $\tau_{noneq}$  for trapped bosonic atoms in an optical lattice potential was considered in Ref.[11] and was shown that the characteristic time of the dynamical restoration of the

phase coherence is a Josephson time  $h/J$ .

*A resonant case.*

One is a resonant mechanism which operates when the modulation frequency  $\omega_M$  is close to the excited internal state of a BEC. A resonant mechanism is determined by change in the population of the ground state due to a resonant driving. In a BEC in a periodic potential the characteristic energetic scale of the band structure [14,15] is the recoil energy  $E_R$ . And, usually,  $E_R$  exceeds  $J$ , and the quasistatic behavior is not valid. However, in this case is more convenient to use a picture that an amplitude - modulated field can be presented as a set of monochromatic fields (main on frequency  $\omega$  and sideband on frequencies  $\omega \pm \omega_M$  ). These fields can induce a Raman transition between the two Bloch bands of a BEC  $s = 0$  and  $s \neq 0$ . A resonant field via a two - photon ( Raman ) transition changes the population of the lowest Bloch band  $s = 0$  of a BEC. In resonant case the system is excited by an off - resonant Raman ( two - photon ) driving with the Rabi frequency  $\Omega_R$ . In an amplitude - modulated optical lattice two standing laser waves of frequencies  $\omega$  and  $\omega - \omega_M$  ( $\omega_M \ll \omega$ ) drive an atom in a Raman scheme. The effective two - photon Rabi frequency is given in terms of the single - photon Rabi frequencies  $\Omega$  and  $\Omega'$  of the fields on the main  $\omega$  and sideband  $\omega' = \omega - \omega_M$  frequencies as  $\Omega_R = \frac{\Omega \cdot \Omega'}{2\Delta}$ . For the amplitude - modulated laser beams with the modulation strength  $\nu$  we have for standing waves  $\Omega_R(\mathbf{x}) = \frac{\nu \Omega^2(\mathbf{x})}{2\Delta} = \frac{\nu V_0(\mathbf{x})}{h}$ , here  $\Delta$  is the far - detuning of the laser beams from the atomic resonance (the closest neighboring optical dipole transition), and  $V_0(\mathbf{x}) = \sum_{j=1}^3 V_{j0} \cos^2 kx_j$  is the optical lattice potential with wave vectors  $k = \frac{2\pi}{\lambda}$  and  $\lambda$  the wavelength of the laser beams on the main frequency, and we are omitted the terms proportional to the small parameter  $\Delta k \cdot x = \frac{\omega_M}{c} x \ll 1$ , where  $\Delta k \equiv k' - k$ , for  $x < \lambda \sim 10^{-4}cm$ ,  $\omega_M \preceq 10^7 s^{-1}$ , this parameter is about  $10^{-7}$ . The eigenstates of a BEC are Bloch waves, and an appropriate superposition of Bloch states yields a set of Wannier functions which are localized on the individual lattice sites. Without a resonant driving the wave functions in a tight binding model can be represented [12] as  $w_s(\mathbf{x}) = Z_s \Psi_s(\mathbf{x})$ , where  $\Psi_s(\mathbf{x})$  is the wave function of the oscillator in a state  $s$ , and  $Z_s$  is a normalization factor. However, for a resonant driving the wave functions are

$w_s^{res}(\mathbf{x}) = \varphi(\mathbf{x})w_s(\mathbf{x})$ , where the function  $\varphi(\mathbf{x})$  is the solution of the Bloch equation of the standard Rabi problem. We consider a steady state driving regime, in which the function  $\varphi(\mathbf{x})$  is determined on the parameters of the field ( the Rabi frequency and the detuning from resonance  $\delta = \omega_M - (\omega_b - \omega_a)$  ). In order to avoid a sufficient population of the excited Bloch band  $s \neq 0$ , we are assume  $\Omega_R < \delta$ , then  $\varphi^2(\mathbf{x}) \simeq 1 - \frac{1}{2}(\frac{\Omega_R(\mathbf{x})}{\delta})^2$ . Using the wave functions  $w_s^{res}$ , in which the size of the ground state oscillator wave function  $l_0$  in a lattice is much less than the lattice period  $d = \lambda/2$ , and following Ref. [12] we find the effective scattering length  $a_{eff}^{res}$  modified by the resonantly amplitude modulation

$$\frac{a_{eff}^{res} - a_{eff}}{a_{eff}} \simeq -\left(\frac{\nu V_0}{h\delta}\right)^2. \quad (3)$$

As a result, we can manipulate the effective scattering length in a BEC in an resonantly amplitude - modulated optical lattice. This tuning is determined by the modulation strength  $\nu$  and the detuning  $\delta$ . By varying the parameter  $\nu/\delta$ , we can tune  $a_{eff}^{res}$  to a given value.

*A nonresonant case.*

Second is a nonresonant mechanism for which we will use a direct time - dependent description. Note that such potential is created by interfering pair of amplitude - modulated laser beams with the modulation strength  $\nu/2$  of the electric field amplitude, and was omitted the quadratic term for the small parameter  $\nu^2/4$ . A nonresonant low - frequency mechanism of manipulation is in operation for a quasistatic regime  $\omega_M < 2\pi\tau_{noneq}^{-1}$ , therefore, if  $V_0$  corresponds to a geometric resonance at  $l_* = |a|$  for attractive interaction then as the time runs, the system goes from an attractive interaction to a repulsive interaction and back again with frequency  $\omega_M$ . Therefore, it is allow to investigate the nonequilibrium dynamics in a BEC. In a nonresonant case  $\Omega_R/\delta \rightarrow 1$  then  $\varphi(\mathbf{x}) = 1$  and we have a time - dependent WKB tunneling exponent  $D_0(t)$ . The WKB tunneling exponent is a nonlinear function on the strength of the periodic potential  $V_0$  which is proportional to the laser intensity . If we have a time - dependent lattice laser intensity, then, although  $\overline{V_0(t)} = V_0$ , due to the nonlinearity of a WKB tunneling exponent a time - averaged  $\overline{D_0(t)}$  will be different from a WKB tunneling exponent  $D_0$  without a modulation. For a small  $\nu$  we have the

time - averaged  $\overline{D_0(t)} = D_0 + \Delta D_0$ , where the bar stands for a time average and  $\Delta D_0$  is proportional to the time - averaged  $\nu^2 \overline{\cos^2 \omega_M t}$  ( the time - averaged of the modulation harmonic oscillation to the lowest even order power ). For a low - frequency nonresonant manipulation we will use a time - dependent potential ( Eq.(2)), and there could be a modulation frequency  $\omega_M$  low enough for quasistatic behavior. Using the WKB tunneling exponent  $D_0 \simeq \exp(-\frac{\pi d}{\hbar} \sqrt{2mV_0})$  and using  $V_0(t)$  (Eq.(2)), we have a time - dependent length scale  $l_*^{non}(t)$  and, correspondingly, a time - dependent effective scattering length  $a_{eff}^{non}(t)$ . Calculating the time - average  $\overline{D_0(t)}$  by expanding in a series about the small parameter  $2mV_0\pi^2 d^2/\hbar^2 < 1$ , we find the time - average length scale  $\overline{l_*^{non}}$ , which is modified by the nonresonant amplitude modulation

$$\frac{\overline{l_*^{non}} - l_*}{l_*} \simeq \frac{\nu^2}{4} \frac{2mV_0\pi^2 d^2}{\hbar^2}. \quad (4)$$

If  $\frac{\nu^2}{4} \frac{2mV_0\pi^2 d^2}{\hbar^2}$  is not small, then we may expand the expression for the time - dependent  $D_0(t)$  using Bessel functions  $I_n(x)$  and the functions  $\cos^n \omega_M t$ . In this case instead the parameter  $\frac{\nu^2}{4} \frac{2mV_0\pi^2 d^2}{\hbar^2}$  we have  $I_2(\nu \frac{\pi d}{\hbar} \sqrt{2mV_0})$ , because, usually  $\frac{\pi d}{\hbar} \sqrt{2mV_0} \gg 1$  and for  $\nu < 0.4$  among  $I_n(x < 2)$  the largest is  $I_2(x)$ .

In summary, the effective scattering length in a BEC in an optical lattice can be manipulated by an amplitude - modulated manner. The effect of the modulation is to modulate the optical lattice potential so that it becomes a time - dependent optical lattice potential depth and, correspondingly, the effective scattering length becomes time - dependent. There are two mechanisms of the manipulation. One, a resonant mechanism is in operation for the modulation frequency closed to the excited internal state of a BEC, and two standing laser waves of main and sideband frequencies of an amplitude - modulated optical lattice induce a Raman transition. Due to resonant Raman driving the population of the lowest Bloch band  $s = 0$  of a BEC changes, therefore, the effective scattering length is also changed. As a result, the effective scattering length is other than without a modulation, this tuning is determined by the square of the modulation strength and the square of the detuning. Second, a nonresonant low - frequency mechanism, for which we use a direct time - depen-

dent description. Due to the nonlinear dependence of the WKB tunneling amplitude on the optical lattice depth for the harmonic modulation of the optical lattice depth the time - averaged tunneling exponent  $\overline{D_0(t)}$  is different from  $D_0$  without a modulation. The proposed mechanism of manipulating the effective scattering length makes it possible to tune it to a given value. As a result, although a time - averaged optical lattice potential with a harmonic modulation is equal to an optical lattice potential without a modulation, the effective scattering length in a BEC confined in an amplitude - modulated optical lattice can be manipulated by the modulation strength. In conclusion, the nonequilibrium dynamics of a BEC can be investigated by manipulating the effective scattering length of a BEC in an amplitude - modulated optical lattice.

I thank A.J.Freeman for encouragement.

## REFERENCES

- [1] M. H. Anderson *et al.*, Science **269**, 198 (1995).
- [2] C. C. Bradley *et al.*, Phys. Rev. Lett. **75**, 1687 (1995).
- [3] K. B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995).
- [4] C. Orzel *et al.*, Science **291**, 2386 (2001).
- [5] M. Greiner *et al.*, Nature (London) **415**, 39 (2002).
- [6] F. S. Cataliotti *et al.*, Science **293**, 843 (2001).
- [7] O. Morsch *et al.*, Phys. Rev. Lett. **87**, 140402 (2001).
- [8] M. Olshanii, Phys. Rev. Lett. **81**, 938 (1998).
- [9] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Phys. Rev. Lett. **85**, 3745 (2000).
- [10] D. S. Petrov, M. Holzmann, and G. V. Shlyapnikov, Phys. Rev. Lett. **84**, 2551 (2000).
- [11] D. S. Petrov and G. V. Shlyapnikov, Phys. Rev. A **64**, 012706 (2001).
- [12] P. O. Fedichev, M. J. Bijlsma, and P. Zoller, Phys. Rev. Lett. **92**, 080401 (2004).
- [13] A. Polkovnikov, S. Sachdev, and S. M. Girvin, Phys. Rev. A **66**, 053607 (2001).
- [14] K. Berg - Sørensen and K. Mølmer, Phys. Rev. A **58**, 1480 (1998).
- [15] D. Diakonov *et al.*, Phys. Rev. A **66**, 013604 (2002).