

Capillary flow in an evaporating sessile drop

Yuri Yu. Tarasevich

*Astrakhan State University, Tatishcheva 20, Astrakhan, 414056, Russia**

An analytical expression of velocity potential inside an evaporating sessile drop with pinned contact line is found.

PACS numbers: 47.55.Dz

I. INTRODUCTION

The desiccated sessile drops attract the attention because of different reasons and possible applications. If a drop of pure liquid dries on a smooth substrate, then the base of the drop shrinks but the contact angle remains fixed. Under evaporation, a drop of colloidal suspension or solution having a strongly anchored three phase line keeps a spherical-cap shape with a constant base. The contact angle decreases with time and the height in each point of the profile decreases. To satisfy the anchoring condition, a flow of liquid has to occur inside the drop. An outward flow in a drying drop is produced when the contact line is pinned so that liquid that is removed by evaporation from the edge of the drop must be replenished by a flow of liquid from the interior [1, 2, 3]. This flow is capable of transferring 100% of the solute to the contact line and thus accounts for the strong perimeter concentration of many stains. Ring formation in an evaporating sessile drop is a hydrodynamic process in which solids dispersed in the drop are advected to the contact line. After all the liquid evaporates, a ring-shaped deposit is left on the substrate that contains almost all the solute. Perhaps, everybody knows a dense, ring-like deposit along the perimeter of a dried drop of coffee or tee, milk or juice on a table.

Exploratory experiments [3] using a variety of carrier fluids, solutes, and substrates indicated that preferential deposition at the contact line is insensitive to a wide range of experimental conditions. Ringlike deposits were observed whenever the surface was partially wet by the fluid irrespective of the chemical composition of the substrate. Different substrates were investigated (metal, polyethylene, roughened Teflon, freshly cleaved mica, ceramic, and silicon). Rings were found in big drops (15 cm) and in small drops (1 mm). They were found with aqueous and non-aqueous (acetone, methanol, toluene, and ethanol) solvents. They were found with solutes ranging in size from the molecular (sugar and dye molecules) to the colloidal (10 μm polystyrene microspheres) and with solute volume fractions ranging from 10^{-6} to 10^{-1} . Likewise, environmental conditions such as temperature, humidity, and pressure could be extensively varied without affecting the ring. Effects due to

solute diffusion, gravity, electrostatic fields, and surface tension forces are negligible in ring formation.

The vertically averaged radial flow of the fluid in a desiccated sessile drop was calculated in the work [1]. The conservation of fluid was utilized. A constant evaporation rate all over drop free surface was assumed.

For the case where the limiting rate is the diffusion of the liquid vapor, the evaporation of the drop rapidly attains a steady state so that the diffusion equation reduces to Laplace's equation [3]. In this case the evaporation rate is larger near the contact line and the resulting ringlike deposit is more concentrated at the edge.

The models [1] and [3] deal with the vertically averaged radial flow of the fluid. The next Section describes how to obtain the space distribution of the fluid flow.

II. MODEL AND RESULTS

Likewise the works [1, 2, 3], let us suppose, that the shape of the drop is a spherical cap. It means that the drop is small and a spherical cap shape induced by surface tension (Bond number smaller than one: $\text{Bo} = g(\rho - \rho_f)d^2\sigma^{-1} \ll 1$, where g is gravitational acceleration, ρ is drop density, ρ_f is surrounding medium density, σ is surface tension, d is drop diameter). The problem of a spherical cap on an impermeable substrate can be replaced by the problem of a lens. Let us consider an extremely simple situation when the cap is a semi-sphere, i.e. the contact angle is 90° .

Under evaporation, the height in each point of the profile decreases. Nevertheless, under room condition the evaporation is a slow process. So, the desiccation time of a 15 mg sessile water drop is larger than 3500 s [4]. This fact gives a possibility to consider a quasi-static process. We will consider a drop with a fixed free surface and the specific boundary conditions. Quasi-static approximation was utilized in [3] to obtain a vapor rate near the free surface, too.

Let us suppose that the evaporation rate is uniform all over drop free surface, i.e. we will consider only begin of the evaporation. The conservation of fluid determines the relationship between the velocity of the free surface, u_n , the normal to the free surface flow of the fluid, v_n , and the rate of mass loss per unit surface area per unit time from the drop by evaporation, J :

$$\rho v_n + J = \rho u_n.$$

*Electronic address: tarasevich@astranet.ru

Let us consider the potential flow, only. The potential of the velocity field, φ , is a solution of the Laplace's equation

$$\Delta\varphi = 0. \quad (1)$$

Also, we will solve the Laplace's equation inside a spherical area using spherical coordinates.

Measurements of the evolution of the drying drop height h with time t can be performed. Let us suppose, that the velocity of the drop height, u_0 , is known. Thus $u_n = u_0 \cos \theta$, where θ is azimuthal angle. The volume of the drop is decreasing due to evaporation:

$$\int (J - \rho u_0 \cos \theta) dS = 0,$$

where the integration goes all over the free surface. Thus

$$J = \frac{u_0 \rho}{2}.$$

The equation of solute conservation yields,

$$v_r(r = R) = \frac{\partial \varphi}{\partial r} \Big|_{r=R} = u_0 \left(\frac{1}{2} - |\cos \vartheta| \right), \quad (2)$$

where R is the radius of the contact base.

We looking for the boundary value problem (1), (2) inside a spherical area ($r \leq R$) as

$$\varphi(r, \vartheta) = \sum_{k=0}^{\infty} A_k \left(\frac{r}{R} \right)^k P_k(\cos \vartheta), \quad (3)$$

where $P_k(\cos \vartheta)$ are Legendre polynomials.

Thus,

$$\frac{\partial \varphi}{\partial r} = \sum_{k=0}^{\infty} k \frac{A_k}{R^k} r^{k-1} P_k(\cos \vartheta). \quad (4)$$

Let us write the right hand side of the equation (2) using Fourier–Legendre series expansion also known as a Generalized Fourier Series expansion:

$$u_0 \left(\frac{1}{2} - |\cos \vartheta| \right) = \sum_{k=0}^{\infty} b_k P_k(\cos \vartheta). \quad (5)$$

Where,

$$b_k = \frac{2k+1}{2} u_0 \int_0^\pi \left(\frac{1}{2} - |\cos \vartheta| \right) P_k(\cos \vartheta) \sin \vartheta d\vartheta.$$

Thus, $b_0 = 0$, $b_{2k+1} = 0$,

$$b_{2k} = u_0 (4k+1) \frac{(-1)^k (2k-2)!}{2^{2k} (k-1)! (k+1)!}, \quad k = 1, 2, \dots$$

Taking into account (2), (4) equals (5), if $r = R$:

$$\sum_{k=0}^{\infty} k \frac{A_k}{R^k} R^{k-1} P_k(\cos \vartheta) = \sum_{k=0}^{\infty} b_k P_k(\cos \vartheta). \quad (6)$$

Eq. 6 is always valid, if $A_k = b_k R/k$. Hence, the coefficients of the series (3) are

$$A_{2k} = u_0 R (4k+1) \frac{(-1)^k (2k-2)!}{2^{2k} k! (k+1)!}, \quad k = 1, 2, \dots$$

The capillary flow carries a fluid from the apex of the drop to the contact line (Fig. 1).




FIG. 1: Capillary flow in a desiccated sessile drop

III. DISCUSSION

It is clear, more realistic assumption, that a drop has a spherical cap shape with an arbitrary contact angle, can not change the qualitative picture of the flow. In particular, the calculations of the vertically averaged velocity for the drops with the contact angles up to 90° [1] did not show the qualitative changes of the flow. If a drop has a spherical cap shape with an arbitrary contact angle the Laplace's equation can be solved using toroidal coordinates. Although, the analytical results are too complicate in this case, and numerical calculations should be performed. Nonuniform evaporation rate can not change the results qualitative, but the line, where $v_r(R) = 0$, has to move to the base.

Recently, the buckling instability was investigated during the drying of sessile drops of colloidal suspensions or of polymer solutions [5, 6, 7]. Under solvent evaporation, disperse particles or polymers accumulate near the vapor–drop interface. The outer layer of the drop

is more concentrated in the polymer and may display a gel or glassy transition and, hence, may form a permeable rigid gelled or glassy skin. This skin behaves like an elastic shell although it does not block the evaporation. This gelled or glassy skin will thus bend as the volume it encloses decreases, leading to large surface distortions.

The processes of skin formation are not the scope of this brief communication. Nevertheless, knowledge of the velocity field inside an evaporating sessile drop is a necessary background to describe the colloidal particles transfer or solute diffusion, and skin formation.

-
- [1] F. Parise and C. Allain, *J. Phys. II France* **6**, 1111 (1996).
[2] R. D. Deegan, O. Bakajin, T. F. Dupont, G. Huber, S. R. Nagel, and T. A. Witten, *Nature (London)* **389**, 827 (1997).
[3] R. D. Deegan, O. Bakajin, T. F. Dupont, G. Huber, S. R. Nagel, and T. A. Witten, *Phys. Rev. E* **62**, 756 (2000).
[4] C. C. Annarelli, J. Fornazero, J. Bert, and J. Colombaria, *Eur. Phys. J. E* **5**, 599 (2001).
[5] L. Pauchard and C. Allain, *Europhys. Lett.* **62**(6), 897 (2003).
[6] L. Pauchard and C. Allain, *C. R. Physique* **4**, 231 (2003).
[7] L. Pauchard and C. Allain, *Phys. Rev. E* **68**, 052801 (2003).