

# Configuration interaction calculation of hyperfine and P,T-odd constants on $^{207}\text{PbO}$ excited states for the electron EDM experiments

A.N. Petrov,<sup>1,\*</sup> A.V. Titov,<sup>1</sup> T.A. Isaev,<sup>1</sup> N.S. Mosyagin,<sup>1</sup> and D. DeMille<sup>2</sup>

<sup>1</sup>*Petersburg Nuclear Physics Institute, Gatchina, Leningrad district 188300, Russia*

<sup>2</sup>*Physics Department, Yale University, New Haven, Connecticut 06520, USA*

We report first configuration interaction calculations of hyperfine constants  $A_{\parallel}$  and the effective electric field  $W_d$  acting on the electric dipole moment of the electron, in two excited electronic states of  $^{207}\text{PbO}$ . A new combined scheme of correlation calculations is also developed and first applied to studying the core properties. The obtained hyperfine constants,  $A_{\parallel} = -3826$  MHz for the  $a(1)$  state and  $A_{\parallel} = 4887$  MHz for the  $B(1)$  state, are in very good agreement with the experimental data,  $-4113$  MHz and  $5000 \pm 200$  MHz, respectively. We find  $W_d = -(6.1_{-0.6}^{+1.8}) \cdot 10^{24}$  Hz/(e · cm) for  $a(1)$ , and  $W_d = -(8.0 \pm 1.6) \cdot 10^{24}$  Hz/(e · cm) for  $B(1)$ .

*Introduction.* The search for the electric dipole moment (EDM) of the electron,  $d_e$ , remains one of the most fundamental problems in physics. Up to now only upper limits were obtained for  $|d_e|$ ; the latest constraint,  $|d_e| < 1.6 \cdot 10^{-27}$  e·cm, was obtained in an experiment on atomic Tl [1]. The search for  $d_e$  in heavy-atom polar molecules was initiated by theoretical investigations [2, 3] on radicals with one unpaired electron. (The first experimental bound on  $d_e$  using such a system was recently obtained, using YbF [4].) It was since noted that experiments on the excited  $a(1)$  [5] or  $B(1)$  [6] states of PbO, which have two unpaired electrons, may make it possible to search for  $d_e$  with sensitivity 2-4 orders of magnitude higher than the current limit. An important feature of such experiments is that the knowledge of the effective electric field,  $W_d$ , seen by an unpaired electron is required for extracting  $d_e$  from the measurements.  $W_d$  can not be obtained in an experiment; rather, electronic structure calculations are required for its evaluation. Recently, a semiempirical estimate of  $W_d$  for the  $a(1)$  state [7] and *ab initio* relativistic coupled cluster (RCC) [8] calculations [9] with the generalized relativistic effective core potential (GRECP) [10, 11] of  $W_d$  for both the  $a(1)$  and  $B(1)$  states were performed. The nonvariational one-center restoration (NOCR) technique [10, 12, 13, 14] was employed for obtaining proper electronic densities close to nuclei after GRECP/RCC calculations of heavy-atom molecules. The semiempirical estimate of Ref. [7],  $|W_d| \geq 12 \times 10^{24}$  Hz/(e·cm), is three times higher than the RCC result (see Table I). Calculations performed in [9] demonstrated the need for a higher level of accounting for correlation in the valence region for the excited  $a(1)$  and  $B(1)$  states. The main problem was that the Fock-space RCC with single and double excitations (RCC-SD) version used in [9] was not optimal for accounting for the nondynamic correlations (see [15] for details of RCC-SD calculations of the Pb atom), though the potential of the RCC method for electronic structure calculations is very high in prospect [8]. The estimated error bounds put

the actual  $W_d$  value between 75% and 150% of the value calculated by the GRECP/RCC/NOCR approach.

The main goal of the present work is to develop a method of calculation which could provide substantially higher accuracy and reliability for  $W_d$  in the excited states of PbO and could be efficiently applied to other heavy-atom molecules. As an accuracy check for the calculation of the electronic structure near the Pb nucleus, the hyperfine constants ( $A_{\parallel}$ ) are also calculated. For this purpose in the present work the SODCI code [16, 17] for the spin-orbit direct configuration interaction (CI) calculation is employed, in which the relativistic (spin-orbit) scheme [18] and new criteria [18, 19] of the configuration selection proposed by us earlier are incorporated. New codes for calculation of a one-electron density matrix with the CI wavefunction and restored four-component molecular spinors have been developed. The 10-electron GRECP/SODCI/NOCR calculations are performed for  $A_{\parallel}$  and  $W_d$  in the  $a(1)$  and  $B(1)$  states of PbO with wavefunctions restored in the Pb core (see [9, 20, 21] for recent two-step calculations).

The space of a SODCI calculation consists of a set of many-electron configurations obtained by selecting the singly- and doubly-excited configurations with respect to some multiconfigurational reference states when using some selection thresholds ( $T$ ) [18, 22]. Because of limited computational resources, all the possible singly- and doubly-excited configurations (without selection) cannot be usually included to the SODCI calculation. The corrections accounting for zero threshold ( $T=0$ ) [22] and higher-order excitations (“full-CI”) [23] allow one to estimate the contribution to the total energy of all the configurations omitted in the SODCI calculation (see below for more details). In the present work we first apply the  $T=0$ -, full-CI- and “outercore correlations”-type corrections to calculating the properties which cannot be obtained from the energy curves (surfaces). The applicability of these corrections is justified and discussed in details in [24]. The developed method has allowed us to reduce

the deviation of the calculated  $A_{\parallel}$  from its experimental value for  $a(1)$  in about five times, from 34% to 7%, as compared to the previous calculation [9]. That deviation for  $B(1)$  is reduced from  $(18 \pm 4)\%$  to  $(2 \pm 4)\%$ , i.e., it is now within the experimental uncertainty  $\pm 4\%$ . The SODCI results without applying the  $T=0$ , full-CI and outercore correlation corrections give the deviations 12% for  $a(1)$  and  $(8 \pm 4)\%$  for  $B(1)$ . As a consequence, substantially more reliable values for  $W_d$  are obtained. Our present error estimate on  $W_d$  is relatively more conservative than in [9] since  $|W_d|$  is increased from the RCC-SD datum on 50% for  $a(1)$  and decreased on 18% for  $B(1)$  (these notable changes are, however, within the error limits declared in [9]). The expression for  $W_d$  is given by Refs. [9, 25, 26]

$$W_d = \frac{1}{\Omega d_e} \langle \Psi_X | \sum_i H_d(i) | \Psi_X \rangle, \quad (1)$$

where  $\Psi_X$  is the wavefunction for  $a(1)$  or  $B(1)$  state, and  $\Omega = \langle \Psi_X | \mathbf{J} \cdot \mathbf{n} | \Psi_X \rangle = \pm 1$ ,  $\mathbf{J}$  is the total electronic momentum,  $\mathbf{n}$  is the unit vector along the molecular axis directed from Pb to O,

$$H_d = 2d_e \begin{pmatrix} 0 & 0 \\ 0 & \sigma \mathbf{E} \end{pmatrix}, \quad (2)$$

$\mathbf{E}$  is the inner molecular electric field,  $\sigma$  are the Pauli matrices. The hyperfine constant  $A_{\parallel}$  is determined by the expression [27]

$$A_{\parallel} = \frac{1}{\Omega} \frac{\mu_{\text{Pb}}}{I} \langle \Psi_X | \sum_i \left( \frac{\boldsymbol{\alpha}_i \times \mathbf{r}_i}{r_i^3} \right)_Z | \Psi_X \rangle, \quad (3)$$

where  $I$  and  $\mu_{\text{Pb}}$  is the spin and magnetic moment of  $^{207}\text{Pb}$ ,  $\boldsymbol{\alpha}_i$  are the Dirac matrices for the  $i$ -th electron, and  $\mathbf{r}_i$  is its radius-vector in the coordinate system centered on the Pb atom.

*Methods and calculations.* The above properties were calculated in two steps. At the *first step*, a 22-electron GRECP for Pb [28] simulating interaction with the explicitly excluded  $1s$  to  $4f$  electrons is used. In addition, the  $5s^2 5p_{1/2}^2 5p_{3/2}^4 5d_{3/2}^4 5d_{5/2}^6$  shells of lead and the  $1s^2$  shell of oxygen were frozen (see [10] for details) and the residual ten electrons were treated explicitly in the subsequent molecular calculations. The basis set on Pb ( $15s16p12d9f$ )/[ $5s7p4d2f$ ] as well as GRECP are the same as those used in paper [9]. The basis set was optimized for calculation of properties determined mainly by the electronic wave function at the Pb core. The description of the basis set generation procedure can be found in Refs. [15, 29]. The correlation-consistent ( $10s5p2d1f$ )/[ $4s3p2d1f$ ] basis of Dunning listed in the MOLCAS 4.1 library [30] was used for oxygen.

The leading  $\Lambda\Sigma$  coupling terms and configurations for the  $a(1)$  and  $B(1)$  states are  $^3\Sigma^+ \sigma_1^2 \sigma_2^2 \sigma_3^2 \pi_1^3 \pi_2^1$  and

$^3\Pi_1 \sigma_1^2 \sigma_2^2 \sigma_3^1 \pi_1^4 \pi_2^1$ , respectively. The molecular orbitals used in the CI calculations are obtained by the restricted active space self-consistent field (RASSCF) method [30, 31] with the spin-averaged part of the GRECP (AGREP) [10], i.e. only scalar-relativistic effects are taken into account in the RASSCF calculation. Because the  $a(1)$  state is of primary interest, molecular orbitals were generated specifically for the lowest  $^3\Sigma^+$  state. This set of orbitals was used for the subsequent CI calculations of both the  $a(1)$  and  $B(1)$  states. In the RASSCF method, orbitals are divided into three active subspaces: RAS1, with a restricted number of holes allowed; RAS2, where all possible occupations are included; and RAS3, with an upper limit on the number of electrons. In this calculation, no more than two holes in RAS1 and two electrons in RAS3 are allowed. Using the  $C_{2v}$  point group classification scheme, two  $A_1$  orbitals in RAS1, six orbitals in RAS2, (two each in  $A_1$ ,  $B_1$ , and  $B_2$  irreps) and 50 (20  $A_1$ , 6  $A_2$ , 12  $B_1$ , and 12  $B_2$ ) in RAS3 subspaces are included.

Next the spin-orbit CI approach with the selected single- and double-excitations from some multiconfigurational reference states (“mains”) [32] is employed on the sets of different  $\Lambda\Sigma$  many-electron spin- and space-symmetry adapted basis functions (SAFs). In the SODCI code, the double  $C_{2v}$  group,  $C_{2v}^*$ , is used to account for the spin and space symmetry of the PbO molecule, instead of the more restrictive symmetry group  $C_{\infty v}^*$ , which could in principle be employed. In the  $C_{2v}^*$  classification scheme, the doubly-degenerate  $a(1)$  and  $B(1)$  states have the components in the  $B_1^*$  and  $B_2^*$  irreducible representations (irreps). The operators  $W_d$  and  $A_{\parallel}$  have nonzero matrix elements only between the wavefunction components from  $B_1^*$  and  $B_2^*$  irreps. So, one must know both these components of the  $a(1)$  and  $B(1)$  states to calculate  $W_d$  and  $A_{\parallel}$  when working in the  $C_{2v}^*$  group. The SAFs from the  $^{2S+1}C_{2v}$ -irreps (constructed on the basis of AGREP/RASSCF pseudoorbitals), singlet ( $^1B_1$ ), triplets ( $^3A_1$ ,  $^3A_2$ ,  $^3B_2$ ) and quintets ( $^5A_1$ ,  $^5A_2$ ,  $^5B_1$ ,  $^5B_2$ ), were included in calculations of the components belonging to the  $B_1^*$  irrep and equivalent calculations were performed for those laying in the  $B_2^*$  irrep. (Alternatively, the components belonging to the  $B_2^*$  irrep can be constructed by acting on those from the  $B_1^*$  irrep by the operator of projection of the total electronic momentum on the molecular axis.) The reference space,  $\{\Phi_I^{(0)}\}_{I=1}^{N_0}$ , consisted of  $N_0 = 2517$  SAFs having the largest coefficients in the probing CI calculation. The single and double excitations from this reference space produce about  $N_{T=0} \approx 175\,000\,000$  SAFs,  $\{\Phi_I^{(1,2)}\}_{I=1}^{N_{T=0}}$ . The molecular Hamiltonian  $\mathbf{H}$  in our calculations is presented as

$$\mathbf{H} = \mathbf{H}^0 + V, \quad (4)$$

where  $\mathbf{H}^0$  is an unperturbed spin-averaged Hamiltonian constructed to be diagonal in the given many-electron

basis set:

$$\mathbf{H}^0 \Phi_I^{(0,1,2)} = E_I^{(0,1,2)} \Phi_I^{(0,1,2)}, \quad (5)$$

and  $\mathbf{V}$  is a perturbation that includes the two-electron operator describing correlations and a one-electron effective spin-orbit operator. Only the most important SAFs from  $\{\Phi_I^{(1,2)}\}$  set, which give contribution in the second-order perturbation theory *by energy* greater than some chosen threshold  $T_i$ :

$$\frac{|\langle \Phi_J^{(1,2)} | \mathbf{V} | \Psi_X^0 \rangle|^2}{E_J^{(1,2)} - \mathcal{E}_X^0} = \delta E_{XJ} \geq T_i, \quad (6)$$

were selected (with all SAFs from the  $\{\Phi_I^{(0)}\}$  set) for the subsequent CI calculation. In the above equation,  $\mathcal{E}_X^0$  and  $\Psi_X^0$  are the eigenvalue and eigenfunction of  $\mathbf{H}$  for  $a(1)$  or  $B(1)$  state in the subspace  $\{\Phi_I^{(0)}\}$ . About 120 000, 500 000, 1 100 000, and 2 000 000 SAFs were selected (see Table I) for the thresholds (in  $10^{-6}$  a.u.)  $T_1=0.1$ ,  $T_2=0.01$ ,  $T_3=0.0025$  and  $T_4=0.001$ , respectively. (Selection on the CI *coefficients* obtained with the first-order perturbation theory [18] was also used in our calculations of  $A_{||}$  and  $W_d$  but the obtained results for the same number of selected SAFs were very close to those obtained using (6) and they are not presented in the paper.) The final wavefunction is written as

$$\Psi_X^{T_i} = \sum_{n=0}^2 \sum_J C_{XJ}^{T_i(n)} \Phi_J^{(n)}, \quad (7)$$

where summation is performed over selected SAFs for a given threshold  $T_i$  and those from the  $\{\Phi_I^{(0)}\}$  set. With this wavefunction, energy and any property  $W$  can be easily calculated:

$$E_X^{T_n} = \langle \Psi_X^{T_n} | \mathbf{H} | \Psi_X^{T_n} \rangle, \quad (8)$$

$$W_X^{T_n} = \langle \Psi_X^{T_n} | \mathbf{W} | \Psi_X^{T_n} \rangle. \quad (9)$$

Then, the linear  $T=0$  [22] and generalized Davidson (full-CI or FCI) [23] corrections are applied to the calculated properties. The  $T=0$  correction extrapolates the results of calculations with two different thresholds  $T_i$  and  $T_j$  to a result of the calculation with all singly- and doubly-excited SAFs included. The full-CI correction approximates the effect of all possible excitations of higher level (for a given number of correlated electrons and fixed basis set). The  $T=0$  correction for energy is given by

$$E_X^{T=0} = E_X^{T_n} + \lambda P_X^{T_n} \Rightarrow \lambda = -(E_X^{T_i} - E_X^{T_j}) / (P_X^{T_i} - P_X^{T_j}), \quad (10)$$

where  $n = i, j$  and

$$P_X^{T_n} = \sum_{J: \delta E_{XJ} < T_n} \delta E_{XJ}. \quad (11)$$

The full-CI corrected energy is given by

$$E_X^{FCI} \approx E_X^{T=0} + (1 - |c_I^{(0)}|^2) \cdot (E_X^{T=0} - \mathcal{E}_X^0), \quad (12)$$

where  $|c_I^{(0)}|^2$  are weights of the reference SAFs in  $\Psi_X^{T_{\min}}$ ,  $T_{\min} = \min(T_i, T_j)$ . If all singly-excited SAFs,  $\{\Phi_I^{(1)}\}$ , are included in calculation and the number of the reference SAFs is large enough, then similar expressions for  $T=0$  and FCI corrections (in more details they will be discussed in [24]) for the *one-electron* property  $W$  can be written:

$$W_X^{T=0} = W_X^{T_n} + \lambda P_X^{T_n} \Rightarrow \lambda = -(W_X^{T_i} - W_X^{T_j}) / (P_X^{T_i} - P_X^{T_j}), \quad (13)$$

where  $n = i, j$ ,

$$W_X^{FCI} \approx W_X^{T=0} + (1 - |c_I^{(0)}|^2) \cdot (W_X^{T=0} - \mathcal{W}_X^0), \quad (14)$$

where  $\mathcal{W}_X^0 = \langle \Psi_X^0 | \mathbf{W} | \Psi_X^0 \rangle$ . Though equations (13) and (14) look similar to (10) and (12), they have essentially different physical sense. More details about the features of constructing the reference space and selection procedure are given in Refs. [18, 24].

Before calculating  $A_{||}$  and  $W_d$ , the shapes of the four-component molecular spinors are restored in the inner core region after the two-component GRECP calculation of the molecule. For this purpose the NOCR method [12] is applied at the *second step* of the calculation. (See [21] for the currently used formulation of the NOCR scheme and [9] for details of restoration of the four-component molecular spinors near the Pb nucleus).

We designate the  $5s, 5p, 5d$  orbitals of lead and  $1s$  orbital of oxygen as the "outercore", and the  $\sigma_1, \sigma_2, \sigma_3, \pi_1, \pi_2$  orbitals of PbO (consisting mainly of  $6s, 6p$  orbitals of Pb and  $2s, 2p$  orbitals of O) as valence. In the CI calculations we take into account only the correlations between valence electrons. At the final stage of our calculation, we estimate the contribution from correlations between valence and outercore electrons (including high-order correlations between outercore electrons) as difference in the results of the corresponding 30- and 10-electron GRECP/RCC calculations. (See also [15] where this correction is applied to the Pb atom). Such correlations are designated in the text and tables as "outercore correlations".

*Results and discussion.* Calculations were performed at two internuclear distances,  $R = 3.8$  a.u. (as in the RCC calculations of [9]), and  $R = 4.0$  a.u. The latter

point is closer to the equilibrium distances both for  $a(1)$  ( $R_e = 4.06$  a.u. [33]) and for  $B(1)$  ( $R_e = 3.914$  a.u. [34]). However, in the RCC calculations [9],  $R = 3.8$  a.u. was used because of a problem with convergence at  $R = 4.0$  a.u. The calculated values with the one-center expansion of the molecular spinors in the Pb core on either  $s$ ,  $s;p$  or  $s;p;d$  partial waves are collected in Table I. The final data are obtained as the result of linear extrapolation to the experimental equilibrium distances.

As was noted in [9] when comparing the RASSCF results with those obtained by the RCC method, the spin-orbit interaction changes  $A_{||}$  and  $W_d$  dramatically, so even the sign for  $W_d$  is changed. One can see a similar picture from comparison of the RASSCF and CI data obtained in the present work. The second point to notice is a significant difference of the results obtained at the internuclear distances  $R = 3.8$  and  $R = 4.0$  a.u., especially, for the  $W_d$  parameter. It is increasing for the  $a(1)$  state and is decreasing for the  $B(1)$  state, by 15% and 30%, respectively. The agreement with the experimental datum attained in the CI calculation of  $A_{||}$  for the  $a(1)$  state at the point  $R = 4.0$  with the threshold  $T_4=0.001$  is 13%. After applying  $T=0$  and FCI corrections and taking into account the outercore correlations the agreement is improved to the level of 9%. The calculated  $A_{||}$  value for the  $B(1)$  state coincides with the measured datum within the experimental uncertainty of 4%. The results with the  $T=0$  correction are close to those with the smallest threshold  $T_4=0.001$ . To check the reliability of the linear  $T=0$  correction [22] for  $A_{||}$  and  $W_d$  we have calculated it for three different pairs of thresholds:  $T_1$  and  $T_2$ ,  $T_2$  and  $T_3$ ,  $T_3$  and  $T_4$ . For  $A_{||}$  all three pairs give the same result within 1% of accuracy. As to  $W_d$ , the result with the  $T=0$  correction for the first pair differs from those for the other two pairs by 8% for  $a(1)$  and by 5% for  $B(1)$ . However, the last two pairs again give the same result within the accuracy of 1% for both states. So, we suggest, that the  $T=0$  limit is determined for our main configurations with an accuracy of 1% for both considered properties. Because the reference space is large enough, we have a small FCI correction. When taking into account outercore contributions at the point  $R = 4.0$  a.u. we used the results of the RCC calculation at the point  $R = 3.8$  a.u. This assumption seems reasonable for several reasons. First, the core should relax less than the valence region when  $R$  is changed. In addition, because of the spatial separation between the core and valence electrons, core-valence correlation contributions should be more stable than valence-valence ones. Finally, since these contributions are relatively small, we expect errors due to this approximation not to be severe.

We next discuss the uncertainty in the calculated  $W_d$  values. (We confine this detailed discussion to the  $a(1)$  state; similar considerations are applied to  $B(1)$ .) Since  $W_d$  is sensitive to the wavefunction and its derivative at the Pb nucleus, it is natural to use the value of  $A_{||}$  (which

is also singular close to the Pb nucleus) as a benchmark for accuracy and stability. Thus, the 10% deviation of the calculated value of  $A_{||}$  from the experimental value represents an obvious lower bound for the accuracy of  $W_d$ . It appears that  $W_d$  is less computationally stable than  $A_{||}$ , however, as may be noted from the variation in values at various stages of the calculation, shown in Table I. Thus, this simple argument may underestimate the error in  $W_d$ . Because of the good convergence of the CI calculation, it appears that the deviation in  $A_{||}$  can be due in part to sensitivity of the results to the value of  $R$ , and in part to incomplete accounting for outercore-valence correlations. Consideration of the former effect leads us to a possible error in  $W_d$  of  $\sim 10\%$ . The latter is more difficult to account for numerically, but (based on experience from similar calculations on HgF, PbF, YbF, and BaF [14, 20, 27, 35]) we believe that the apparent insensitivity of  $A_{||}$  to outercore-valence correlations (compared to the case for  $W_d$ ) is an artifact unlikely to persist in more advanced calculations. Finally, we note that every improvement to the calculation (increasing correlation threshold; inclusion of core-valence correlation; increasing value of  $R$  towards the equilibrium value) actually increases  $|W_d|$ . Thus, the true value of  $|W_d|$  is not likely to be much lower than the present calculated value. The linear extrapolation of the calculated  $A_{||}$  and  $W_d$  to the experimental equilibrium distance for  $a(1)$ ,  $R_e = 4.06$  a.u. [33], gives us  $-3826$  MHz and  $-6.1 \cdot 10^{24}$  Hz/(e · cm), respectively, and the deviation of the extrapolated  $A_{||}$  value from the experiment now is only 7%. Accounting for the above arguments, the estimated error bounds put the actual  $W_d$  value between 90% and 130% of our final value. We obtain  $W_d$  for  $B(1)$  by linear interpolation of the data obtained at  $R = 3.8$  and  $4.0$  a.u. to the experimental equilibrium distance,  $R_e = 3.914$  a.u. [34], yielding  $W_d = -8.0 \cdot 10^{24}$  Hz/(e · cm), with an estimated error range of  $\pm 20\%$ . Similar interpolation for  $A_{||}$  on  $B(1)$  gives 4887 MHz, which is within the uncertainty of the current experimental value [36],  $5000 \pm 200$  MHz.

*Conclusion.* A new combined method for accurate calculating core properties in excited states of heavy-atom molecules is developed. It has allowed us to obtain accurate  $A_{||}$  and  $W_d$  values for the  $a(1)$  and  $B(1)$  states of PbO in spite of their complexity. The developed method and codes open an opportunity of reliable theoretical study of whole multitude of core properties in such computationally challenging objects as low-lying states of heavy-atom systems with valence  $p$ -electrons and dense spectra.

*Acknowledgments.* The present work is supported by U.S. CRDF grant RP2-2339-GA-02 and RFBR grant 03-03-32335. A.P. is grateful to the Ministry of Education of the Russian Federation (grant PD02-1.3-236). T.I. thanks INTAS for grant YSF 2001/2-164. N.M. was supported by the grants of Russian Science Support Foundation and the governor of Leningrad district. D.D. ac-

TABLE I: Calculated parameters  $A_{\parallel}$  (in MHz) and  $W_d$  (in  $10^{24}\text{Hz}/(e \cdot \text{cm})$ ) for the  $a(1)$  and  $B(1)$  states of  $^{207}\text{PbO}$  at internuclear distances 3.8 and 4.0 a.u. The experimental values of  $A_{\parallel}$  are  $-4113$  MHz in  $a(1)$  [37], and  $5000 \pm 200$  MHz in  $B(1)$  [36].

State Parameters Expansion	$a(1)$ $\sigma_1^2\sigma_2^2\sigma_3^2\pi_1^3\pi_2^1$ $^3\Sigma_1$					$B(1)$ $\sigma_1^2\sigma_2^2\sigma_3^1\pi_1^4\pi_2^1$ $^3\Pi_1$				
	$A_{\parallel}$			$W_d$		$A_{\parallel}$			$W_d$	
	s	s,p	s,p,d	s,p	s,p,d	s	s,p	s,p,d	s,p	s,p,d
(T is in $10^{-6}$ a.u.)										
<b>Internuclear distance <math>R = 3.8</math> a.u.</b>										
10e-RASSCF	-894	-1505	-1503	0.73	0.70				0.0	0.0
10e-RCC-SD [9]			-2635	-2.93	-3.05			3878	-11.10	-10.10
30e-RCC-SD [9]			-2698		-4.10			4081	-9.10	-9.70
<i>outercore (30e-RCC-SD - 10e-RCC-SD)</i>			<i>-63</i>		<i>-1.05</i>			<i>203</i>		<i>0.40</i>
10e-CI (reference) <sup>a</sup>	-406	-1877	-1874	-0.74	-0.83	731	3785	3805	-7.67	-7.17
10e-CI ( $T=0.1$ )	-472	-2930	-2926	-2.12	-2.21	393	4051	4074	-9.85	-9.40
10e-CI ( $T=0.01$ )	-430	-3222	-3218	-3.03	-3.13	371	4320	4344	-10.16	-9.72
10e-CI ( $T=0.0025$ )	-412	-3304	-3300	-3.44	-3.54	359	4411	4436	-10.46	-10.02
10e-CI ( $T=0.0012$ )	-407	-3332	-3328	-3.58	-3.69	359	4449	4474	-10.62	-10.18
10e-CI + $T=0$			-3387		-4.01			4555		-10.52
<i>10e-CI + <math>T=0</math> + FCI</i>			<i>-3446</i>		<i>-4.13</i>			<i>4582</i>		<i>-10.64</i>
<b>FINAL</b> (10e-CI + $T=0$ + FCI + outercore)			<b>-3509</b>		<b>-5.18</b>			<b>4785</b>		<b>-10.24</b>
(T is in $10^{-6}$ a.u.)										
<b>Internuclear distance <math>R = 4.0</math> a.u.</b>										
10e-RASSCF	-770	-1384	-1383	1.05	1.00				0.0	0.0
10e-CI (reference)	-459	-2026	-2025	-0.64	-0.72	966	4127	4150	-6.69	-6.22
10e-CI ( $T=0.1$ )	-479	-3125	-3124	-2.34	-2.44	525	4332	4357	-7.79	-7.35
10e-CI ( $T=0.01$ )	-449	-3458	-3458	-3.50	-3.61	495	4565	4590	-7.38	-6.94
10e-CI ( $T=0.0025$ )	-426	-3536	-3536	-3.97	-4.08	481	4636	4662	-7.45	-7.02
10e-CI ( $T=0.001$ )	-422	-3571	-3571	-4.19	-4.31	480	4666	4692	-7.49	-7.07
10e-CI + $T=0$			-3625		-4.65			4739		-7.15
<i>10e-CI + <math>T=0</math> + FCI</i>			<i>-3689</i>		<i>-4.81</i>			<i>4762</i>		<i>-7.18</i>
<b>FINAL</b> (10e-CI + $T=0$ + FCI + outercore) <sup>b</sup>			<b>-3752</b>		<b>-5.86</b>			<b>4965</b>		<b>-6.78</b>

<sup>a</sup> “Reference” means that the CI calculation was performed with the reference (main) SAFs only.

<sup>b</sup> It is assumed that the outercore contribution at the internuclear distance  $R = 4.0$  a.u. is approximately the same as is at the point  $R = 3.8$  a.u.

knowledges support from NSF grant PHY0244927 and the David and Lucile Packard Foundation.

\* Electronic address: anpetrov@pnpi.spb.ru

- [1] B. C. Regan, E. D. Commins, C. J. Schmidt, and D. DeMille, Phys. Rev. Lett. **88**, 071805/1 (2002).
- [2] O. P. Sushkov and V. V. Flambaum, Sov. Phys.-JETP **48**, 608 (1978).
- [3] V. G. Gorshkov, L. N. Labzovsky, and A. N. Moskalyov, Sov. Phys.-JETP **49**, 209 (1979).
- [4] J. J. Hudson, B. E. Sauer, M. R. Tarbutt, and E. A. Hinds, Phys. Rev. Lett. **89**, 023003/1 (2002).
- [5] D. DeMille, F. Bay, S. Bickman, D. Kwall, D. Krause, Jr., S. E. Maxwell, and L. R. Hunter, Phys. Rev. A **61**, 052507/1 (2000).
- [6] D. Egorov, J. D. Weinstein, D. Patterson, B. Friedrich, and J. M. Doyle, Phys. Rev. A **63**, 030501(R)/1 (2001).
- [7] M. G. Kozlov and D. DeMille, Phys. Rev. Lett. **89**, 133001 (2002).
- [8] U. Kaldor, E. Eliav, and A. Landau, in *Recent Advances in Relativistic Molecular Theory*, edited by K. Hirao and Y. Ishikawa (World Scientific, Singapore, 2004), pp. 283–327.
- [9] T. A. Isaev, A. N. Petrov, N. S. Mosyagin, A. V. Titov, E. Eliav, and U. Kaldor, Phys. Rev. A **69**, 030501(R)/1 (2004).
- [10] A. V. Titov and N. S. Mosyagin, Int. J. Quantum Chem. **71**, 359 (1999).
- [11] A. V. Titov and N. S. Mosyagin, Russ. J. Phys. Chem. **74**, Suppl. 2, S376 (2000), [arXiv: physics/0008160].
- [12] A. V. Titov, Ph.D. thesis, St.-Petersburg (Leningrad) State University, Russia (1985).
- [13] A. V. Titov, Int. J. Quantum Chem. **57**, 453 (1996).
- [14] A. V. Titov, N. S. Mosyagin, and V. F. Ezhov, Phys. Rev. Lett. **77**, 5346 (1996).
- [15] T. A. Isaev, N. S. Mosyagin, M. G. Kozlov, A. V. Titov, E. Eliav, and U. Kaldor, J. Phys. B **33**, 5139 (2000).
- [16] R. J. Buenker and S. Krebs, in *Recent Advances in Multireference Methods*, edited by K. Hirao (World Scientific, Singapore, 1999), pp. 1–29.
- [17] A. B. Alekseyev, H.-P. Liebermann, and R. J. Buenker, in *Recent Advances in Relativistic Molecular Theory*, edited

- by K. Hirao and Y. Ishikawa (World Scientific, Singapore, 2004), pp. 65–105.
- [18] A. V. Titov, N. S. Mosyagin, A. B. Alekseyev, and R. J. Buenker, *Int. J. Quantum Chem.* **81**, 409 (2001).
- [19] N. S. Mosyagin, A. V. Titov, R. J. Buenker, H.-P. Liebermann, and A. B. Alekseyev, *Int. J. Quantum Chem.* **88**, 681 (2002).
- [20] N. S. Mosyagin, M. G. Kozlov, and A. V. Titov, *J. Phys. B* **31**, L763 (1998).
- [21] A. N. Petrov, N. S. Mosyagin, T. A. Isaev, A. V. Titov, V. F. Ezhov, E. Eliav, and U. Kaldor, *Phys. Rev. Lett.* **88**, 073001/1 (2002).
- [22] R. J. Buenker and S. D. Peyerimhoff, *Theor. Chim. Acta* **39**, 217 (1975).
- [23] P. J. Bruna, S. D. Peyerimhoff, and R. J. Buenker, *Chem. Phys. Lett.* **72**, 278 (1980).
- [24] A. V. Titov et al., to be published.
- [25] M. G. Kozlov, V. I. Fomichev, Y. Y. Dmitriev, L. N. Labzovsky, and A. V. Titov, *J. Phys. B* **20**, 4939 (1987).
- [26] M. Kozlov and L. Labzovsky, *J. Phys. B* **28**, 1931 (1995).
- [27] Y. Y. Dmitriev, Y. G. Khait, M. G. Kozlov, L. N. Labzovsky, A. O. Mitrushenkov, A. V. Shtoff, and A. V. Titov, *Phys. Lett. A* **167**, 280 (1992).
- [28] N. S. Mosyagin, A. V. Titov, and Z. Latajka, *Int. J. Quantum Chem.* **63**, 1107 (1997).
- [29] N. S. Mosyagin, E. Eliav, A. V. Titov, and U. Kaldor, *J. Phys. B* **33**, 667 (2000).
- [30] K. Andersson, M. R. A. Blomberg, M. P. Fülscher, G. Karlström, R. Lindh, P.-A. Malmqvist, P. Neogrady, J. Olsen, B. O. Roos, A. J. Sadlej, et al. (1999), quantum-chemical program package “MOLCAS”, Version 4.1.
- [31] J. Olsen and B. O. Roos, *J. Chem. Phys.* **89**, 2185 (1988).
- [32] R. J. Buenker and S. D. Peyerimhoff, *Theor. Chim. Acta* **35**, 33 (1974).
- [33] F. Martin, R. Bacis, J. Verges, J. Bachar, and S. Rosenwaks, *Spectrochim. Acta* **44A**, 889 (1988).
- [34] K. P. Huber and G. Herzberg, *Constants of Diatomic Molecules* (Van Nostrand-Reinhold, New York, 1979).
- [35] M. G. Kozlov, A. V. Titov, N. S. Mosyagin, and P. V. Souchko, *Phys. Rev. A* **56**, R3326 (1997).
- [36] D. Kawai, Y. Gurevich, and D. DeMille, to be published.
- [37] L. R. Hunter, S. E. Maxwell, K. A. Ulmer, N. D. Charney, S. K. Peck, D. Krause, S. Ter-Avetisyan, and D. DeMille, *Phys. Rev. A* **65**, 030501(R) (2002).