

# Linear media in classical electrodynamics and the Post constraint

Friedrich W. Hehl<sup>1,2</sup> and Yuri N. Obukhov<sup>1,3</sup>

<sup>1</sup> Institut für Theoretische Physik, Universität zu Köln  
50923 Köln, Germany

<sup>2</sup> Department of Physics and Astronomy, University of Missouri-Columbia  
Columbia, MO 65211, USA

<sup>3</sup> Department of Theoretical Physics, Moscow State University  
117234 Moscow, Russia

3 November 2004, *file PostCon9.tex*

## Abstract

The Maxwell equations are formulated in a generally covariant and metric-free way in 1+3 and subsequently in 4 dimensions. For this purpose, we use the excitations  $\mathcal{D}$ ,  $\mathcal{H}$  and the field strengths  $E$ ,  $B$ . A local and linear constitutive law between excitations and field strengths is assumed, with a constitutive tensor  $\chi^{ijkl} = -\chi^{jikl} = -\chi^{ijlk}$  of 36 components. The properties of this tensor are discussed. In particular, we address the validity of the Post constraint, a subject that is very much under discussion. In this connection, the Tellegen gyrator, the axion field, and the “perfect electromagnetic conductor” of Lindell & Sihvola are compared with each other.

Keywords: Electrodynamics, linear medium, constitutive tensor, Post constraint, axion field, Tellegen gyrator, skewon field, classical field theory

PACS: 03.50.De, 46.05.+b, 14.80.Mz

## 1 Introduction

Electrical engineers as well as physicists are familiar with the Maxwell equations

$$\underline{d}\mathcal{D} = \rho, \quad \underline{d}\mathcal{H} - \dot{\mathcal{D}} = j, \quad (1)$$

$$\underline{d}B = 0, \quad \underline{d}E + \dot{B} = 0, \quad (2)$$

here expressed [1, 4, 5, 19, 39] in exterior differential forms in terms of the electric and magnetic excitations  $\mathcal{D}$ ,  $\mathcal{H}$  and the electric and magnetic field strength  $E$ ,  $B$ , respectively. With  $\underline{d}$  we denote

the 3-dimensional exterior derivative and with a dot the time derivative. The sources of the inhomogeneous equations are the electric charge density  $\rho$  and the electric current density  $j$ . These equations are expressed in a generally covariant form, that is, they are valid in *arbitrary curvilinear coordinates*. Moreover, they are *metric-free* and thus the concept of orthogonality is not used at all.

As shown in the literature, see [40, 34, 5], the Maxwell equations can alternatively be written in terms of components as

$$\partial_a \mathcal{D}^a = \rho, \quad \epsilon^{abc} \partial_b \mathcal{H}_c - \dot{\mathcal{D}}^a = j^a, \quad (3)$$

$$\partial_a B^a = 0, \quad \epsilon^{abc} \partial_b E_c + \dot{B}^a = 0, \quad (4)$$

with  $a, b, \dots = 1, 2, 3$  and  $\epsilon^{abc} = \pm 1, 0$  as the totally antisymmetric Levi-Civita symbol. Here  $\mathcal{D}^a, B^a, j^a$  are vector densities,  $E_a, \mathcal{H}_a$  covectors, and  $\rho$  is as scalar density. Also in this formulation the general covariance and the condition of being free of the metric are manifest. Note, since no metric is used, we must not raise and lower indices with the help of the metric at this stage.

The Maxwell equations need to be supplemented by constitutive relations. And this for the description (i) of empty spacetime (the vacuum) and (ii) of material media. We will discuss here the simplest case, namely local and linear media. Local in space as well as in time, that is, the excitations at every point in space and time depend only on the field strength at the very same point.

A general local and linear constitutive relation reads (in the conventions of [5]),

$$\mathcal{H}_a = -\mathfrak{C}^b_a E_b + \mathfrak{B}_{ba} B^b, \quad (5)$$

$$\mathcal{D}^a = -\mathfrak{A}^{ba} E_b + \mathfrak{D}_b^a B^b. \quad (6)$$

The generalized permittivity matrix  $\mathfrak{A}^{ba}$  and the generalized impermeability matrix (the reciprocal of the permeability)  $\mathfrak{B}_{ba}$  are well known. Less known, but well established in the electrodynamics of continuous media, see [31], are the magneto-electric matrices  $\mathfrak{C}^b_a$  and  $\mathfrak{D}_b^a$ , which describe magnetic-electric cross effects (like the Faraday effect or optical activity). Altogether, these matrices encompass  $4 \times 9 = 36$  independent components. Since  $\mathcal{H}_a, \mathcal{D}^a$  and  $E_b, B^b$  are real, the matrices  $\mathfrak{A}^{ba}$  etc. are also real. The subject of our article will be the discussion of certain possible algebraic constraints between the components of the constitutive matrices  $\mathfrak{A}^{ba}, \mathfrak{B}_{ba}, \mathfrak{C}^b_a, \mathfrak{D}_b^a$ . The most notorious one is the so-called Post constraint, which requires the sum of the traces of the matrices  $\mathfrak{C}^b_a$  and  $\mathfrak{D}_b^a$  to vanish, i.e.,  $\mathfrak{C}^a_a + \mathfrak{D}_a^a = 0$ . There is an ongoing dispute mainly in the electrical engineering community, see Lakhtakia & Weiglhofer [12, 13, 14, 15, 16, 50, 51] and Lindell, Sihvola, Tretyakov, et al. [20, 21, 22, 23, 24, 38, 43, 49] whether this constraint is valid or not. We will address this question.

The constitutive matrices are functions of space  $x^a$  and time  $t$  in general. In other words, they are *fields*  $\mathfrak{A}^{ba} = \mathfrak{A}^{ba}(x^a, t)$ , etc. For applications to the case (i), to empty spacetime, they can be constants if a flat Minkowski spacetime and Cartesian coordinates are prescribed. However, if a Riemannian spacetime of general relativity is assumed or just a 4-dimensional differential manifold, then clearly the constitutive matrices are fields. Similarly, in case (ii), for material media, a homogeneous medium requires constant constitutive matrices, as soon as inhomogeneous media are allowed, the constitutive matrices are fields again. We may call them generalized permittivity/impermeability fields.

Before we turn to a 4-dimensional discussion of the Maxwell equations, we want to comment on the constitutive relations (5),(6). In the next section we will show that for reasons of 4-dimensional

covariance (5),(6) represent the appropriate constitutive relations. However, in the engineering as well as in the physics literature, we often find  $\mathcal{D}^a$  and  $B^a$  expressed in terms of  $E_b$  and  $H_b$ . If we exclude singular cases, i.e.,

$$\det \mathfrak{B} \neq 0, \quad (7)$$

then (5) can be multiplied with  $\mathfrak{B}^{-1}$ , solved with respect to  $B^a$ , and substituted into (6). In this way, we find, suppressing indices,

$$\mathcal{D} = (-\mathfrak{A} + \mathfrak{D}\mathfrak{B}^{-1}\mathfrak{C}) E + (\mathfrak{D}\mathfrak{B}^{-1}) \mathcal{H}, \quad (8)$$

$$B = (\mathfrak{B}^{-1}\mathfrak{C}) E + (\mathfrak{B}^{-1}) \mathcal{H}. \quad (9)$$

Thus, the the  $EB$ -formalism in (5),(6) is transformed to the  $E\mathcal{H}$ -formalism of (8),(9). Both formalisms are equivalent, provided a singular case can be excluded. The natural linear constitute relations are represented by (5),(6), as was shown by Post [34] and O'Dell [31], e.g., and as we will discuss in the next section. Hence we will concentrate on (5),(6) and will consider (8),(9) only as secondary.

## 2 Electrodynamics in 4-dimensional spacetime

The true structure of classical electrodynamics can be more clearly recognized if we present it in a 4-dimensional way. If we define excitation, electric current, and field strength according to

$$H = \mathcal{D} - \mathcal{H} \wedge dt, \quad J = \rho - j \wedge dt, \quad (10)$$

$$F = B + E \wedge dt, \quad (11)$$

then the 4-dimensional Maxwell equations (1) and (2) read

$$dH = J, \quad (12)$$

$$dF = 0. \quad (13)$$

This shows explicitly that they are invariant under 4-dimensional coordinate/frame transformations and independent of the metric of spacetime, see [5]. If we require metric independence, we get a well-prescribed way of formulating the Maxwell equations. Since the metric of spacetime is the gravitational potential in general relativity, it is gratifying to know that there is a gravity-free way of formulating the Maxwell equations. Accordingly, the Maxwell equations (12),(13) are valid in a flat Minkowski spacetime (in any coordinates), in a curved Riemannian spacetime, and even in the Riemann-Cartan spacetime of the Poincaré gauge theory of gravitation. Needless to say that this beautiful formalism has great practical value for applying electrodynamics in accelerated reference frames, for instance.

The excitation  $H$  is a *twisted* 2-form, the field strength  $F$  an *untwisted* 2-form, see the discussion of Post in [36]. We decompose them in their components according to  $H = H_{ij} dx^i \wedge dx^j / 2$  and  $F = F_{ij} dx^i \wedge dx^j / 2$ . Here  $x^i$  are spacetime coordinates with  $i, j, \dots = 0, 1, 2, 3$ . Analogously, the current  $J$  as a twisted 3-form decomposes as  $J = J_{ijk} dx^i \wedge dx^j \wedge dx^k / 6$ . Thus, in the formalism of tensor analysis, (12) and (13) can alternatively be written as

$$\partial_k \check{H}^{ik} = \check{J}^i, \quad (14)$$

$$\partial_{[i} F_{kl]} = 0, \quad (15)$$

with  $\check{H}^{ik} := \epsilon^{iklm} H_{lm}/2$  and  $\check{J}^i := \epsilon^{iklm} J_{klm}/6$ . Here  $\epsilon^{iklm} = \pm 1, 0$  is the 4-dimensional contravariant Levi-Civita symbol. The covariant Levi-Civita symbol, which we will use below, is denoted by a circumflex:  $\hat{\epsilon}_{iklm} = \pm 1, 0$ . Incidentally, since no metric is available for the raising and lowering of indices at this stage, we have to distinguish these two symbols.

In the constitutive law (5),(6), we assumed *locality* and *linearity*. In 4 dimensions this translates into

$$H = \kappa(F), \quad (16)$$

where  $\kappa$  is a local and linear operator. Accordingly,

$$H_{ij} = \frac{1}{2} \kappa_{ij}{}^{kl} F_{kl} \quad \text{with} \quad \kappa_{ij}{}^{kl} = -\kappa_{ji}{}^{kl} = -\kappa_{ij}{}^{lk}. \quad (17)$$

Here  $\kappa_{ij}{}^{kl}(x)$  is the twisted constitutive tensor of type  $\left[ \begin{smallmatrix} 2 \\ 2 \end{smallmatrix} \right]$ . It can characterize (i) either empty spacetime or (ii) a material medium. It has at most 36 independent components. In general, it is a field, that is, it can describe an inhomogeneous spacetime or an inhomogeneous medium.

The relativistic invariance properties are crucial for deciding between the  $EB$ - and  $E\mathcal{H}$ -formalisms. From the point of view of a 4-dimensional spacetime,  $E$  and  $B$  form the invariant field strength  $F$ ; likewise,  $\mathcal{D}$  and  $\mathcal{H}$  form the invariant excitation  $H$ . Accordingly, the linear constitutive law (5),(6) is a relativistically invariant statement, as it is seen from (16). This law involves the constitutive matrices  $\mathfrak{A}$ ,  $\mathfrak{B}$ ,  $\mathfrak{C}$ , and  $\mathfrak{D}$  that are combined into a covariant object – the constitutive tensor  $\kappa_{ij}{}^{kl}$ . In contrast, the pairs  $\mathcal{D}$ ,  $B$  and  $E$ ,  $\mathcal{H}$  do *not* form any 4-dimensional relativistic objects, and, as a result, the constitutive relation in the form of (8), (9) involves non-covariant objects with unclear physical meaning. Therefore, the  $EB$ -scheme of (5),(6) is, from a relativistic point of view, to be preferred against the  $E\mathcal{H}$ -scheme of (8),(9), see O’Dell [31], Chap.2, Sec.1.2, pp.23–24. If  $\det \mathfrak{B} \neq 0$ , both systems are formally equivalent; still, (8),(9), as a historical artifact, should be phased out from use.

One can decompose the constitutive tensor  $\kappa_{ij}{}^{kl}$  into its irreducible pieces under the linear group acting in the tangent and the cotangent space, respectively. Contraction is the only tool for such a decomposition. Following Post [35], we can define the contracted twisted tensor of type  $\left[ \begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]$

$$\kappa_i{}^k := \kappa_{il}{}^{kl}, \quad (18)$$

with 16 independent components. The second contraction yields the twisted scalar function

$$\kappa := \kappa_k{}^k = \kappa_{kl}{}^{kl} \quad (19)$$

(also called pseudo- or axial-scalar). The traceless piece

$$\not\kappa_i{}^k := \kappa_i{}^k - \frac{1}{4} \kappa \delta_i^k \quad (20)$$

has 15 independent components. These pieces (19) and (20) can now be subtracted from the original constitutive tensor. Then we find the following three irreducible pieces:

$$\begin{aligned} \kappa_{ij}{}^{kl} &= {}^{(1)}\kappa_{ij}{}^{kl} + {}^{(2)}\kappa_{ij}{}^{kl} + {}^{(3)}\kappa_{ij}{}^{kl} \\ &= {}^{(1)}\kappa_{ij}{}^{kl} + 2 \not\kappa_{[i}{}^{[k} \delta_{j]}^l] + \frac{1}{6} \kappa \delta_{[i}^k \delta_{j]}^l. \end{aligned} \quad (21)$$

By construction,  ${}^{(1)}\kappa_{ij}{}^{kl}$  is totally traceless:

$${}^{(1)}\kappa_{il}{}^{kl} = 0. \quad (22)$$

Thus, we split  $\kappa_{ij}{}^{kl}$  according to  $36 = 20 + 15 + 1$ , since the  ${}^{[2]}$  tensor  ${}^{(1)}\kappa_{ij}{}^{kl}$  is subject to the 16 constraints (22) and hence carries  $36 - 16 = 20$  components. We call  ${}^{(1)}\kappa_{ij}{}^{kl}$  the *principal* (or the metric-dilaton) part of the constitutive tensor. Without such a term, the propagation of electromagnetic waves is ruled out, see the discussion subsequent to eq.(53). We further identify the two other irreducible parts  ${}^{(2)}\kappa_{ij}{}^{kl}$  and  ${}^{(3)}\kappa_{ij}{}^{kl}$  as *skewon* and *axion* fields, respectively. Conventionally, the skewon and the axion fields are introduced with different factors by

$$\mathcal{S}_i{}^j := -\frac{1}{2} \not\kappa_i{}^j, \quad \alpha := \frac{1}{12} \kappa. \quad (23)$$

If we substitute (21) into (17)<sub>1</sub> and use (23), then the general linear constitutive law can be written as

$$H_{ij} = \frac{1}{2} {}^{(1)}\kappa_{ij}{}^{kl} F_{kl} + 2 \mathcal{S}_{[i}{}^k F_{j]k} + \alpha F_{ij}. \quad (24)$$

It has been shown recently [17] that taking the linear spacetime relation for granted, one can derive a Riemannian light cone provided one forbids birefringence in vacuum. However, if one allows for the skewon field, then the light cone is dissolved and we find birefringence and more general type of optical effects [30].

Let us now turn to some examples for case (i), the vacuum.

### Maxwell-Lorentz vacuum electrodynamics

If we assume a Riemannian metric  $g_{ij}$  with Lorentzian signature, then vacuum electrodynamics is recovered by

$${}^{(1)}\kappa_{ij}{}^{kl} = \lambda_0 \sqrt{-g} \hat{\epsilon}_{ijmn} g^{mk} g^{nl}, \quad \mathcal{S}_i{}^j = 0, \quad \alpha = 0. \quad (25)$$

Here  $\lambda_0 = \sqrt{\epsilon_0/\mu_0}$  is the vacuum admittance and  $g := \det g_{kl}$ . The principal part  ${}^{(1)}\kappa_{ij}{}^{kl}$  corresponds to the Hodge star operator  $\star$ . The Maxwell equations can then be written as<sup>1</sup>

$$\lambda_0 d^\star F = J, \quad dF = 0. \quad (26)$$

### Axion electrodynamics

If we admit additionally the axion field  $\alpha \neq 0$ , then  $H = \lambda_0 \star F + \alpha F$ . This framework is called axion (Maxwell-Lorentz) electrodynamics, see Ni [25, 26, 27] and Wilczek [52], e.g.:

$$\lambda_0 d^\star F + (d\alpha) \wedge F = J, \quad dF = 0. \quad (27)$$

It is as if the current  $J$  picked up an additional piece depending on the gradient of the axion field. The real part of Kiehn's *chiral vacuum* [9] is a subcase, for  $\alpha = \text{const}$ , of axion electrodynamics.

Decomposing the inhomogeneous equation in time and space, we have ( $c$  is the speed of light and  $\underline{\star}$  the 3-dimensional Hodge star)

$$\frac{\lambda_0}{c} \underline{d}^\star E = \rho - (\underline{d}\alpha) \wedge B, \quad (28)$$

$$\lambda_0 c \underline{d}^\star B - \frac{\lambda_0}{c} \partial_t^\star E = j + (\underline{d}\alpha) \wedge E + (\partial_t \alpha) B. \quad (29)$$

---

<sup>1</sup>The inhomogeneous equation in indices reads  $\lambda_0 \partial_j (\sqrt{-g} g^{ik} g^{jl} F_{kl}) = \check{J}^i$ .

Possible experimental verifications were suggested by Sikivie and others, see [44, 45, 32].

### Pure axion field and the gyrator

As a degenerate special case, we can also consider the pure (“stand-alone”) axion field with  ${}^{(1)}\kappa_{ij}{}^{kl} = {}^{(2)}\kappa_{ij}{}^{kl} = 0$ . Then,

$$H = \alpha F \quad \text{or} \quad \mathcal{H} = -\alpha E, \quad \mathcal{D} = \alpha B, \quad (30)$$

and the Maxwell equations read

$$(d\alpha) \wedge F = J \quad \text{and} \quad dF = 0. \quad (31)$$

This is a special case of axion electrodynamics, namely (27) with  $\lambda_0 = 0$ . Here we have a case in which the constitutive matrix  $\mathfrak{B}$  in (5) vanishes. Consequently,  $\det \mathfrak{B} = 0$  and the  $E\mathcal{H}$ -version (8),(9) of the constitutive law loses its meaning.

Seemingly the first person to discuss (and to reject) a constant pure axion field was Schrödinger [41], p.25, penultimate paragraph, and, as a non-constant field, Dicke [3]. The framework (30),(31) corresponds to Tellegen’s *gyrator* [46, 47] and to Lindell & Sihvola’s *perfect electromagnetic conductor* (PEMC) [20, 21].

Lindell & Sihvola define their PEMC by using directly the relation  $H = \alpha F$ . Tellegen [46, 47] considered a linear network of two terminal pairs (four-pole or two-port):

$$v_1 = a_{11} i_1 + a_{12} i_2, \quad (32)$$

$$v_2 = a_{21} i_1 + a_{22} i_2. \quad (33)$$

The  $v$ ’s represent voltages, the  $i$ ’s currents, and the (frequency dependent)  $a$ ’s resistances. Since the field strengths are related to the voltage ( $E \stackrel{\text{SI}}{\sim} V/m$ ,  $B \stackrel{\text{SI}}{\sim} Vs/m^2$ ) and the excitations to the current ( $\mathcal{H} \stackrel{\text{SI}}{\sim} A/m$ ,  $\mathcal{D} \stackrel{\text{SI}}{\sim} As/m^2$ ), our translation into the linear medium picture reads:

$$E = a_{11} \mathcal{D} + a_{12} \mathcal{H}, \quad (34)$$

$$B = a_{21} \mathcal{D} + a_{22} \mathcal{H}. \quad (35)$$

Tellegen had a different translation, since he used the  $E\mathcal{H}$ -formalism of (8),(9). Tellegen [46], eq.(10), defined his gyrator by

$$v_1 = -\sigma i_2, \quad (36)$$

$$v_2 = \sigma i_1. \quad (37)$$

The gyrator “rotates” currents into voltages and vice versa. In our medium picture we have

$$E = -\sigma \mathcal{H}, \quad (38)$$

$$B = \sigma \mathcal{D}, \quad (39)$$

i.e., the excitations are rotated into the field strengths. As a comparison with (30) shows, this corresponds to the axion field. Lindell & Sihvola [20] have shown that by using the  $E\mathcal{H}$ -formalism and by performing a suitable limiting transition (since  $\det \mathfrak{B} = 0$ ), one can also arrive at (38),(39). Consequently, it is established that the medium analogue of Tellegen’s gyrator is the axion field — and both are equivalent to the PEMC of Lindell & Sihvola.

The gyrator has been first realized for microwaves by the use of premagnetized ferrites. A corresponding figure with detailed explanations can be found in the review article of Hogan [6], Fig.12.

### 3 The constitutive tensor $\chi^{ijkl}$ and its properties

We can raise the two lower indices of the constitutive tensor  $\kappa_{ij}{}^{kl}$  with the help of the contravariant Levi–Civita symbol  $\epsilon^{ijmn}$ . Thus, we can introduce the constitutive tensor *density* of spacetime,

$$\chi^{ijkl} := \frac{1}{2} \epsilon^{ijmn} \kappa_{mn}{}^{kl}, \quad (40)$$

which is equivalent to  $\kappa_{ij}{}^{kl}$  but more widely in use. The notation  $\chi$  was chosen in order to conform with Post’s convention [34]. The constitutive law can now be written in the conventional form as

$$\check{H}^{ij} = \frac{1}{2} \chi^{ijkl} F_{kl}, \quad (41)$$

see Post [34], eq.(6.12). The constitutive tensor with all 36 components has also been discussed in an equivalent way by Lindell in his book [19], see also [42, 11, 43, 38, 23, 24, 20, 21].

We can decompose  $\chi^{ijkl}$  irreducibly under the linear group, too. With the expressions

$${}^{(1)}\chi^{ijkl} = \frac{1}{2} \epsilon^{ijmn} {}^{(1)}\kappa_{mn}{}^{kl}, \quad (42)$$

$${}^{(2)}\chi^{ijkl} = \frac{1}{2} \epsilon^{ijmn} {}^{(2)}\kappa_{mn}{}^{kl} = -\epsilon^{ijm[k} \kappa_m{}^{l]}, \quad (43)$$

$${}^{(3)}\chi^{ijkl} = \frac{1}{2} \epsilon^{ijmn} {}^{(3)}\kappa_{mn}{}^{kl} = \frac{1}{12} \epsilon^{ijkl} \kappa, \quad (44)$$

we find<sup>2</sup>

$$\underbrace{\chi^{ijkl}}_{36} = \underbrace{{}^{(1)}\chi^{ijkl}}_{\text{principal } 20} + \underbrace{{}^{(2)}\chi^{ijkl}}_{\text{skewon } 15} + \underbrace{{}^{(3)}\chi^{ijkl}}_{\text{axion } 1}. \quad (45)$$

The irreducible pieces carry the additional symmetries

$${}^{(1)}\chi^{ijkl} = {}^{(1)}\chi^{klij}, \quad {}^{(2)}\chi^{ijkl} = -{}^{(2)}\chi^{klij}, \quad {}^{(3)}\chi^{ijkl} = {}^{(3)}\chi^{[ijkl]}. \quad (46)$$

Clearly, the *principal* part with its 20 independent components is the one discussed by Post [34]. The *skewon* part with its 15 components vanishes if one assumes the existence of a Lagrangian 4-form from which the spacetime relation can be derived in full, see below. However, this assumption is not a necessary, as we have shown [5]. Finally, the *axion* piece with only 1 independent component is totally antisymmetric:

$${}^{(3)}\chi^{ijkl} = \alpha \epsilon^{ijkl}. \quad (47)$$

In Post’s book [34], eq.(6.18), this piece is forbidden; therefore, Lakhtakia and Weiglhofer [14] called it the Post constraint:

$${}^{(3)}\chi^{ijkl} = 0 \quad \text{or} \quad \alpha = 0 \quad (\text{Post constraint}). \quad (48)$$

---

<sup>2</sup>We want to underline that the irreducible decomposition of  $\chi^{ijkl}$  is *algebraically* similar the decomposition of a curvature tensor in a Riemann-Cartan space (Riemannian space with a metric compatible connection). Of course, only in a Riemann-Cartan space the curvature tensor has 36 independent components and can carry a totally antisymmetric piece  $R^{[ijkl]}$ . This piece, by the first Bianchi identity, is directly related to Cartan’s torsion. But let us stress, these are just algebraic analogies, not more and not less. Premetric electrodynamics, as represented by eqs.(1) and (2), does neither couple to the metric nor to the connection of spacetime. Thus also torsion is not involved. Moreover, the spacetime relation eventually defines a metric, but that’s all.

Post even carries on and requires  $\partial_i \alpha = 0$ , see his eq.(6.19). The reasons he gave for the latter constraint are incomprehensible to us. Therefore, we will not dwell on it any further.

We want now to turn to criteria that will help us to decide which pieces of the constitutive tensor are reasonable from a physical point of view. Our kinematical analysis so far gave no convincing argument why any of its 36 components should not exist in nature. In the sequel, we will distinguish between case (i), the vacuum, and case (ii), material media.

In electrodynamics, after formulating the Maxwell equations, one has to specify the energy-momentum density of the electromagnetic field. Also this can be done in a metric-free environment, see [5]. The *energy-momentum* 3-form in (holonomic) coordinates reads

$$\Sigma_i = \frac{1}{2} (F \wedge e_i \rfloor H - H \wedge e_i \rfloor F) , \quad (49)$$

with  $e_i = \partial_i$  as the 4-dimensional vector basis of the tangent space. Let us denote the basis of the 4-dimensional 3-forms by  $\hat{\epsilon}_k = \epsilon_{klmn} dx^l \wedge dx^m \wedge dx^n / 6$ . If we then decompose the energy-momentum 3-form with respect to  $\hat{\epsilon}_k$ , that is,  $\Sigma_i = \mathcal{T}_i^k \hat{\epsilon}_k$ , we find the conventional Minkowski energy-momentum tensor

$$\mathcal{T}_i^k = \frac{1}{4} \delta_i^k F_{lm} \check{H}^{lm} - F_{il} \check{H}^{kl} . \quad (50)$$

Of course, (49) and (50) are equivalent representations of the energy-momentum distribution.

The representation (49) is most convenient for showing that the axion piece, even as field, doesn't carry electromagnetic energy-momentum. The constitutive law  $H = \alpha(x)F$  yields  $F \wedge e_i \rfloor H = F \wedge e_i \rfloor (\alpha F) = \alpha F \wedge e_i \rfloor F$ . Together with the second piece in (49), this vanishes. For the principal piece it can be shown that it leads to positive energy, see [8], also the skewon piece has non-vanishing contributions in general, see Table 1. We are talking here about the *electromagnetic* energy. If the axion field contributed an own kinetic term à la  $\sim g^{ik} (\partial_i \alpha) (\partial_k \alpha) / 2$  to the Lagrangian, then it would carry own energy.

Table 1: Contribution of each irreducible part  ${}^{(r)}\chi$  of the constitutive tensor density to the electromagnetic energy-momentum, to the Lagrangian, and to the **T**amm-**R**ubilar tensor density, see [5]. Our results are consistent with the findings of Kiehn [9, 10].

Irreducible part	Contributes to		
	energy-momentum	Lagrangian	TR tensor density
principal ${}^{(1)}\chi$	yes	yes	yes
skewon ${}^{(2)}\chi$ or $\mathcal{G}$	yes	no	yes
axion ${}^{(3)}\chi$ or $\alpha$	no	yes	no

This brings us to the Lagrangian of the electromagnetic field. The Maxwell equations (12),(13) can be derived from electric charge and magnetic flux conservation and the energy-momentum current (49) via the Lorentz force density. A Lagrangian is not needed in this procedure. The same is true for the local and linear constitutive law (in the sense of irreversible thermodynamics).

However, one can find an electromagnetic *Lagrangian* 4-form

$$V = -\frac{1}{2} H \wedge F. \quad (51)$$

By substituting  $H = \kappa(F)$  into the Lagrangian, the piece  ${}^{(2)}\kappa(F)$  drops out because of the “anti-Onsager” symmetry (46)<sub>2</sub>. This shows that a possible skewon piece is related to *dissipative* effects. For material media this is certainly a possibility. For empty spacetime it may be a legitimate hypothesis.

We will now turn to the propagation of the electromagnetic field (of “light”). In the geometrical optics limit, we find for the wave covector  $q$  the Fresnel equation [29, 5]

$$\mathcal{G}^{ijkl}(\chi) q_i q_j q_k q_l = 0, \quad (52)$$

with the (metric-free) Tamm-Rubilar tensor density

$$\mathcal{G}^{ijkl}(\chi) := \frac{1}{4!} \hat{\epsilon}_{mnpq} \hat{\epsilon}_{rstu} \chi^{mnr(i} \chi^{j|ps|k} \chi^{l)qtu}. \quad (53)$$

By simple algebra it is possible to prove that  $\chi^{[ijkl]}$  drops out from the Tamm-Rubilar tensor:  $\mathcal{G}^{ijkl}({}^{(1)}\chi + {}^{(2)}\chi + {}^{(3)}\chi) = \mathcal{G}^{ijkl}({}^{(1)}\chi + \chi^{(2)})$ . Furthermore,  $\mathcal{G}^{ijkl}({}^{(3)}\chi) = 0$ . In other words, the axion part  ${}^{(3)}\chi^{ijkl} = \epsilon^{ijkl}\alpha(x)$ , in the geometrical optics limit, does *not* influence the propagation of light. For the skewon piece we find  $\mathcal{G}^{ijkl}({}^{(2)}\chi) = 0$ . But even more far-reaching, by straightforward algebra, see [5], it can be shown that  $\mathcal{G}^{ijkl}({}^{(2)}\chi + {}^{(3)}\chi) = 0$ . Consequently, for  ${}^{(1)}\chi = 0$ , the Fresnel equation collapses and there is no orderly wave propagation in the geometrical optics limit. Therefore, the existence of conventional wave propagation requires  ${}^{(1)}\chi \neq 0$ , that is, the principal part of the constitutive tensor must be non-vanishing.

Although from Table 1 one may have an impression that the axion field is to a great extent irrelevant in experiments, it *does* affect physical observations. In particular, the important phenomenon of electromagnetic waves is certainly not exhausted by geometrical optics. Specifically, in the framework of axion electrodynamics (27) with a constant principal part  ${}^{(1)}\chi^{ijkl}$ , Itin [7] (see also Carroll et al. [2]) has shown that the axion field does influence electromagnetic waves if one looks into exact solutions. Assuming the usual plane wave ansatz for the electromagnetic field strength  $F = f e^{i\varphi}$ , with the constant amplitude 2-form  $f$  and the phase function  $\varphi(x^i)$ , we find a generalization of the Fresnel equation (52) in the form of [7]

$$\mathcal{G}^{ijkl}(\chi) q_i q_j q_k q_l - \chi^{ijkl} (\partial_i \alpha) (\partial_l \alpha) q_j q_k = 0. \quad (54)$$

Here  $q_i = \partial_i \varphi$  is the wave covector, as usual. This extended Fresnel equation is no longer an homogeneous algebraic equation, and the axion field contributes explicitly to the second term. This necessarily leads, in the presence of the axion field, to the birefringence of the waves.

As another manifestation of the axion field, let us consider the Lorentz force density  $f_i$  and the balance equation related to it (see [5], Sec. B5.1):

$$f_i = d\Sigma_i + X_i, \quad \text{with} \quad X_i := -\frac{1}{2} (F \wedge \mathcal{L}_i H - H \wedge \mathcal{L}_i F). \quad (55)$$

This is the force density that an electromagnetic field exerts on charges and currents. The axion field indeed drops out from the first term, which contains the electromagnetic energy-momentum

current (49). However, the second term is nontrivial. Using the elementary properties of the Lie derivative  $\mathcal{L}_i$ , we straightforwardly find for the constitutive law  $H = \lambda_0 \star F + \alpha F$ ,

$$X_i = -\frac{1}{2}(\partial_i \alpha) F \wedge F. \quad (56)$$

This additional force density arises from the axion field.<sup>3</sup>

## 4 Constitutive tensor split in space and time

We will now come back to the constitutive relations (5),(6). We will again work with a 3–vector formalism. Again, we formulate it in a *metric independent* way. We decompose the electromagnetic field with respect to the 3–dimensional coframe  $\vartheta^a$  and the 2–form basis  $\hat{e}_b = \hat{e}_{bcd} \vartheta^c \wedge \vartheta^d/2$ :

$$\mathcal{H} = \mathcal{H}_a \vartheta^a, \quad E = E_a \vartheta^a, \quad (57)$$

$$\mathcal{D} = \mathcal{D}^b \hat{e}_b, \quad B = B^b \hat{e}_b. \quad (58)$$

Then the constitutive relations (5),(6) can be put in the form of a matrix equation [5],

$$\begin{pmatrix} \mathcal{H}_a \\ \mathcal{D}^a \end{pmatrix} = \begin{pmatrix} \mathfrak{C}^b_a & \mathfrak{B}_{ba} \\ \mathfrak{A}^{ba} & \mathfrak{D}_b^a \end{pmatrix} \begin{pmatrix} -E_b \\ B^b \end{pmatrix} =: \begin{pmatrix} \kappa \end{pmatrix} \begin{pmatrix} -E_b \\ B^b \end{pmatrix}. \quad (59)$$

Moreover, by using the Levi-Civita symbol, see (40), we find for the  $\chi$  tensor density

$$\begin{pmatrix} \chi \end{pmatrix} = \begin{pmatrix} \mathcal{B}_{ab} & \mathcal{D}_a^b \\ \mathcal{C}^a_b & \mathcal{A}^{ab} \end{pmatrix}. \quad (60)$$

If we compare (60) with (17)<sub>1</sub>, then, by straightforward algebra, the constitutive  $3 \times 3$  matrices  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$  can be related to the 4-dimensional constitutive tensor density (40) by

$$\mathfrak{A}^{ba} = \chi^{0a0b}, \quad \mathfrak{B}_{ba} = \frac{1}{4} \hat{e}_{acd} \hat{e}_{bef} \chi^{cdef}, \quad (61)$$

$$\mathfrak{C}^a_b = \frac{1}{2} \hat{e}_{bcd} \chi^{cd0a}, \quad \mathfrak{D}_a^b = \frac{1}{2} \hat{e}_{acd} \chi^{0bcd}. \quad (62)$$

Let us now employ the irreducible decomposition. For the principal part of  $\kappa$  or  $\chi$ , respectively, we take the notation of Post [34], namely  $\varepsilon^{ab} = \varepsilon^{ba}$  (6 independent components),  $\mu_{ab}^{-1} = \mu_{ba}^{-1}$  (6 components), and  $\gamma^a_b$ , with  $\gamma^c_c = 0$  (8 components). The skewon part we parametrize according to

$$\mathcal{S}_i^j = \begin{pmatrix} -s_c^c & m^a \\ n_b & s_b^a \end{pmatrix}, \quad (63)$$

with, in 3 dimensions, the tensor skewon field  $s_a^b$  (9 independent components), the vector skewon  $m^a$  (3 components), and the covector skewon  $n_a$  (3 components). If we insert all of this in  $\kappa$ , we eventually find

$$\begin{pmatrix} \kappa \end{pmatrix} = \underbrace{\begin{pmatrix} \gamma^b_a & \mu_{ab}^{-1} \\ -\varepsilon^{ab} & \gamma^a_b \end{pmatrix}}_{\text{principal part 20 comp.}} + \underbrace{\begin{pmatrix} -s_a^b + \delta_a^b s_c^c & -\hat{e}_{abc} m^c \\ \varepsilon^{abc} n_c & s_b^a - \delta_b^a s_c^c \end{pmatrix}}_{\text{skewon part 15 comp.}} + \alpha \underbrace{\begin{pmatrix} \delta_a^b & 0 \\ 0 & \delta_b^a \end{pmatrix}}_{\text{axion part 1 comp.}}. \quad (64)$$

<sup>3</sup>There are proposals to combine the latter force term with the usual term  $d\Sigma_i$ , thus arriving at a modified energy-momentum current [2, 7]. However, this only works under the additional assumption of the constancy of the gradient of the axion field,  $d(\partial_i \alpha) = 0$ , and results in a non gauge invariant expression for the energy-momentum.

According to (59), we can evaluate the matrix elements:

$$\mathcal{H}_a = (\mu_{ab}^{-1} - \hat{\epsilon}_{abc} m^c) B^b + (-\gamma^b{}_a + s_a{}^b - \delta_a^b s_c{}^c) E_b - \alpha E_a, \quad (65)$$

$$\mathcal{D}^a = (\varepsilon^{ab} - \epsilon^{abc} n_c) E_b + (\gamma^a{}_b + s_b{}^a - \delta_b^a s_c{}^c) B^b + \alpha B^a. \quad (66)$$

Recall that  $\varepsilon^{ab} = \varepsilon^{ba}$ ,  $\mu_{ab}^{-1} = \mu_{ba}^{-1}$ , and  $\gamma^c{}_c = 0$ . Incidentally,  $\alpha$  is a 4-dimensional (axial) scalar, whereas  $s_c{}^c$  is only a 3-dimensional scalar. The cross-term  $\gamma^a{}_b$  is related to the Fresnel-Fizeau effects. The skewon contributions  $m^c, n_c$  are responsible for electric and magnetic Faraday effects, respectively, whereas the skewon terms  $s_a{}^b$  describe optical activity.

## 5 Constitutive tensor and physics

Up to here, we haven't said much about which piece of the constitutive tensor may be realized in nature. We just made a "kinematical" investigation of the most general local and linear constitutive tensor with its 36 independent components. From an analysis of the Tamm-Rubilar tensor we found that  ${}^{(1)}\chi^{ijkl} \neq 0$  is necessary for a non-degenerate propagation of electromagnetic waves in the geometrical optics limit. Only then the Tamm-Rubilar tensor doesn't vanish and the Fresnel equation exists. By the same method we find that, in the same limit,  ${}^{(3)}\chi^{ijkl} \neq 0$  doesn't disturb the wave propagation at all.

The *skewon* part  ${}^{(2)}\chi^{ijkl}$ , in its full generality, was discussed, amongst others, by Sihvola & Lindell [43, 19] and by us [5]. We studied already the possible effects of the skewon field on light propagation [30]. But we should mention Nieves and Pal [28] who introduced, besides the vacuum impedance  $1/\lambda_0$  and the velocity of light  $c$ , a third constant for the vacuum by means of a specially chosen skewon piece. Nieves and Pal assume (in our notation)

$$s_a{}^b = \frac{s}{2} \delta_a^b, \quad m^a = 0, \quad n_a = 0 \quad (\text{Nieves \& Pal}). \quad (67)$$

This corresponds to a spatially isotropic skewon field. By (67), a constraint is given that restricts the class of allowed reference frames. Lakhtakia<sup>4</sup> calls it "*isotropic chirality which is Lorentz reciprocal*". The Post constraint doesn't forbid it since the Nieves & Pal piece relates to the skewon part, whereas the violation of the Post constraint to the axion part. Lakhtakia points out that  $s \neq 0$  is valid for certain *materials*, but not for vacuum. Furthermore, others (see [5]) put forward the hypothesis that there can be a fourth electromagnetic constant that makes the medium "*nonreciprocal (but still isotropic)*". It is, of course, the axion field. With (67), the constitutive law (65),(66) reads

$$\mathcal{H}_a = -(s + \alpha) E_a + {}^{(1)}\mathcal{H}_a(E, B), \quad (68)$$

$$\mathcal{D}^a = {}^{(1)}\mathcal{D}^a(E, B) - (s - \alpha) B^a, \quad (69)$$

where we used the notation  ${}^{(1)}\mathcal{H} := {}^{(1)}\kappa(F)$  and  ${}^{(1)}\mathcal{D} := {}^{(1)}\kappa(F)$ . We rewrite (68),(69) for the case of a Maxwell-Lorentzian principal piece in exterior calculus as

$$\mathcal{H} = \mu^{-1} \lrcorner B - (s + \alpha) E, \quad (70)$$

$$\mathcal{D} = \varepsilon \lrcorner E - (s - \alpha) B. \quad (71)$$

---

<sup>4</sup>Private communication.

Note how the isotropic skewon field and the axion field act with different signs on  $E_a$  and  $B^a$ . Thereby they can be distinguished phenomenologically. We stress that the axion field  $\alpha$  is a premetric, 4-dimensional scalar with twist and as such of fundamental potential importance. In contrast, the skewon field  $s$  is only a 3-dimensional scalar valid in a constrained class of reference frames.—

What is then the *the state of the Post constraint*? Of course, the somewhat abstract concepts of a pure axion field, the Tellegen gyrator, and the PEMC of Lindell & Sihvola all violate the Post constraint, that is,  $\alpha \neq 0$ , whereas the other pieces of the constitutive tensor vanish at the same time.

One argument in favor of the Post constraint is the following (see [50]): With  ${}^{(1)}H := {}^{(1)}\kappa(F)$  and  ${}^{(2)}H := {}^{(2)}\kappa(F)$ , the inhomogeneous Maxwell equation for the linear constitutive law reads

$$dH = d({}^{(1)}H + {}^{(2)}H) + (d\alpha) \wedge F = J, \quad (72)$$

where we used already the homogeneous Maxwell equation  $dF = 0$ , see also (27). If the axion piece of the constitutive tensor is *constant*, i.e.,  $d\alpha = 0$ , then  $\alpha$  cannot couple to the Maxwell equations. However, this is only true for an infinitely extended material. As soon as the material has finite size, the axion piece  $\alpha$  jumps at the surfaces of the material — and this jump of  $\alpha$  couples to the Maxwell equations. In other words, the axion field can be seen along those jump surfaces by means of electromagnetic waves beyond the geometrical optics limit. In order to make this explicit, let us assume that a two-dimensional surface  $S$  divides the space into two parts. Let now both half-spaces be filled with homogeneous material media both of which are characterized by constant but unequal values of the axion field:  $\alpha_1$ , say, in the first material and  $\alpha_2$  in the second one. Then, if we take as an example the pure gyrator case with the constitutive law (30), we find from the jump conditions on the boundary (see [5], Sec. B.4.3),

$$[\alpha] n_a B^a \Big|_S = \tilde{\rho}, \quad [\alpha] \epsilon^{abc} n_b E_c \Big|_S = -\tilde{j}^a. \quad (73)$$

Here  $n_a$  are the components of the covector normal to the surface  $S$ , and  $[\alpha] := \alpha_2 - \alpha_1$  is the jump of the axion field on the boundary. Thus, even though the constant axion drops out from the Maxwell equations inside both homogeneous regions, it pops up in the form of the surface charge density  $\tilde{\rho}$  and the surface current density  $\tilde{j}^a$  induced on the boundary  $S$ . Both are proportional to the jump of the axion  $[\alpha]$ . In the more general case of axion electrodynamics, see (27), the left-hand sides in (73) will include the jump of the normal components of the electric excitation  $\mathcal{D}^a$  and the jump of the tangential components of the magnetic excitation  $\mathcal{H}_a$ , respectively. Hence our argument can be generalized straightforwardly.

This enables us to measure the value of the axion field of a homogeneous material medium. In this sense, the axion field of a material medium, our case (ii), always couples to the inhomogeneous Maxwell equation, and the argument mentioned above no longer applies. In the case of spacetime, also a curved one, the argument is correct if really everywhere  $d\alpha = 0$ . As soon as the axion field becomes space and/or time dependent, also in case (i) the quoted argument loses its meaning.

For empty spacetime, our case (i), the Post constraint, according to experimental evidence, is fulfilled. However, the relaxation of the Post constraint leads to the concept of the axion field. According to our Table 1, this field is very elusive since it neither shows up in the electromagnetic energy-momentum tensor nor in the Tamm-Rubilar-tensor, i.e., it doesn't influence light propagation in the limit of geometrical optics. Hence it is hard to detect it. A whole industry is at work to

find the axion field, see the new data on elementary particles [32], p. 389. So far without success. There is no theoretical reason known to us that would forbid the axion field. Quite the opposite. From a theoretical point of view, it appears as a quite natural extension of Maxwell-Lorentz electrodynamics. But, as mentioned, the axion hasn't been found so far.

For material media, our case (ii), the situation is different. Gytrators have been engineered, see [6]. Consequently, there is little doubt that materials can be constructed that violate the Post constraint. The Tellegen model with parallel or antiparallel electric and magnetic dipoles looks reasonable, see Tretyakov et al. [49]. In the meantime, Tretyakov et al. [48] built an artificial *Tellegen particle* and verified its nonreciprocal magnetoelectric behavior. "Natural" material media exist that violate the Post constraint, namely  $\text{Cr}_2\text{O}_3$  and in  $\text{Fe}_2\text{TeO}_6$  in static magnetic fields, see de Lange & Raab [18]. Accordingly, model calculations [33, 38] and experiments [18] (see also [37]) show that the Post constraint is valid for many materials; however, it is definitely *violated* in some rare cases. Thus, the Post constraint as a general dogma should be buried with all due honors.

## Acknowledgment

We would like to thank R.M. Kiehn, Ismo Lindell, and E. Jan Post for numerous fruitful discussions via email. We also appreciate having received from them pre- and reprints of their and others work. We are grateful to Frank Gronwald, Yakov Itin, and Roger Raab for helpful remarks. Financial support from the DFG (HE-528/20-1) is gratefully acknowledged.

## References

- [1] P. Bamberg and S. Sternberg, *A Course in Mathematics for Students of Physics*, Vol. 2 (Cambridge University Press: Cambridge, 1990).
- [2] S.M. Carroll, G.B. Field, and R. Jackiw, *Limits on a Lorentz- and parity-violating modification of electrodynamics*, *Phys. Rev.* **D41** (1990) 1231-1240.
- [3] R.H. Dicke, *The theoretical significance of experimental relativity*, (Gordon and Breach: New York, 1964).
- [4] T. Frankel, *The Geometry of Physics – An Introduction* (Cambridge University Press: Cambridge, 1997).
- [5] F.W. Hehl and Yu.N. Obukhov, *Foundations of Classical Electrodynamics — Charge, Flux, and Metric*. Birkhäuser, Boston (2003).
- [6] C.L. Hogan, *The ferromagnetic Faraday effect at microwave frequencies and its applications*, *Rev. Mod. Phys.* **25** (1953) 253–262.
- [7] Y. Itin, *Carroll-Field-Jackiw electrodynamics in the pre-metric framework*, *Phys. Rev.* **D70** (2004) 025012 (6 pages) [arXiv:hep-th/0403023].
- [8] Y. Itin and F.W. Hehl, *Is the Lorentz signature of the metric of spacetime electromagnetic in origin?* *Annals of Physics (NY)* **312** (2004) 60–83 [arXiv:gr-qc/0401016].
- [9] R.M. Kiehn, *The chiral vacuum*, preprint 1997, updated 2002 (11 pages), see *Cartan's Corner* <http://www22.pair.com/csdc/car/carhomep.htm>.
- [10] R.M. Kiehn, *Non-equilibrium and irreversible electrodynamics* November 2003 (76 pages), see <http://www.cartan.pair.com>.
- [11] R.M. Kiehn, G.P. Kiehn, and J.B. Roberds, *Parity and time-reversal symmetry breaking, singular solutions, and Fresnel surfaces*, *Phys. Rev.* **A43** (1991) 5665–5671.

- [12] A. Lakhtakia, *On the genesis of the Post constraint in modern electromagnetism* (15 pages) [arXiv:physics/0403042] (March 2004).
- [13] A. Lakhtakia, *Ten years past the Post*. In: Proc. SPIE (Int. Soc. for Optical Engineering), Vol. **5508** *Complex Mediums V: Light and Complexity*, M.W. McCall, G. Dewar (eds.). August 2004, pp. 85–94.
- [14] A. Lakhtakia and W.S. Weiglhofer, *Are linear, nonreciprocal, biisotropic media forbidden?* IEEE Trans. Microw. Theory Techn. **42** (1994) 1715–1716.
- [15] A. Lakhtakia and W.S. Weiglhofer, *On a constraint on the electromagnetic constitutive relations of nonhomogeneous linear media*, IMA J. of Appl. Math. **54** (1995) 301–306.
- [16] A. Lakhtakia and W.S. Weiglhofer, *Lorentz covariance, Occam’s razor, and a constraint on linear constitutive relations*, Phys. Lett. **A213** (1996) 107–111; erratum **A222** (1996) 459.
- [17] C. Lämmerzahl and F.W. Hehl, *Riemannian light cone from vanishing birefringence in premetric vacuum electrodynamics*, Phys. Rev. D, to be published 2004/05 [arXiv.org/gr-qc/0409072].
- [18] O.L. de Lange and R.E. Raab, *Post’s constraint for electromagnetic constitutive relations*, J. Optics **A3** (2001) L23–L26.
- [19] I.V. Lindell, *Differential Forms in Electromagnetics*. IEEE Press, Piscataway, NJ, and Wiley-Interscience (2004).
- [20] I.V. Lindell and A.H. Sihvola, *Perfect electromagnetic conductor* (6 pages) Preprint September 2004, submitted to JEMWA.
- [21] I.V. Lindell and A.H. Sihvola, *Transformation method for problems involving perfect electromagnetic conductor (PEMC) structures* (13 pages) Preprint October 2004, submitted to IEEE Trans. Antennas Propag.
- [22] I.V. Lindell, A.H. Sihvola, S.A. Tretyakov, A.J. Viitanen, *Electromagnetic Waves in Chiral and Bi-Isotropic Media*. Artech House, Boston (1994).
- [23] I.V. Lindell and K.H. Wallén, *Differential-form electromagnetics and bi-anisotropic Q-media*, J. Electromagn. Waves and Appl. **18** (2004) 957–968.
- [24] I.V. Lindell and K.H. Wallén, *Generalized Q-media and field decomposition in differential-form approach*, J. Electromagn. Waves and Appl. **18** (2004) 1045–1056.
- [25] W.-T. Ni, *A non-metric theory of gravity*. Dept. Physics, Montana State University, Bozeman. Preprint December 1973. The paper is available via <http://gravity5.phys.nthu.edu.tw/webpage/article4/index.html>.
- [26] W.-T. Ni, *Equivalence principles and electromagnetism*, Phys. Rev. Lett. **38** (1977) 301-304.
- [27] W.-T. Ni, *Equivalence principles and precision experiments*. In *Precision Measurement and Fundamental Constants II*, B.N. Taylor, Taylor B.N. W.D. Phillips, eds. Nat. Bur. Stand. (US) Spec. Publ. 617, US Government Printing Office, Washington, DC (1984).
- [28] J.F. Nieves and P.B. Pal, *The third electromagnetic constant of an isotropic medium*, Am. J. Phys. **62** (1994) 207-216.
- [29] Yu.N. Obukhov, T. Fukui, and G.F. Rubilar, *Wave propagation in linear electrodynamics*. Phys. Rev. D **62** (2000) 044050 [arXiv:gr-qc/0005018].
- [30] Yu.N. Obukhov and F.W. Hehl, *On possible skewon effects on light propagation* [arXiv:physics/0409155].
- [31] T.H. O’Dell, *The Electrodynamics of Magneto-Electric Media*, North-Holland, Amsterdam (1970).
- [32] Particle Data Group (S. Eidelman et al.), *Review of particle physics*, Phys. Lett. **592** (2004) 1–1109.
- [33] S. Ponti, J.A. Reyes and C. Oldano, *Homogeneous models for bianisotropic crystals*, J. Phys.: Condens. Matter **14** (2002) 10173–10182.
- [34] E.J. Post, *Formal Structure of Electromagnetics – General Covariance and Electromagnetics* (North Holland: Amsterdam, 1962, and Dover: Mineola, New York, 1997).
- [35] E.J. Post, *The constitutive map and some of its ramifications*, Ann. Phys. (NY) **71** (1972) 497-518.

- [36] E.J. Post, *Quantum Reprogramming — Ensembles and Single Systems: A Two-Tier Approach to Quantum Mechanics* (Kluwer: Dordrecht, 1995).
- [37] R.E. Raab, *Some unmeasured crystal properties*, Cryst. Res. Technol. **38** (2003) 202–214.
- [38] R.E. Raab and A.H. Sihvola, *On the existence of linear non-reciprocal bi-isotropic (NRBI) media*, J. Phys. **A30** (1997) 1335–1344.
- [39] P. Russer, *Electromagnetics, Microwave Circuit and Antenna Design for Communications Engineering* (Artech House: Boston, 2003).
- [40] J.A. Schouten, *Tensor Analysis for Physicists*, 2nd ed. reprinted (Dover: Mineola, New York 1989).
- [41] E. Schrödinger, *Space-Time Structure* (Cambridge University Press: Cambridge, 1954).
- [42] A.K. Schultz, R.M. Kiehn, E.J. Post, and J.B. Roberds, *Lifting of the four-fold EM degeneracy and PT asymmetry*, Phys. Lett. **A74** (1979) 384–386.
- [43] A.H. Sihvola and I.V. Lindell, *Material effects on bi-anisotropic electromagnetics*, IEICE Trans. Electron. (Tokyo) **E78-C** (1995) 1383–1390.
- [44] P. Sikivie, *Experimental tests of the “invisible” axion*, Phys. Rev. Lett. **51** (1983) 1415–1417.
- [45] P. Sikivie, ed., *Axions ’98*, in: *Proc. of the 5th IFT Workshop on Axions*, Gainesville, Florida, USA. Nucl. Phys. **B** (Proc. Suppl.) **72** (1999) 1–240.
- [46] B.D.H. Tellegen, *The gyrator, a new electric network element*, Philips Res. Rep. **3** (1948) 81–101.
- [47] B.D.H. Tellegen, *The gyrator, an electric network element*, Philips Technical Review **18** (1956/57) 120–124. Reprinted in H.B.G. Casimir and S. Gradstein (eds.) *An Anthology of Philips Research*. Philips’ Gloeilampenfabrieken, Eindhoven (1966) pp.186–190.
- [48] S.A. Tretyakov, S.I. Maslovski, I.S. Nefedov, A.J. Viitanen, P.A. Belov, A. Sanmatrin, *Artificial Tellegen particle*, Electromagnetics **23** (2003) 665–680.
- [49] S.A. Tretyakov, A.H. Sihvola, A.A. Sochava, C.R. Simovski, *Magnetolectric interactions in bi-anisotropic media*, J. Electromag. Waves Appl. **12** (1998) 481–497.
- [50] W.S. Weiglhofer, *On a medium constraint arising directly from Maxwell’s equations*, J. Phys. **A27** (1994) L871–L874.
- [51] W.S. Weiglhofer and A. Lakhtakia, *The Post constraint revisited*, Arch. Elektron. Übertrag. **52** (1998) 276–279.
- [52] F. Wilczek, *Two applications of axion electrodynamics*, Phys. Rev. Lett. **58** (1987) 1799–1802.

=====