

# **Fine structure of histograms of alpha-activity measurements depends on direction of alpha particles flow and the Earth rotation: experiments with collimators.**

**Simon E. Shnoll<sup>\*\*†</sup>, Konstantin I. Zenchenko<sup>†</sup>, Iosias I. Berulis<sup>‡</sup>, Natalia V. Udaltsova<sup>\*</sup> and Ilia A. Rubinstein<sup>§</sup>**

<sup>\*</sup> Institute of Theoretical and Experimental Biophysics of RAS, Pushchino, Moscow Region, 142290, Russian Federation

<sup>†</sup> Department of Physics, Moscow State University, Vorob'evy Gory, Moscow, 119992, Russian Federation

<sup>‡</sup> Radioastronomical Station, Pushchino, 142290, Russian Federation

<sup>§</sup> *Skobeltsyn Institute of Nuclear Physics, Moscow State University, Moscow, 119992, Russian Federation*

[Shnoll@iteb.ru](mailto:Shnoll@iteb.ru), [udanat@yahoo.com](mailto:udanat@yahoo.com)

## **Summary**

**The fine structure of histograms of measurements of  $^{239}\text{Pu}$  alpha-activity varies periodically, and the period of these variations is equal to sidereal day (1436 minutes). The periodicity is not observed in the experiments with collimator that restricts the alpha particles flow to be oriented to the Polar Star. Based on this study and other independent data, such as measurements conducted by the Arctic expedition, and similarity of the histograms in processes observed at different locations at the same local time, the conclusion was made, that the fine structure of statistical distributions of the observed processes depends on the celestial sphere.**

## **Introduction**

During many years, we explore changes in the fine structure of the histograms for consequent measurements of parameters of various physical processes. We call such changes as “macroscopic fluctuations” (**MF**). (Udaltsova, N.V. et al. 1987; Shnoll S.E. et al.1992, 1998, 2000; Fedorov M.V. et al.2003).

Comparing various physical processes (from measurements of noise in gravitation-gradient antenna and rates of biochemical reactions, to radioactive decay measurements) observed at different geographic locations at the same local (longitudinal) time, we have found the following:

1. Shapes of histograms in such different processes are similar at the same local time with high likelihood ( Shnoll S.E.2001).
2. The shape of histograms varies regularly in time. Here are some observations:
  - a. The histograms of the adjacent time intervals are similar with highest likelihood: this is the "neighbor zone effect".

b. For different locations similar histograms are observed most likely at the same local time: this fact indicates that histogram shape depends on the rotation of the Earth around its axis.

c. The recurrence of the similar shape of the histogram is observed with the period 1436 minutes, i.e. sidereal day.

These effects can be explained by the fact that, while the Earth is rotating around its axis and moves at the same time around the Sun, different sections of the Earth's surface are exposed to different sectors of the celestial sphere (Shnoll S.E., 2001; Fedorov et al. 2003).

The assumption that the shape of histograms depends on the celestial sphere "above the place of measurements", leads to the next ones: "the neighbour zone effect" and the about-daily variations of the histogram shape disappear at the geographical poles.

This hypothesis was confirmed in the study of  $^{239}\text{Pu}$  alpha-activity during the Arctic expedition at the Arctic Ocean. This expedition took place in August – October, 2000; the measurements were conducted at 82 degree of the North latitude, aboard the ship "The Academician Fedorov" ( Shnoll S.E.et al., 2003).

Additional confirmations of this assumption were obtained in the study of collimators restricting alpha particle flow in certain directions. The results of these experiments are presented in this article.

### **Objects and methods**

The most convenient object for the MF studies is the process of radioactive decay, especially alpha decay. Alpha decay is definitely immune to any trivial factors. It is possible to register separate acts of radioactive decay as a "binary" phenomenon (in logic of 0, 1).

This possibility was exercised in the specially constructed sensitive devices designed by one of the authors (I.A.R.). These devices provide transformation of the registered alpha activity data into time-series datasets with absent low-frequency trends. The results of the measurements according to all criteria correspond to "white noise" with Poisson distribution.


We used two identical devices for alpha-radioactivity measurements of  $^{239}\text{Pu}$  samples. Both devices provided rigid fixation of the source position in respect to the detector. The collimator (which "cuts out" a jet of alpha particles in a specified direction) may be placed between the source and the detector. One of the devices had the collimator oriented toward the Polar Star. The other device in one case had no collimator and the detector registering alpha particles emitting from the entire surface of the source which was West-oriented, in other case it had the collimator oriented to the West.

The distance between the source and the detector was 12 mm for both devices. While flying this distance, alpha particles lose about 10% of their energy. As the result, the energy of particles reaching the detector was  $\sim 4$  MeV. The registration threshold was  $\sim 1,6$  MeV; it completely excluded any possible influence of detector noise as well as humidity and air density variations.

A crystal generator (131 MHz frequency) was used for measuring time intervals. The power voltage of the converter was stabilized. The instability of the registration threshold was about  $\pm 6\%$  for the temperature range from minus  $30^{\circ}\text{C}$  to plus  $50^{\circ}\text{C}$ .


**Fig. 1** shows the histogram constructed for 60 of the results of consecutive measurements of alpha-activity of the  $^{239}\text{Pu}$  sample. The duration of 1 measurement is 1 second. X-axis represents

the measured parameter (in this case, the number of registered events of alpha decay). Y-axis shows the number of measurements in respect to the observed level of activity.




**Fig. 1**  
**A. Non-smoothed “inconsistent” histogram, 60 measurements.**  
**B. The same histogram, after single smoothing.**  
**C. The same histogram smoothed 15 times.**

As shown the actual value of the ordinate is quite small, not more than 4. The number of classes (in this case, it is 60) used for constructing these histograms is compatible to the number of measurements (in this case, it is also 60). As a result the original histogram is pockmarked. Such histograms are called "inconsistent". It is hard to find any regularity in change of shape in such histograms. However, things alter when we smooth these insufficient histograms using the method of moving averages. Look at **Fig. 1-B**; it is the same histogram after single smoothing. The result of 15 times smoothing is shown at **Fig. 1-C**. It is evident that, as a result of smoothing, the specific shape becomes visible. In the following discussion we consider just these histograms



**Fig. 2. Non-smoothed histograms constructed for non-overlapped sections of time-series ; every histogram represents 60 results of alpha-activity measurements in the  $^{239}\text{Pu}$  sample, the time interval for each histogram is 1 minute. Axes are the same as at Fig. 1.**

**Fig. 2** represents a number of non-smoothed inconsistent histograms constructed for the first 25 sections of time-series results of alpha-activity measurements in the  $^{239}\text{Pu}$  sample. (these sections are not overlapping). Each section covers 60 measurements within 1 minute in total. One couldn't find evident regularities for these distributions.



**Fig.3.** The same histograms as at **Fig. 2** smoothed 15 times. Axes are the same as at **Fig. 2**.

Fig. 3 represents the same histograms as at **Fig. 2**, after 15 times smoothing. The similar form of some histograms becomes evident. Is such a similarity accidental? To answer this question, we have to compare thousands of possible combinations of histogram pairs.

For this purpose, we used the computer program "Histogram Manager" (HM) designed by Edwin Pozharsky (Shnoll S.E. et al. 1998a). This program provides construction of histograms, their smoothing, imposition upon each other, some linear transformations, as well as the accumulation of the records of all selected similar pairs in a special file. The main function of this program is the calculation and construction of the frequency distribution of selected similar histograms against time-intervals between them. Nevertheless an expert performs the most important part of the analysis - the diagnosis "similar – not similar".

The comparison of many thousands of histogram pairs is a job extremely time and labor consuming. There have been many attempts to replace an expert with an adequate computer program, but all the attempts failed. The various methods were used - from regular calculations of correlation coefficients, statistical criteria estimation, to application of neural networks and Wavelet-analysis. The results of application of all these methods are still inferior to a visual estimation.

Some efforts were made to eliminate the bias in the diagnostic process. The program HM does this through the randomization of histogram numbers. In this case, the expert is blinded to the histogram numbers; so the distribution of the time-intervals between the similar histograms obtained by the expert may not be biased.

The main purpose of the study was investigation of the association between the realization of similar histograms and the value of time interval between them. Based on the results of histogram comparison, we construct the distribution of a number of similar pairs in respect to the length of the interval between them.

Many thousands of histogram pairs were compared to get the reliable distributions for similar pairs of histograms in respect to time intervals.

The estimation of statistical significance of maximums of these distributions was based on hypergeometric distribution (Pearson E.S. et al. 1958.) As shown in the work cited (N.V.Udaltsova see in: Shnoll S.E. et al. 2004), the Poisson distribution may be used as a general, broader estimation of the significance. We will utilize this method for conclusions that have a fundamental value.

## Results

From January 31 through July 11, 2002, and from February 19 through December 2003, ten sets of round-o-clock measurements of  $^{239}\text{Pu}$  alpha-activity were conducted at the Institute of Theoretical and Experimental Biophysics, RAN (RUSSIAN ACADEMY OF SCIENCE), Pushchino (latitude:  $54^{\circ} 50'$  North; longitude:  $37^{\circ} 38'$  East). Measurements were taken every second using two independent identical devices. One of the devices was equipped with the collimator oriented toward the Pole Star, the other one was either equipped with a collimator oriented to the West, or without collimator with detector and a radioactive sample plate oriented to the West. Each set of measurements continued about one month. The results of one-second measurements were summarized into one-minute rates. Every 60 one-minute numbers were combined into a histogram, so each histogram was built on one-hour interval of time-series. As a result we “replaced” a set of radioactive decay measurements by a set of consecutive one-hour histograms. After that we made a one-by-one comparison of smoothed histogram shapes and constructed a frequency chart of similar histograms against the time interval between them.

The results of experiments of 2002 – 2003 are presented in the table 1 and at the **fig 4**. In the **Table 1** counts of similar histograms are presented against the time interval between them. Measurements were conducted with no collimator (February 17, 2002 – July 11, 2002), and with collimator oriented to the West (February 19, 2003 – December 2, 2003).

**Table 1.**

**Counts of similar histograms of measurements of  $^{239}\text{Pu}$  alpha-activity by time interval between histograms (a flat detector without the collimator was used).**

Interval (hours)	17 Feb 2002	18 Mar 2002	15 Apr 2002	14 Jun 2002	11 Jul 2002	19 Feb 2003	19 Feb 2003 a	18 Mar 2003	15 Apr 2003	2 Dec 2003	Total
<b>1</b>	<b>118</b>	<b>79</b>	<b>86</b>	<b>66</b>	<b>76</b>	<b>79</b>	<b>71</b>	<b>89</b>	<b>93</b>	<b>133</b>	<b>890</b>
<b>2</b>	65	36	44	35	64	42	34	47	38	78	<b>483</b>
<b>3</b>	76	30	39	36	50	34	31	25	31	48	<b>400</b>
<b>4</b>	60	38	53	34	27	31	31	34	23	41	<b>372</b>
<b>5</b>	45	40	28	31	40	36	38	55	39	75	<b>427</b>
<b>6</b>	27	32	20	21	42	28	24	37	26	68	<b>325</b>
<b>7</b>	34	48	29	17	46	33	32	36	27	63	<b>365</b>
<b>8</b>	45	30	29	22	24	38	36	38	18	67	<b>347</b>
<b>9</b>	37	32	27	20	31	31	28	34	21	68	<b>329</b>
<b>10</b>	41	36	29	35	33	44	25	47	18	55	<b>363</b>
<b>11</b>	51	44	57	30	49	47	34	32	19	74	<b>437</b>
<b>12</b>	37	34	42	27	45	44	38	26	28	64	<b>385</b>
<b>13</b>	34	36	39	23	40	43	34	31	17	67	<b>364</b>


14	52	22	38	20	50	52	43	34	39	90	440
15	33	32	33	23	38	36	32	36	34	83	380
16	52	39	35	13	53	36	34	39	28	75	404
17	62	39	45	22	44	41	42	32	28	103	458
18	38	45	43	25	34	47	25	41	24	68	390
19	45	39	38	17	39	21	24	28	15	51	317
20	34	39	46	26	45	21	29	28	17	66	351
21	39	33	45	27	34	34	39	37	22	48	358
22	44	42	36	32	35	57	58	61	25	56	446
23	71	37	57	55	54	58	65	62	51	97	607
24	91	111	77	64	106	85	97	93	77	156	957
25	53	49	52	55	60	42	44	50	36	70	511
26	39	35	57	35	43	25	28	25	20	50	357
27	37	33	23	28	42	19	15	38	22	66	323
Total	1360	1110	1147	839	1244	1104	1031	1135	836	1980	

Let us look at the results of the first column of measurements (17 FEB 2002) in detail. There were selected 118 pairs of similar histograms related to the interval of 1 hour. For the interval of 2 hours, there were only 65 similar pairs. It is the "neighbour zone effect": the adjacent histograms have the highest chance to be similar. While the interval between histograms increases, the number of similar pairs decreases. However, at 24 hours, this number jumps high (to 91 pair of similar histograms).

Altogether, for the first cell, 697 histogram pairs with the interval of one hour were compared, that included experimental data obtained for almost a month (698 hours, ~ 29 days). The overall number of histogram pairs compared for the first column (as well as for each other column) was over 18000; a total count of selected similar histograms was 1360 (the lower line of the Table 1), i.e., about 7% of number of compared pairs.

Estimation of statistical significance based on hyper-geometric distribution gives the probability of random realization of the "neighbour zone effect" (interval 1 hour) for the first column as  $p < 10^{-9}$ . For the interval of 24 hours, it is  $p < 10^{-8}$ . The general estimation based on Poisson distribution shows that these two maximums far exceed 99% upper confidence limit.

The total number of pair comparisons of histograms in all ten sets of measurements in the Table 1 was approximately 180,000. The total counts of similar histograms for each time interval are shown in the last "Total" column and at the Fig.4. 890 similar pairs correspond to the first interval ( $p < 10^{-8}$ ). The count of similar histograms significantly decreases for larger intervals (to 390-430) and then it rises sharply for the interval of 24 hours (to 957,  $p < 10^{-8}$ ). The probability that these effects are random is extremely small.



**Fig. 4. Total counts of similar 60-minute histograms for measurements of alpha-activity  $^{239}\text{Pu}$  without collimator. X-axis represents time interval between similar histograms. Y-axis shows a number of similar pairs.**


Table 2 and Fig. 5 represent counts of similar histograms for experiments with the collimator oriented toward the Polar Star. The difference with Table 1 and Fig.4 is obvious: there is neither "neighbour zone effect", nor daily periodicity.

**Table 2.**

**Counts of similar histogram pairs of measurements of  $^{239}\text{Pu}$  alpha-activity with the collimator oriented toward the Polar Star regarding the length of time interval between histograms.**

Interval (hours)	17 Feb 2002	18 Mar 2002	15 Apr 2002	14 Jun 2002	11 Jul 2002	19 Feb 2003	19 Feb 2003 a	18 Mar 2003	15 Apr 2003	2 Dec 2003	Total
1	51	46	73	37	55	47	53	56	25	63	506
2	66	46	55	39	54	24	30	42	20	47	423
3	74	56	45	39	64	33	37	40	14	35	437
4	71	44	62	29	47	27	33	26	13	28	380
5	64	53	64	45	78	41	44	41	16	57	503
6	55	46	55	35	56	40	41	34	21	45	428
7	57	47	57	32	68	24	41	31	25	55	437
8	62	50	66	23	71	25	42	40	15	58	452
9	67	49	58	25	71	22	34	44	16	47	433
10	60	50	81	25	54	34	23	46	19	57	449
11	87	46	72	35	52	29	19	31	16	52	439
12	76	34	72	35	55	40	31	41	16	39	439
13	52	50	62	27	48	48	32	41	19	33	412

14	82	33	53	25	57	46	44	37	32	57	466
15	63	40	73	18	51	34	25	36	24	46	410
16	74	37	75	25	56	37	26	49	18	47	444
17	95	47	79	34	41	27	22	38	16	43	442
18	71	51	67	24	61	33	21	42	26	44	440
19	74	53	66	29	50	21	36	70	17	42	458
20	67	47	63	34	48	34	38	56	17	53	457
21	92	44	76	30	53	55	36	41	24	52	503
22	83	52	53	33	49	45	46	44	24	46	475
23	75	42	67	32	59	58	49	42	30	62	516
24	76	47	67	35	62	33	36	56	37	74	523
25	72	51	72	36	59	37	34	44	40	66	511
26	86	40	57	28	71	46	36	39	22	56	481
Total	1852	1201	1690	809	1490	940	909	1107	562	1304	



**Fig.5. Counts of similar 60-minute histograms regarding the time interval between them for measurements of alpha-activity  $^{239}\text{Pu}$  with collimator oriented to the Polar Star. Axes as at the Fig. 4.**


As we see at given figures, a chance of repeated appearance of the same shape of histogram is essentially different in measurements with the collimator oriented to the Polar Star and measurements without collimator or when collimator oriented to the West. In the measurements without collimator, or with the collimator oriented to the west, the obvious effect of “neighbor zone” is observed, as well as about-24-hour period. In the measurements with the collimator oriented toward the Polar Star a likelihood of repeated appearance of the same shape of histogram is about the same along time – there is neither “neighbor zone”, nor about-daily periodicity. Estimation of probability to obtain such differences in total distributions gives very small values.



About-daily periodicity in appearance of the same shape of histogram in the measurements without collimator, or when collimator oriented to the West, is equal exactly to 1436 minutes, i.e. sidereal day. This period was observed in one-minute histograms constructed for 60 one-second measurements. In these experiments we compared several sets of one-minute histograms. We considered pairs with intervals of 1434 to 1442 minutes between histograms. Each set had 698 histograms. In each experiment, for each column of Table 3, 6282 histogram pairs have been compared, and a count of similar histograms for each time interval has been written into corresponding cell. The average proportion of similar histograms was about 5%. The frequency of appearance of similar histograms at the interval of 1436 minutes is more than 2 times higher than average frequency for other intervals. Estimated probability of random appearance of such effect is less than  $10^{-6}$ .

**Table 3. Comparison of one-minute histograms of  $^{239}\text{Pu}$  alpha-activity: 24-hour period in histogram shape resemblance is equal 1436 minutes, i.e. sidereal day.**

Number of similar histogram pairs						
Interval (minutes)	3-4 Jul 2003	5-6 Jul 2003	12-13 Jul 2003	22-23 Jun 2003	24-25 Jun 2003	Total
1434	19	27	31	25	23	125
1435	31	31	36	42	35	175
<b>1436</b>	<b>97</b>	<b>92</b>	<b>68</b>	<b>95</b>	<b>82</b>	<b>434</b>
1437	45	43	35	51	46	220
1438	45	32	28	33	33	171
1439	29	27	25	26	32	139
1440	35	38	21	38	23	155
1441	19	15	25	16	20	95
1442	9	13	11	19	18	70
Total	<b>329</b>	<b>318</b>	<b>280</b>	<b>345</b>	<b>312</b>	<b>1584</b>



**Fig. 6. Count of similar histogram pairs for measurements of  $^{239}\text{Pu}$  alpha-activity with collimator oriented to the West. About-daily period of appearance of the same histogram shape is equal to 1436 minutes. X-axis is intervals between histograms in minutes. Y-axis is total number of similar histogram pairs.**

## Conclusions

The results exclude any trivial explanation. False effects are excluded not only by strict following to the principle of “equality of other conditions”( *ceteris paribus* principle) – the only difference between “case” and “control” is direction of alpha particles registered in experiments. It is absolutely impossible to affect the radioactive decay itself. It is paradoxical that in experiments conducted on  $54^\circ$  North latitude with collimator oriented toward the Polar Star the shapes of histograms vary in time by the same way as histograms received for measurements near the North Pole (Shnoll S.E., 2003).

It may indicate that probability of flow of alpha particles is not the same for different directions, and variations of this probability in time depend on changes in the surrounding time-space continuum. The presence of clear periodicity in these changes with the sidereal period of 1436 minutes (Shnoll S.E., 2001), the higher frequency of the appearance of similar histograms at the same local (longitudinal) time, the disappearance of daily periodicity near the North Polar, and, finally, presented above results of the experiments with collimators, - all these facts suggest the dependence of the observed effects on the state of the celestial sphere. The registration of such dependence is not sufficient for declaring a hypothesis, which might explain its "mechanism". However, in this article our task is only to register the significance of these astonishing phenomena.

## Literature

1. Udaltsova, N.V., Kolombet, V.A., and Shnoll, S.E., A Possible Cosmophysical Origin of Macroscopic Fluctuations in Various Processes, Puschino: ONTI NtsBI, 1987 (in Russian)

2. Pearson E.S., Hartley H.O. Biometrika Tables for statisticians, Vol 1. Cambridge Univ. Press, 1958.
3. Shnoll S.E., Udaltsova N.V., Kolombet V.A., Namiot V.A., and Bodrova N.B.:1992, Patterns in the discrete distributions of the results of measurements (cosmophysical aspects) *Biophysics* **37**(3),378-398.
4. Shnoll S.E.: 1995, The form of the spectra of states realized in the course of macroscopic fluctuations depends on the rotation of the Earth about its axis, *Biophysics* **40**(4),857-866
5. Shnoll S.E. , Kolombet V.A., Zenchenko T.A. , Shnoll S.E., Kolombet V.A., Pozharski E.V., Zvereva I.M. and Konradov A.A.: 1998a , Cosmophysical origin of “macroscopic fluctuations”, *Biophysics* **43**(5).864-870
6. Shnoll S.E. , Kolombet V.A. , Pozharski E.V. , Zenchenko T.A. , Zvereva I.M. and Konradov A.A. :1998b, Realization of discrete states during fluctuations in macroscopic processes, *Physics-Uspehi* **162**(10), 1129-1140.
7. Shnoll S.E. , Pozharski E.V. , Zenchenko T.A. , Kolombet V.A. , Zvereva I.M. and Konradov A.A.: 1999, Fine structure of distributions in measurements of different processes as affected by geophysical and cosmophysical factors, *Phys.& Chem. Earth A: Solid Earth & Geod.***24**(8), 711–714.
8. Shnoll S.E. , Zenchenko T.A. , Zenchenko K.I., Pozharski E.V., Kolombet V.A., and Konradov A.A.: 2000, Regular variation of the fine structure of statistical distributions as a consequence of cosmophysical agents, *Physics-Uspehi* **43**(2), 205-209,
9. Shnoll,S.E.: 2001, Discrete distribution patterns: arithmetic and cosmophysical origins of their macroscopic fluctuations, *Biophysics* **46**(5),733-741.
10. Fedorov M.V., Belousov L.V., Voeikov V.L., Zenchenko T.A., Zenchenko K.I., Pozharskii E.V., Konradov A.A. and Shnoll S.E.: 2003, Synchronous changes in dark current fluctuations in two separate photomultipliers in relation to Earth rotation, *Astrophysics and Space Science*. **283**:3-10.
11. Shnoll S.E., Rubinstein I.A., Zenchenko K.I. , Zenchenko T.A., Konradov A.A , Shapovalov S.N., Makarevich A.V., Gorshkov E.S., and Troshichev O.A.: 2003, Dependence of “Macroscopic Fluctuations” on Geographic Coordinates (by Materials of Arctic and Antarctic Expeditions), *Biophysics* **48**(6),1123-1131
12. Shnoll S.E.,Zenchenko K.I., Berulis I.I., Udaltsova N.V., Zhirkov S.S., Rubinstein I.A.: 2004, The Dependence of “Macroscopic Fluctuations” on Cosmophysical Factors. Spatial Anizotropy, *Biophysics* **49**(1) ,132-139

## ACKNOWLEDGEMENTS

We are indebted to M.N. Kondrashova and L.A. Blumenfeld who inspired, encouraged, and discussed this work. Special thanks are to Y.M. Popov who built the mechanical part of detectors used in this study. We are grateful to our colleagues, T.A. Zenchenko, A.A. Konradov, and S.S. Zhirkov for collaboration and assistance; and to V. K. Lyapidevskii, B.M. Vladimirsky, V.A.Tverdislov, A.P.Levich, B.V. Komberg, A.S. Kondrashov, F.A. Kondrashov, D.P. Kharakoz, F.I. Ataulakhanov, V.N. Morozov, I.M. Dmitrievsky, Yu.S.Vladimirov, D.S.Chernavsky and B.U. Rodionov for valuable discussions. Financial support obtained from V.P. Tikhonov and T. Peterson and their interest to this work are gratefully acknowledged.

Correspondence and requests for materials should be addressed to S.E.S. (e-mail: [shnoll@iteb.ru](mailto:shnoll@iteb.ru)).