The spectra of mixed ³He-⁴He droplets

S. Fantoni

SISSA and INFM DEMOCRITOS National Simulation Center, Via Beirut 2-4, I-34014 Trieste, Italy

R. Guardiola

Departamento de Física Atómica y Nuclear, Facultad de Física, E-46100-Burjassot, Spain

J. Navarro

IFIC (CSIC-Universidad de Valencia), Apartado Postal 22085, E-46071-Valencia, Spain

A. Zuker

Institute de Recherches Subatomiques, IN2P3-CNRS/Université Louis Pasteur, F-67037 Strasbourg, France

(Dated: November 15, 2018)

The diffusion Monte Carlo technique is used to calculate and analyze the excitation spectrum of 3 He atoms bound to a cluster of 4 He atoms, by using a previously determined optimum filling of single-fermion orbits with well defined orbital angular momentum L, spin S and parity quantum numbers. The study concentrates on the energies and shapes of the three kinds of states for which the fermionic part of the wave function is a single Slater determinant: maximum L or maximum S states within a given orbit, and fully polarized clusters. The picture that emerges is that of systems with strong shell effects whose binding and excitation energies are essentially determined by averages over configuration at fixed number of particles and spin, *i.e.*, by the monopole properties of an effective Hamiltonian.

PACS numbers: 36.40.-c 61.46.+w

I. INTRODUCTION

The study of liquid helium in confined geometries is currently an active area of experimental and theoretical research.^{1,2} Helium droplets are weakly bound quantum systems, as a consequence of their small atomic mass and the particular form of the associated van der Waals interaction. For ⁴He, clusters are bound for any number of atoms, while for ³He, it takes about 30 - 32 atoms^{3,4} to form a bound system, due to the larger zero-point motion and the Pauli Principle. The case of mixed ³He-⁴He clusters is very interesting, since they are made of particles with different statistics and masses interacting through the same potential. The theoretical calculations predict the existence of instability islands for a sufficiently small number of ⁴He atoms.^{5,6,7,8}

Experimentally, small helium clusters are produced by free jet expansion of the gas. Their mass is then measured by diffraction through a transmission grating followed by a mass spectrometer detector.⁹ Pure ⁴He clusters, containing up to eight atoms, and mixed clusters, containing one ³He and up to six ⁴He atoms, have been detected using a grating with 100 nm period,¹⁰ and even the very weakly bound dimer ⁴He₂ has been unambiguously detected.¹¹ The experimental setup of Ref. 9 has been improved to detect droplet sizes up to 25-30 amu, and very small mixed systems have been definitely identified.¹²

There have been several theoretical studies of a single ³He atom in a medium size ⁴He cluster, either through a density functional approach^{13,14} or microscopic methods.^{15,16,17} All ⁴He_{N_B}³He clusters form bound states for $N_B \geq 2$. The excess in kinetic energy pushes the

³He atom to the surface, resulting in a a quasi twodimensional wave function similar to the Andreev state describing one ³He impurity in ⁴He bulk. Recently, the ordering of the single-particle orbital states has been established.¹⁸ Larger combinations of ³He and ⁴He atoms have been studied employing a non-local finite range density functional.^{14,19,20} Previous variational microscopic studies of these mixed systems have been carried out for the ${}^{4}\text{He}_{2}{}^{3}\text{He}_{2}$ cluster, 21 and for clusters with $N_{B} \leq 8$, $N_F < 20.5$ More recently, the diffusion Monte Carlo (DMC) method has been applied to droplets with $N_F \leq$ 3, $N_B \leq 17$ in Refs. 6,7, and with $N_F \leq 20$, $N_B \leq 8$ in Ref. 8. The most important result from microscopic calculations^{5,8} is the prediction of instability regions, specially when the number of ⁴He atoms is small; nevertheless, a core with five or more ⁴He atoms is able to bind any number N_F of ³He atoms.

Our purpose is to analyze the ground state and the low-lying excited states of the ³He atoms in a mixed helium droplet, by using the DMC method, and relying on the single-particle orbital orderings obtained previously¹⁸ from the study of a single fermion. The resulting spacings resemble those of the rotational spectrum of a diatomic molecule, where the ⁴He core plays the role of one atom and the ³He is the other atom. At low energies each level is uniquely classified by its angular momentum, but vibrational-like excitations appear at higher energies for heavy enough clusters. The adopted filling order for large number of ³He atoms is therefore 1s 1p 1d ..., with some attention being paid to a possible 2s intruder. We shall concentrate on clusters with eight and twenty bosons, varying the number of fermions. This moderately large number of bosons ensures the existence of several bound excited levels. All calculations have been made using the HFD-(B) potential of Aziz and co-workers²² for the He-He interaction.

The plan of the paper is as follows. In Section II we give some general details concerning the DMC calculation. Section III collects single-fermion results needed later. Section IV deals with the case of two fermions: this system is simple enough to be analyzed in depth for a large set of quantum numbers, shedding light in particular on the single-particle ordering. Section V is devoted to stable mixed clusters, for the specific cases of maximum spin and maximum orbital angular momentum. The case of fully polarized clusters is also considered. In Section VI we exhibit the calculated one- and two-body distribution functions, both for normal and fully polarized mixed clusters. Finally, in Section VII we summarize our findings and draw some general conclusions.

II. THE DIFFUSION MONTE CARLO METHOD

The DMC^{23,24} method is based on an importance sampling wave function, with the double role of controlling the variance of the ground state energy and to incorporate both the required statistics and the desired spin and angular momentum quantum numbers. In this work we use the same form employed previously.⁸ The trial wave function is written as the product of four factors,

$$\Psi_T = \Phi_{BB} \Phi_{BF} \Phi_{FF} D_F \tag{1}$$

corresponding to a Jastrow form for the boson-boson (BB), boson-fermion (BF), and fermion-fermion (FF) parts, and the Slater determinantal part D_F for the fermions. Each Jastrow term is a translationally invariant and symmetric wave function, with the structure

$$\Phi_{MN} = \prod_{i,j} e^{f_{MN}(r_{ij})} \tag{2}$$

where indices M, N represent bosons (B) or fermions (F), and indices i, j run over the corresponding atoms. The product runs over all different pairs i < j if M = N, and no restrictions apply when $M \neq N$. We have used a simple but nevertheless physically complete representation

$$f_{MN}(r) = -\frac{1}{2} \left(\frac{b_{MN}}{r}\right)^{\nu_{MN}} - r \, p_{MN}, \qquad (3)$$

depending on three parameters, b, p and ν , in general different for each of the three BB, BF and FF pairs. The short range coefficients b and ν have been fixed, independently of the number of bosons or fermions, to the values $\nu = 5.2$ for any subset MN, $b_{BB} = 2.95$ Å, $b_{BF} = 2.90$ Å, and $b_{FF} = 2.85$ Å. The long range part $r p_{MN}$ entering Eq. (3), has the role of confining the system and fixing roughly its size. The three parameters p_{MN} are adjusted for each droplet by minimizing its ground state energy. The last factor D_F in the trial wave function is the determinantal part, which will be discussed later on for each specific case. The description of the trial wave function is completed by the inclusion of the Feynman-Cohen backflow²⁵ in the fermionic exponential tail as well as in the radial dependence of the Slater factor. Following Ref. 26, we have replaced each fermion coordinate \mathbf{r}_i by a transformed coordinate

$$\hat{\mathbf{r}}_{i} = \mathbf{r}_{i} + \sum_{i \neq j} \eta(r_{ij}) \left(\mathbf{r}_{i} - \mathbf{r}_{j}\right)$$
(4)

For the backflow function $\eta(r)$ we choose the mediumrange form used in Ref. 27:

$$\eta(r) = \frac{\lambda}{r^3} \tag{5}$$

keping the same value for the parameter $\lambda = 5 \text{ Å}^3$.

Using the model described above for the importance sampling guiding function, a real-time DMC evolution has been carried out employing a $\mathcal{O}(\tau^3)$ approximation to the Green function, with time slice $\tau = 0.00025 \text{ K}^{-1}$, for a total of 200 blocks of 200 steps each. A block average was used in order to diminish the unavoidable correlations of the DMC method, with the aim of obtaining a reasonable estimate of the variance. An initial population of 1000 walkers lead, on the average, to a total number of forty millions samples.

In spite of this block averaging, the resulting estimate of the variance turned out to be quite optimistic, *i.e.*, much smaller than reasonably expected. As a consequence, we opted for a very costly but safe procedure, by carrying out ten independent calculations with the same time slice and the same number of samples, but with randomly selected initial set of walkers. The resulting ten results lead to the *true* variance (or so we hope). The numbers quoted in what follows correspond to this prescription.

Due to the presence of the Slater determinants, the importance sampling function is not definite positive. In our calculations, the random process has been constrained using the so-called fixed node approximation: any walker attempting to cross a nodal surface is neglected. As it has been shown,^{29,30} the use of this approximation leads to an upper bound to the lowest energies.

III. THE ONE FERMION SYSTEM

The system made of a single ³He atom plus a drop of ⁴He atoms has been recently investigated.¹⁸ For the sake of completeness we summarize in this section the most relevant results for the analysis of systems with more fermions.

As there is only one fermion, the Slater determinant in Eq. (1) becomes a single-particle wave function $\phi_{n\ell m}(\mathbf{r})$ of radial quantum number n and orbital angular momentum ℓ with projection m. Translational invariance is en-

sured by referring the fermion coordinate to the bosonic center of mass.

Different values of ℓ select a specific angular momentum subspace in which the DMC procedure will drive the wave function so as to minimize the energy. Thus, the radial part of the fermionic wave function can be taken to have the form

$$\phi_{1\ell m}(r) = r^{\ell} Y_{\ell m}(\hat{r}), \tag{6}$$

which ensures that there are no radial nodes (n = 1). These single-particle wave functions do not contain a radial confining term, because it is already included in the Jastrow part of the importance sampling wave function. Notice that within the DMC procedure it is not possible to obtain excited levels with n > 1, because this will require to impose a strict orthogonality on the physical n = 0 ground state. Nevertheless, one may use an indirect procedure based in the moment method which will provide an upper bound to the energy of the radial excited levels.

Table I displays the energy of these one-fermion states, as a function of the single-fermion quantum numbers. The quoted energies are slightly different from those of Ref. 18 because of the improved statistics of the present calculation. The ground state energies of the pure bosonic droplets are also given, so as to define the *separation energies*

$$\epsilon_{n\ell} = E_{n\ell}(N_B, N_F = 1) - E(N_B, N_F = 0), \quad (7)$$

also quoted in the Table. Positive values refer to unbound levels. The energy obtained for the $n = 2, \ell = 0$ excited level is above the dissociation limit for $N_B = 8$ or very close to it for $N_B = 20$.

TABLE I: Energies of droplets with eight and twenty bosons and one fermion, for several values of the angular momentum. The row labelled $N_F = 0$ is the system without fermions. The columns labelled ϵ are the separation energies. Values in italics correspond to unbound levels.

Config	$N_B =$	= 8	$N_B = 20$		
	E(K)	$\epsilon(K)$	E(K)	ϵ (K)	
$N_F = 0$	-5.14(1)		-33.76(2)		
1s	-6.08(1)	-0.94	-35.55(1)	-1.79	
1p	-5.60(1)	-0.46	-35.15(2)	-1.39	
1d	-4.98(1)	0.16	-34.55(2)	-0.89	
2s	-5.10	0.06	-33.80	-0.04	

The single-particle wave functions of Eq. (6) will be used later to construct model wave functions for systems with two or more fermions. Unfortunately the moment method does not provide a wave function for the radial excitation, and for the calculations of the following sections we will use the simple form

$$\phi_{n=2,\ell=0}(r) = r^2, \tag{8}$$

without any radial node. The lack of nodes may be a deficiency, but the truly important point is that the simple form chosen is linearly independent of the 1s state.

IV. THE TWO FERMION SYSTEM

Adding two ³He atoms to a core of ⁴He atoms results in a system deeply resembling the helium atom. To a large extent, the bosonic sub-cluster plays the role of the atomic nucleus, with the two ³He atoms corresponding to the electrons. There are two families of levels, singlet (S = 0) and triplet (S = 1). Each of the states is characterized by the configuration, the orbital angular momentum L, the spin S and the parity P.

Specific two-fermion states are constructed from the single-particle wave functions (6) by coupling the angular momentum part to the required quantum numbers and by symmetrizing the singlet or antisymmetrizing the triplet radial wave functions. In general one does not obtain real wave functions, but the remedy is simple. For M = 0 the result is real. Otherwise construct two cases for M and -M and either add or subtract them. The wave function thus constructed has no good L_z , but still good total orbital angular momentum, and should have no effect on the computed energy.

For the sake of completeness we list in Table II the

TABLE II: Space part of importance sampling singlet(upper part) and triplet (lower part) wave functions for the two-fermion drop.

L^P	Config	Ψ (SINGLET)
0^{+}	$1s^2$	1
0^{+}	$1p^2$	$\mathbf{r}_1 \cdot \mathbf{r}_2$
0^{+}	$1d^2$	$3(\mathbf{r}_1 \cdot \mathbf{r}_2)^2 - r_1^2 r_2^2$
1^{-}	1s1p	$z_1 + z_2$
		$z_1r_2^2 - 3z_2(\mathbf{r}_1\cdot\mathbf{r}_2)z_2r_1^2 - 3z_1(\mathbf{r}_1\cdot\mathbf{r}_2)$
2^{+}	1s1d	$x_1^2 + x_2^2 - y_1^2 - y_2^2$
2^{+}	$1p^2$	$x_1x_2 - y_1y_2$
2^{+}	$1d^2$	$(x_1^2 - y_1^2)r_2^2 + (x_2^2 - y_2^2)r_1^2 -$
		$3(x_1x_2-y_1y_2)(\mathbf{r}_1\cdot\mathbf{r}_2)$
2^{-}	1p1d	$x_1x_2z_2 - x_2^2z_1 - y_2(y_1z_2 - y_2z_1) +$
		$x_2x_1z_1 - x_1^2z_2 - y_1(y_2z_1 - y_1z_2)$
3^{-}	1p1d	$x_1(y_2^2 - x_2^2) + 2x_2y_1y_2 +$
		$x_2(y_1^2 - x_1^2) + 2x_1y_1y_2$
4^{+}	$1d^2$	$(x_1^2 - y_1^2)(x_2^2 - y_2^2) - 4x_1x_2y_1y_2$
		$r_1^2 + r_2^2$
		$z_1 r_2^2 + z_2 r_1^2$
L^P	Config	Ψ (TRIPLET)

$1^{-} 1s1p$	$z_1 - z_2$
$1^{-} 1p1d$	$z_1r_2^2 - 3z_2(\mathbf{r}_1\cdot\mathbf{r}_2) - z_2r_1^2 + 3z_1(\mathbf{r}_1\cdot\mathbf{r}_2)$
1^{+} $1p^{2}$	$x_1 z_2 - x_2 z_1$
$1^{+} 1d^{2}$	$(x_1z_2-x_2z_1)(\mathbf{r}_1\cdot\mathbf{r}_2)$
$2^+ 1s1d$	$x_1^2 - x_2^2 - y_1^2 + y_2^2$
2^{-} 1p1d	$x_1x_2z_2 - x_2^2z_1 - y_2(y_1z_2 - y_2z_1) -$
	$x_2x_1z_1 + x_1^2z_2 + y_1(y_2z_1 - y_1z_2)$
$3^+ 1d^2$	$x_1^2(y_2^2 - z_2^2) - x_2^2(y_1^2 - z_1^2) + y_1^2z_2^2 - y_2^2z_1^2$
$3^{-} 1p1d$	$x_1(y_2^2 - x_2^2) + 2x_2y_1y_2 -$
	$x_2(y_1^2 - x_1^2) - 2x_1y_1y_2$
$0^+ \ 1s2s$	$r_1^2 - r_2^2$
$1^ 1p2s$	$z_1 r_2^2 - z_2 r_1^2$


specific forms thus obtained, and used as importancesampling wave functions to drive the DMC stochastic procedure.

TABLE III: Binding energies, in K, for several states with two fermions and $N_B = 8$ (upper table) and $N_B = 20$ (lower table), classified by the configuration and angular momentum quantum numbers. Values in bold face correspond to the physically interesting states. The dissociation limit is 6.08 K ($N_B = 8$) and 35.55 K ($N_B = 20$). The statistical errors of the energies are between 0.01 and 0.02 K. The row labelled *Eff.* is the prediction of the non-interacting fermion model.

				$N_B = 8$				
$L^P S$	$1s^2$	1s1p	$1p^2$	1s1d	1p1d	$1d^2$	1s2s	1p2s
$0^{+}0$	7.09		6.12			4.90	7.07	
$1^{-}0$		6.61			5.49			6.49
$2^{+}0$			6.15	5.99		4.88		
$2^{-}0$					5.50			
$3^{-}0$					5.48			
$4^{+}0$						4.85		
$0^{+}1$							5.89	
$1^{+}1$			6.19			5.06		
$1^{-}1$		6.65			5.65			6.46
$2^{+}1$				6.01				
2^{-1}					5.52			
$3^{+}1$						4.89		
$3^{-}1$					5.45			
Eff.	7.02	6.54	6.06	5.92	5.44	4.82	6.02	
			1	$V_B = 20$)			
$L^P S$	$1s^2$	1s1p	$1p^2$	1s1d	1p1d	$1d^2$	1s2s	1p2s
$0^{+}0$	37.33		36.70			35.45	37.33	
$1^{-}0$		37.06			36.05			36.98
$2^{+}0$			36.74	36.47		35.54		
$2^{-}0$					36.19			
$3^{-}0$					36.17			
$4^{+}0$						35.55		
							35.68	
$0^{+}1$								
$1^{+}1$			36.74			35.79		
$1^{+}1$ $1^{-}1$		37.05	36.74		36.27	35.79		36.99
$ 1^{+1} \\ 1^{-1} \\ 2^{+1} $		37.05	36.74	36.49		35.79		36.99
1^{+1} 1^{-1} 2^{+1} 2^{-1}		37.05	36.74	36.49	36.27 36.20			36.99
$ \begin{array}{c} 1^{+1} \\ 1^{-1} \\ 2^{+1} \\ 2^{-1} \\ 3^{+1} \end{array} $		37.05	36.74	36.49	36.20	35.79 35.66		36.99
1^{+1} 1^{-1} 2^{+1} 2^{-1}	37.34	37.05 36.94	36.74	36.49			35.59	36.99

The energies obtained from these configurations are displayed in Table III, for $N_B = 8$ and $N_B = 20$. Some among these results lie above the dissociation limit (*i.e.*, the energy of the lowest bound state with the same number of bosons but with a single fermion) and do not correspond strictly to truly bound systems.

It has to be understood that in the results quoted in these two tables, only a single number per row does have a physical sense: the DMC algorithm improves systematically the importance sampling wave function, but because of the use of the fixed-node approximation, that improvement only provides a variational upper bound to the energy for each of the subspaces with well defined L, S and P quantum numbers. To give an example: the state ${}^{1}S$ may have projections on the $1s^{2}$, $1p^{2}$, $1d^{2}$ and FIG. 1: The two-fermion spectrum for $N_B = 8$ and $N_B = 20$, classified accordingly to the configuration. Energies are in Kelvin. The horizontal line is the dissociation limit. Note that apart from the energy shift, the scales of the two plots are the same.




other shell-model states, but the mixing will probably not be constructed along the DMC stochastic procedure. Thence, the DMC physically relevant results are those with larger value of the binding energy for each row. The other configurations with smaller energy are possible interacting configurations, and presumably an optimized linear combination within each row could provide a better binding energy.

Note that among the states involving the 2s shell none of them has maximum binding except 1s2s :³S because the configuration is unique. For $N_B = 8$ this level is not bound, but it is the last bound one for $N_B = 20$. For the other cases the DMC optimized energies prefer 1s over 2s orbits, but 1s2s :³S survives because $1s^2$:³S violates the exclusion principle.

Concentrating on the normal shells (1s, 1p, 1d) it appears that the binding energies depend basically on the configuration, and are almost independent of the coupling within the configuration. Figs. 1 and 2 give an idea of this near independence, both for energies and radii. Within a central-field shell-model description, this fact indicates that the *residual interaction* between the two ³He atoms is very small, with energies close to the ones

FIG. 2: The root mean square radius (in Å) of bosons and fermions for two-fermion systems, referred to the full centerof-mass of the drop, as a function of the configuration. The two plots correspond to $N_B = 8$ (upper figure) and $N_B = 20$ (lower figure). The lines displaying an almost horizontal lines represent the boson radii, and the varying lines represent the fermion radii.



provided by the simple non-interacting picture

$$E(n_1\ell_1, n_2\ell_2) = E_0 + \epsilon_{n_1\ell_1} + \epsilon_{n_2\ell_2},$$

where E_0 is the energy of the boson core and $\epsilon_{n\ell}$ the separation energies defined in Eq. (7) and given in Table I. The corresponding values appear in the last row of Table III, with the exception of the last column, the 1p2sconfiguration which actually has evolved close to the 1s1pconfiguration.

The rough picture of the two-fermion drops as basically non-interacting fermions bound to a rigid ⁴He cluster will be refined in Section VI by introducing an effective monopole interaction.

V. CLUSTERS WITH MORE THAN TWO FERMIONS: BINDING ENERGIES

In this Section we consider selected states with a fixed number of bosons and an increasing number of fermions, up to 18 for normal clusters, *i.e.*, with fermions with spin up and down, and up to 9 for fully polarized clusters, with only spin up fermions.

A. The Slater determinant

The fermionic factor D_F must have good angular momentum and spin quantum numbers, and it must be properly antisymmetrized and translationally invariant. We take it to be a product of two Slater determinants, one for each spin orientation. The obvious way to build them up is with single-particle functions $\phi_{n\ell m}$ generated by a central field which dictates a natural filling order. In general, good total orbital angular momentum L and total spin S demand a linear combination of determinants. If we insist on a single product, the construction is quite cumbersome for shells with high ℓ (see *e.g.* Ref. 28), though simple when dealing with s and p shells.

The main problem in establishing a reasonable shellmodel description of a system containing ³He atoms, both in pure as well as in mixed drops, is the lack of phenomenological information about the central field. There are two familiar schemes common to other fermionic systems: the shell ordering 1s, 2s, 2p, 3s, 3p..., characteristic of atoms, and the harmonic oscillator sequence 1s, 1p, (2s, 1d), (2p, 1f)..., used in light and medium atomic nuclei. The principal quantum number follows different rules in both schemes. For the latter the parenthesis indicate degenerate orbitals.

In the previous Sections we have explained why the $1s, 1p, 1d, 2s, \ldots$ ordering was adopted for the calculations. The information comes from the analysis of a single ³He atom bound to a medium-size bosonic drop, which has been studied by density functional methods^{13,14} and by microscopic methods based either in variational wave functions^{15,17} or in DMC techniques.¹⁸ All these studies suggest that the fermion may be viewed as a particle bound to a potential well, centered in the bosonic drop, but which is appreciably different from zero (and attractive) only in a rather wide region near the surface of the drop, but goes to zero near the center of the drop as well as at long distances. This single-particle potential gives a special level ordering based on the orbital angular momentum, 1s, 1p, 1d, 1f..., with almost degenerate single-particle energies. Moreover, the fermion has a very small probability penetrating the boson drop. The same scheme results from the study of one ³He atom in liquid ⁴He, giving rise to the so-called Andreev states, as well as from the study of many ³He atoms attached to a large core of ⁴He atoms.²⁰ Note, however, that *intruder* levels, like 2s, may appear for a sufficiently large number of fermions and a sufficiently small number of bosons.

Among the possible wave functions related to a given configuration (the so-called *terms* in Atomic Physics) we have chosen the two simplest cases: a) maximum total spin on which maximum total orbital angular momentum is built; b) maximum orbital angular momentum on which maximum spin is built. The resulting wavefunctions are products of two determinants, one for each spin orientation. Other choices demand linear combination of Slater determinants.

As the exponential tail in the trial wave function has

the role of roughly confining the system, we can construct the Slater determinants considering only the angular momentum part of the single-particle functions, as well as the spin part. As mentioned above, we have used the harmonic polynomials $\phi_{\ell m}(\mathbf{r}) = r^{\ell} Y_{\ell m}(\Omega)$ as single-particle functions. The determinants so constructed are translationally invariant wave functions, in the sense that they only depend on the $3N_F - 3$ relative coordinates $\mathbf{r}_i - \mathbf{r}_j$. This fact is of crucial importance, particularly when describing systems with a small number of constituents.

The way of constructing the required determinants is very simple. Take for instance the maximum spin case. Once the innermost shells have been filled, the remaining ³He atoms occupy the $\phi_{\ell\ell}$, $\phi_{\ell\ell-1}$... spin up states, until the angular momentum states are exhausted; then the same procedure is followed to fill out the spin down states. The spin S of the resulting determinant has the maximum value allowed for the occupancy of the shell, and its orbital angular momentum is $L = |L_z|$. In general the value of the determinant is a complex number, not very adequate for the DMC algorithm. The solution is as in the two-fermion case: to use the sum or difference (whichever is non vanishing) of the determinants with $L_z = L$ and $L_z = -L$. The importance sampling wave function has well defined S, S_Z and L, but not L_z . Nevertheless, due to the rotational symmetry of the Hamiltonian, this has no influence on the energy values. However the need of computating determinants with complex matrix elements still remains, with the consequent slowing down of the numerical calculations.

By using this procedure we have calculated states with up to $N_F = 18$, corresponding to the complete filling of the three lowest shells. We have also considered states in which all spins are up, *i.e.*, fully polarized fermions, with the maximum number $N_F = 9$. The procedure to construct the determinants in this case is an obvious adaptation of the one described above.

In previous work^{5,8} we used a conventional *Cartesian* ordering, in particular $\{x^2, y^2, z^2, xy, xz, yz\}$, whereby the single-particle orbitals are a mixture of 2s and 1d wave functions. As a consequence the differences with the present calculations—associated to changes in the nodal structures—become significant when the *d*-shell starts to be filled.

B. Binding energies of normal clusters

Table IV presents the values of the binding energies corresponding to two situations, $N_B = 8$ and $N_B = 20$, for values of N_F from 0 to 18. All these values have been computed with the level ordering discussed above, and for two possible couplings: S_{max} , where in each shell particles are aligned to maximum spin S, and then to to maximum orbital angular momentum L; L_{max} , where particles are first aligned to maximum L, and then maximum S. In Table IV, whenever there are two entries for a given cluster, the upper row corresponds to S_{max} , and the lower one to L_{max} . Maximum spin is quite uniformly favoured but the splitting of the two computed levels is always smaller than 0.3-0.4 K.

TABLE IV: Binding energies (in K) of mixed clusters with 8 and 20 bosons and up to 18 fermions. Whenever there are two entries for a given cluster, the upper row corresponds to the S_{max} coupling: particles aligned to maximum S and then to maximum L. The lower row is for the L_{max} case: maximum L first and then maximum S. In the last column the results obtained in Ref. 8 for $N_B = 8$ are also displayed.

	$\int dx = 0 \text{ are also displayed.}$									
	N_F	Conf	L	S	$N_B = 8$	$N_B = 20$	$N_B = 8 \ [8]$			
	0		0	0	5.14(0)	33.76(1)	5.13(2)			
	1	$1s^1$	0	1/2	6.08(0)	35.55(1)	6.07(2)			
	2	$1s^2$	0	0	7.09(0)	37.32(1)	7.05(2)			
	3	$1p^1$	1	1/2	7.72(0)	38.88(1)	7.69(2)			
	4	$1p^2$	1	1	8.44(1)	40.47(2)	8.42(2)			
			2	0	8.40(1)	40.44(2)				
	5	$1p^3$	0	3/2	9.25(1)	42.14(1)	9.23(2)			
			2	1/2	9.20(1)	42.08(2)				
	6	$1p^4$	1	1	10.09(1)	43.72(3)	10.03(3)			
			2	0	10.04(1)	43.71(2)				
	7	$1p^5$	1	1/2	11.00(1)	45.40(2)	11.03(3)			
	8	$1p^6$	0	0	12.00(1)	47.07(2)	12.03(3)			
-	9	$1d^1$	2	1/2	12.49(1)	48.37(2)	12.33(3)			
	10	$1d^2$	3	1	13.02(1)	49.62(2)	12.74(3)			
			4	0	12.97(1)	49.64(4)				
	11	$1d^3$	3	3/2	13.65(1)	51.03(4)	13.20(4)			
			5	1/2		51.01(3)				
	12	$1d^4$	2	2	14.42(1)	52.46(5)	13.71(5)			
			6	0	14.19(1)	52.23(4)				
	13	$1d^5$	0	5/2	15.26(2)	53.99(2)	14.20(4)			
			6	1/2	14.96(2)	53.71(3)				
	14	$1d^6$	2	2	16.01(1)	55.37(4)	14.88(4)			
			6	0	15.74(2)	55.19(4)				
	15	$1d^7$	3	3/2	16.77(2)	56.83(3)	15.73(5)			
			5	1/2	16.64(1)	56.70(5)	. /			
	16	$1d^8$	3	1	17.63(2)	58.34(4)	16.55(4)			
			4	0	17.62(2)	58.29(4)				
	17	$1d^9$	2	1/2	18.64(3)	59.94(3)	17.44(13)			
	18	$1d^{10}$	0	0	19.74(3)	61.56(5)	18.49(5)			
-						. /	. /			

The present results for $N_F \leq 8$ should coincide, and indeed they do, with previous calculations based in the cartesian ordering of the single-particle states.⁸ Beyond $N_F = 8$ the calculations of Ref. 8 used an uncontrolled mixture of 1d and 2s states, as explained above, thus corresponding to different importance sampling functions. The last column of Table IV displays the results obtained in Ref. 8 for $N_B = 8$ clusters. One should keep in mind that both results are based on the DMC method within the fixed node approximation, so that in both cases the obtained energies are actually upper bounds to the real ones. The present binding energies for $N_F > 8$ are slightly higher that the previous results, the gain being of 0.16 K for $N_F = 9$ and monotonically increasing up to 1.40 K for $N_F = 18$, in the case of $N_B = 8$. This apparently modest increase (up to 6%) may be relevant for the boundaries of the stability chart of mixed drops. In any case, it gives support to the level ordering used

in the present work, as the associated importance sampling function provides a better variational bound than the previous ones.

FIG. 3: Binding energies (in K) for $N_B = 8$ and $N_B = 20$, as a function of the number of fermions N_F . The long horizontal lines correspond to normal clusters, and the short lines to polarized clusters. Note that apart from the energy shift, the scales of the two figures are the same.





FIG. 4: Separation energies, or ionization potentials, (in K) for normal S_{max} clusters as a function of the number of fermions N_F . The lower curve correspond to 8 bosons, and the upper curve to 20 bosons. Dashed lines are just an eye guide.



The binding energies of Table IV, for normal clusters,

have been plotted in Fig. 3, together with those for polarized clusters [to be discussed in Section VD (Table V)]. The scale is such that values for the normal clusters are superimposed.

Basically the energies grow linearly with the number of fermions but finer details emerge when looking at the fermion chemical potential

$$\mu_F(N_F) = E(N_B, N_F - 1) - E(N_B, N_F), \qquad (9)$$

which is plotted in Fig. 4 for two values of N_B and $N_F \leq 18$. Observe the sudden drop of μ_F after $N_F = 2$ and 8, corresponding to the closure of the 1s and 1p shells. The relative minimum appearing at $N_F = 14$ is at first somewhat puzzling. Its origin will become clear in Section VI.


C. The stability map revisited

As it has been mentioned in the introduction, one of the most appealing properties of the mixed He clusters is the existence of *instability islands*, namely regions around selected values of (N_B, N_F) in which the system is not bound. These regions were discovered after Many-Body computations based on self-adjustable variational functions constructed with Jastrow factors supplemented by 2p-2h and 3p-3h Configuration Interaction correlations.⁵ Afterwards, the calculation was refined by means of the DMC method,⁸ confirming the previous findings. Given that both calculations provide really with only upper bounds to the energy, and having observed the improvement of the present DMC approach, based upon a different ansatz for the determinantal part of the importance sampling guiding wave function, we have revised the previous calculation, just to check and eventually improve the limits of the instability regions.

Indeed, in the previous subsection we have seen that the present level ordering leads to a noticeable energy gain in the 1*d*-shell with respect to previous works. This fact suggests that some of the clusters previously qualified as *metastable*, *i.e.* systems with negative energy but less bound than clusters with a smaller number of fermions, could be in fact stable. Indeed, a new computation near the beginning of the 1*d*-shell indicates two new bound systems, namely the clusters $(N_B = 3, N_F = 11)$ and $(N_B = 4, N_F = 9)$. Special attention has been paid to the cluster $(N_B = 1, N_F = 18)$, corresponding to full 1s, 1p and 1d shells, but our finding is that this cluster is not bound.

The stable clusters are displayed in Fig. 5, which supersedes the results previously obtained in Ref. 8. From the experimental point of view, we suggest the study of the regions $N_B \leq 4$ and $N_F \leq 5$ to ascertain the stability limits. The measurements will require an improved mass resolution, at least 1 amu for clusters up to 25-30 amu.

FIG. 5: Stability map of mixed clusters. Solid squares represent truly bound states, and open squares represent metastable states.



D. Polarized mixed clusters

We have also considered fully polarized mixed systems. The values obtained for their binding energies are shown in Table V for $N_B = 8$ and 20. In both cases the number of fermions $N_F \leq 9$ is limited by the complete filling of the 1s, 1p and 1d orbits. The asterisk in Table V corresponds to situations in which the binding energy of the fully polarized system $E_{\uparrow}(N_B, N_F)$ is smaller than the energy $E(N_B, N_F - 1)$ of the normal cluster with one fermion less. In the other cases the results correspond to true bound states.

The energies of polarized clusters are compared with those of normal clusters in Figure 3. Polarized mixed drops are always less bound than the unpolarized cluster with the same number of fermions. In other words, they are excited states.

The energy differences between normal an polarized clusters take values around 0.25 K ($N_B = 8$) and 0.15 K ($N_B = 20$) per fermion. It is worth mentioning that theoretical calculations for liquid ³He provide a difference of around 0.10 K per particle at the equilibrium density and -0.10 K at densities close to the solidification one.³¹ In other words, the preferred phase would be the polarized one at high densities. This anomalous behavior has been interpreted as a side effect related to improper nodal surfaces for the unpolarized systems. The particle density of the fermionic phase in our case is much smaller than that of the fermionic liquid, and in consequence we cannot ascertain if our shell-model filling scheme will present such an anomalous crossing at higher densities.

In the same figure one may appreciate a sudden change for the differences $\Delta_E = E_{\text{normal}}(N_F) - E_{\text{pol}}(N_F)$ at $N_F = 4$, which has a simple interpretation: it corresponds to the filling of the 1*p* shell for the polarized case and the related jump in the ionization potential after adding a new fermion.

As for the unbound clusters, all of them have very high spin, and their binding energy is larger than the energy of the polarized cluster with one fermion less. Therefore it is very likely that the set of polarized states corresponds to a stable branch above the dissociation limit, analogous to the so called *displaced terms* in Atomic Physics.


The separation energies for the polarized clusters, displayed in Fig. 6, follow basically the same pattern than the separation energies for the normal clusters. Again

TABLE V: Binding energies (in K) for fully polarized clusters, for 8 and 20 bosons. The ³He single-particle configuration is indicated in the second column. Results marked with an asterisk correspond to clusters with a binding smaller than the cluster with a fermion less according to Table IV.

				0	
N_F	Config.	L	S	$N_B = 8$	$N_B = 20$
0		0	0	5.14(1)	33.76(1)
1	$1s^1$	0	1/2	6.08(1)	35.55(1)
2	$1s^11p^1$	1	1	6.65(1)	37.05(2)
3	$1s^{1}1p^{2}$	1	3/2	7.30(1)	38.61(2)
4	$1s^{1}1p^{3}$	0	2	8.06(1)	40.22(1)
5	$1s^11p^31d^1$	2	5/2	8.26(1) *	41.37(2)
6	$1s^11p^31d^2$			8.53(1) *	42.57(2)
7	$1s^11p^31d^3$	3	7/2	8.96(2) *	43.83(3) *
8	$1s^1 1p^3 1d^4$	2	4	9.53(2) *	45.19(3) *
9	$1s^11p^31d^5$	0	9/2	10.19(2) *	46.63(2) *

they recall the atomic ionization potentials with the sudden drop once a given shell is closed. Note that for polarized fermions the closure of shells occurs at $N_F = 1$ $(1s), N_F = 4 (1p)$ and $N_F = 9 (1d)$.

FIG. 6: Separation energies for polarized clusters.



VI. EFFECTIVE MONOPOLE INTERACTION ANALYSIS

In this section we shall treat the results of our calculations as data, and assume the validity of the shell-model scheme to find an effective one- plus two-body Hamiltonian that could reproduce them. This two-body part will only refer to fermions as the boson cluster will be assumed as a fixed core that generates the single particle energies in Table I, leading to a one body potential

$$U = \sum_{s} n_s \varepsilon_s$$

where n_s is the number of particles in shell s. The sum is extended to the occupied shells.

The two-body part is defined by matrix elements

$$V_{rs,tu}^{LS} = \langle rs : (LS) | V | tu : (LS) \rangle$$

where $|tu:(LS)\rangle$ is a two-particle state in shells t and s, coupled to orbital angular momentum L and spin S, properly antisymmetrized and normalized.

To simplify matters, the two-body part of the full Hamiltonian will be separated into monopole (m) and multipole (M) contributions $H = H_{mS} + H_M$. The monopole Hamiltonian H_{mS} is defined by the property of giving the *average* energy of configurations at fixed number of particles n_r and spin \mathbf{S}_r . The closed shells and the one-particle and one-hole states built on them are configurations with a single state (we call this set $cs \pm 1$). Hence, their energies are entirely given by H_{mS} . For the other Slater determinants entering our calculations H_m gives an average value that will be split by the multipole term H_M . We shall assume (and check) that the influence of H_M is small, and simply neglect it.

The extraction of effective interaction averages goes back in time, and an important earlier reference is the work of French.³² The form we shall use here came later³³ and was used to describe shell formation properties in nuclear physics.³⁴ A forthcoming review article³⁵ contains comprehensive information on formal properties of H_m and H_M .

The average matrix elements are defined as

$$V_{rs} = \frac{\sum_{LS} V_{rsrs}^{LS} (2L+1)(2S+1)(1+(-1)^{L+S} \delta_{rs})}{\sum_{LS} (2L+1)(2S+1)(1+(-1)^{L+S} \delta_{rs})}$$
$$V_{rs}^{S} = \frac{\sum_{L} V_{rsrs}^{LS} (2L+1)(1+(-1)^{L+S} \delta_{rs})}{\sum_{L} (2L+1)(1+(-1)^{L+S} \delta_{rs})}$$
(10)

where V_{rs} is the full (scalar) average of two-body matrix elements, whereas V_{rs}^{S} are (vector) averages at fixed S. It is convenient to introduce the following combinations

$$a_{rs} = \frac{1}{4}(3V_{rs}^1 + V_{rs}^0), \quad b_{rs} = V_{rs}^1 - V_{rs}^0$$
(11)

$$V_{rs} = a_{rs} - \frac{3}{4} \frac{\delta_{rs}}{D_r - 1} b_{rs}$$
(12)

where $D_r = 2(2l_r + 1)$ is the maximum number of particles in the shell. The standard result is then

$$H_{mS} = U + \sum_{r \le s} \frac{1}{(1 + \delta_{rs})} \bigg[a_{rs} n_r (n_s - \delta_{rs}) + b_{rs} \left(\mathbf{S}_r \cdot \mathbf{S}_s - \frac{3n_r}{4} \delta_{rs} \right) \bigg],$$
(13)

where \mathbf{S}_r is the total spin operator corresponding to the particles of shell r, $\mathbf{S}_r = \sum_{i \in r} \sigma_i/2$, and the $3n_r/4$ substraction ensures the two-body nature of H_{ms} by making the spin contribution vanish for *single particle* states. It has the drawback of producing a non-zero values at closed shells and single hole states. Therefore it is preferable to rewrite

$$H_{mS} = U + \sum_{r \le s} \frac{1}{(1+\delta_{rs})} \left[V_{rs} n_r (n_s - \delta_{rs}) + b_{rs} \left(\mathbf{S}_r \cdot \mathbf{S}_s - \frac{3n_r (D_r - n_r)}{4(D_r - 1)} \delta_{rs} \right) \right]$$
(14)

The counter-terms in the second line now ensure its vanishing at the closed shell as well as at one-particle and one-hole states. As a consequence, their energies are fully given by the first line in Eq. (14), which we refer to as H_m from now on. The advantage of this operation is that it decouples the determination of the V_{rs} and b_{rs} centroids, so that we can proceed with the former first, as they are the ones that give the global features.

In principle, the six necessary centroids— V_{ss} , V_{sp} , V_{sd} , V_{pp} , V_{pd} and V_{dd} —could be extracted from Table III. However, this parameter-free choice has large uncertainties and it is better to reserve it as a consistency check with the results of a more precise fit to the energies in Table IV, which we call E_i , i = 0, 18. It is very instructive to start doing the fit by hand, *i.e.*, step by step.

Upon filling, the closed shells become new "cores": $E_2 = E_{Cs}, E_8 = E_{Cp} \dots$ The single particle energies are taken from Table I. Then

 $E_2 = E_{Cs} = E_0 + 2\varepsilon_s + V_{ss}$. Extract V_{ss}

 $E_3 = E_{Cs} + \epsilon_p + 2V_{sp} \equiv E_{Cs} + \bar{\epsilon}_p$. Extract V_{sp}

To extract V_{pp} , we do not rely on E_4 , because it is not purely given in terms of centroids, but on

$$E_7 = E_{Cs} + 5\bar{\varepsilon}_p + 10V_{pp}$$
 or

$$E_8 = E_{Cp} = E_{Cs} + 6\bar{\varepsilon}_p + 15V_{pp}$$

The fit becomes overdetermined, signalling a problem with some basic assumption about the effective interaction, which we shall try to identify later. As of now let us settle for a compromise value of V_{pp} . The next step is $E_9 = E_{Cp} + \varepsilon_d + 2V_{sd} + 6V_{pd} \equiv E_{Cp} + \overline{\varepsilon}_d$, that determines $V_{sd} + 3V_{pd}$. As the two matrix elements will only appear in this linear combination, the number of parameters is reduced to five. Finally, for V_{dd} we have the same compromise problem we had for V_{pp} . To find reasonable values for V_{pp} and V_{dd} it was decided to do an overall fit of the five parameters. In principle, the idea does not seem very sound because three parameters are *apparently* well determined. As we shall see, this may not be the case, and the numerical fit will turn out to be sound.

The hand-made fit involves $cs \pm 1$ states that are common to the S_{max} and L_{max} cases in Table IV. In doing the numerical fit, only the S_{max} states were included. The results for the binding energies in the $N_B = 8$ clusters are given in Fig. 7. The agreement is quite excellent, but for the fully polarized case, also shown, there are significant discrepancies that can be cured by introducing the full H_{mS} through a single parameter $b_{rs} = b$ so that the contribution of the second line in Eq. (14) becomes $b[S(S+1)/2 - \sum_r 3n_r(D_r - n_r)/4(D_r - 1)]$. The results of the fit are given in Table VI. The fitted and calculated curves become nearly undistinguishable. As noted at the end of Section VB, the full energies are rather smooth patterns that tell us little about details. As a first approximation, Fig. 7 for the normal S_{max} clusters is reasonably well represented by a straight line, which would be the analogue of the famous Bethe-Weizsäcker "Liquid drop" formula for nuclei. The truly sensitive quantities are the separation energies (chemical potentials) in Eq. (9) and Fig. 4. And, indeed, the true test

FIG. 7: Comparison of DMC binding energies with the monopole Hamiltonian with spin (continuous line) and without spin terms (dashed line). Circles correspond to the computed DMC values for normal S_{max} clusters. The lower line corresponds to the fully polarized case, and the upper group to the normal clusters. In both cases is $N_B = 8$.




TABLE VI: The fitted centroids in K

	N_B	= 8	$N_B =$	= 20
	Value	Error	Value	Error
V_{ss}	0.073	0.006	-0.019	0.007
V_{sp}	0.079	0.002	0.080	0.003
V_{pp}	0.081	0.002	0.031	0.003
$(V_{sd} + 3V_{pd})/4$	0.078	0.001	0.045	0.001
V_{dd}	0.069	0.001	0.045	0.002
b	0.071	0.005	0.063	0.007

of the monopole description comes in Fig. 8. A smooth linear approximation to the binding energies would result in a constant. By introducing H_m there is an enormous improvement in that the shell effects at closures are well reproduced (dashed line). However, the more detailed pattern between closures demands the S(S + 1) term in the full H_{mS} : the agreement with DMC becomes truly quantitative (full line).

The numerical fit was made for the normal case we have called S_{max} , but Table IV contains another normal mode, L_{max} . As was noted, the hand-made fit is the same for both couplings, and it gives results that are almost as good as the numerical fit for S_{max} , and very good ones for L_{max} . But here the numerical fit also does a slightly better job, shown in Fig. 9, where H_{mS} is seen to reproduce beautifully the staggering pattern between S = 0 and S = 1/2 states (referred to as L_{max} case in Table IV). Here we are faced with some interesting physics: the numerical fit was chosen to find good compromise values for V_{pp} and V_{dd} , but it does slightly better than a fit restricted to those over-determined parameters. The hint is that the other three parameters are not as well determined as the "hand-made" fit suggests. The most likely reason is to be found in size effects: as fermions are added, the overall radius evolves, and for a self-bound system it should go asymptotically as $N_F^{1/3}$. [A strong indication in this sense will be found in

FIG. 8: Comparison of calculated separation energies with the monopole Hamiltonian with spin (H_{mS} , continuous line) and without spin terms (H_m , dashed line), for S_{max} states. Circles correspond to the computed DMC values. The lower group corresponds to $N_B = 8$, and the upper group to $N_B = 20$.





Fig. 10.] Therefore, the effective matrix elements should also evolve in a way our simplified H_{mS} ignores: The numerical fit then emerges as the sound and natural way to define a best compromise value, not only for V_{pp} and V_{dd} but for all the monopole parameters.

FIG. 9: Same as Fig. 8 but for clusters with L_{max} .



Finally, Table VII compares the centroids V_{rs}^0 and V_{rs}^1 obtained with the fit to the 19 normal S_{max} clusters with the values obtained directly from the DMC calculation of two-fermion states, again with a very good agreement (within the large errors of about 0.04 K associated to the latter), thus confirming the consistency of the effective interaction interpretation.

It appears that very hard DMC calculations lead to results amenable to a very simple and cogent interpretation in terms of the monopole Hamiltonians H_{mS} . The S(S + 1) contribution is particularly interesting: Fig. 7 suggests the idea that for large enough number of fermions the polarized clusters could become ground states. Though this is only a speculation, it may be also be taken as a strong invitation to push the study of mixed clusters much further.

TABLE VII: The two-fermion centroids as obtained from the effective interaction (columns labelled Eff) compared with the DMC computed values.


	$N_B = 8$								
	S=	=0	S=1						
	Eff	DMC	Eff	DMC					
\mathbf{SS}	7.09	7.09							
sp	6.57	6.61	6.64	6.65					
pp	6.10	6.14	6.17	6.19					
sd	5.94	5.99	6.02	6.01					
pd	5.46	5.49	5.54	5.51					
dd	4.84	4.86	4.91	4.94					
		$N_B =$	20						
	S=	=0	S=1						
	Eff	DMC	Eff	DMC					
\mathbf{SS}	37.32	37.33							
$^{\mathrm{sp}}$	36.97	37.06	37.04	37.05					
pp	36.58	36.73	36.64	36.74					
sd	36.44	36.47	36.50	36.49					
pd	36.04	36.15	36.10	36.19					
dd	35.54	35.54	35.61	35.70					

VII. THE SHAPE OF MIXED CLUSTERS

A. Normal mixed clusters

In this section we present several figures related to the shape of mixed clusters. In Fig. 10 there are the values of the root mean square radii for bosons and fermions referred to the center-or-mass of the cluster, for the selected cases $N_B = 8$ and 20. There are some fluctuations, probably related to the use of the mixed estimator method to compute these radii, and thus depending on the quality of the importance sampling wave function. Apart from these fluctuations, the most noticeable properties which emerge from these plots are the almost constancy of the bosonic radii and the smooth growing of the fermionic radii. This manifests clearly the representation of the cluster as a quite rigid bosonic core with an halo of fermions.

FIG. 10: The values of the root mean square radii (in Å) for bosons (squares) and fermions (circles) referred to the centerof-mass of the cluster, as a function of the number of fermions N_F in the cluster. The number of bosons are fixed to $N_B = 8$ and 20.



This picture is confirmed by the plots of Fig. 11, where

the one-body distributions of bosons and fermions with respect to the center of mass of the mixed cluster are displayed. These distributions are given by

$$\rho_M(r) = \langle \Psi | \sum_{i=1}^{N_M} \delta(\mathbf{r} - [\mathbf{r}_i - \mathbf{R}]) | \Psi \rangle,$$

where label M stands for B (bosons) or F (fermions), N_M is the number of atoms of the given species, and Ris the center-of-mass of the full drop. Given that these distributions may have an angular dependence for open shells, we have computed the spherical average of them. The distributions are normalized to the number of particles of a given species,

$$\int \rho_M(r) d\mathbf{r} = N_M.$$

We observe in Fig. 11 that bosons are located in the same central region, being slightly compressed as the number of fermions increases. This shrinking is more important for the light $N_B = 8$ cluster, and is almost negligible when $N_B = 20$. With respect to the distributions of fermions, they are clearly located at the surface of the bosonic subcluster, with a small penetration near the center of the drop in the case of $N_B = 8$, more important for larger values of the number of fermions. In the case of $N_B = 20$ the dominating picture is that of a rigid core of bosons with a fermionic halo.

A complementary information about the shape of clusters is provided by the two-body distributions,

$$\rho_M(\mathbf{r}, \mathbf{r}') = \frac{2}{N_M(N_M - 1)} \langle \Psi | \sum_{i < j}^{N_M} \delta(\mathbf{r} - \mathbf{r}_i) \delta(\mathbf{r}' - \mathbf{r}_j) | \Psi \rangle,$$

which is normalized to 1.

Because of the finite size of the system under consideration, this distribution function depends on two coordinates, \mathbf{r} and \mathbf{r}' , or, equivalently, on the distance of the center-of-mass of the pair $((\mathbf{r} + \mathbf{r}')/2)$ to the center-of-mass of the system and the relative distance $(\mathbf{r} - \mathbf{r}')$ of the two particles, thus producing a function very difficult to plot. In order to get a more friendly quantity we have averaged the above two-body distribution with respect to its center-of-mass and for the remaining dependence we have computed the spherical average. The reduced pair distribution so obtained is now normalized to 1,


$$\int \rho_{12}(r)d\mathbf{r} = 1$$


B. Polarized mixed clusters

There are noticeable similarities between the distribution functions related to polarized systems and those corresponding to normal clusters. The subcluster of bosons

FIG. 11: The density distributions (in Å⁻³) of bosons and fermions with respect to the center-of-mass of the cluster, for the two selected $N_B = 8$ and 20 cases.

FIG. 12: The boson-boson and fermion-fermion distributions (in Å⁻³), for the two selected cases $N_B = 8$ and 20.






is again hard, its radius being practically independent of the number of spin-aligned fermions N_F . Differences with respect to the unpolarized cluster appear when comparing the root mean square radii of fermions, as shown if Fig. 13 where the fermion halo in the polarized clusters is larger than in the unpolarized case.

Something similar happens with the density distributions of bosons or fermions with respect to the center of mass. The former remains basically unaltered when the number of fermions grows, and the latter follows the same pattern that in the case of unpolarized clusters.

There is, however, a remarkable fact in connection with the two-body distributions and, specifically, the fermionfermion distributions. These distributions are shown in Fig. 14, for the two selected cases N = 8 and 20. As can be seen there, the fermion-fermion distributions are very different from those obtained for the normal systems: the rise near 4 Å is much less pronounced and the range is much larger, suggesting that the fermions are either less correlated or subjected to a long-range correlation.

In order to appreciate the differences in the density distribution functions for normal and polarized clusters, we have plotted them in Figs. 15 for $N_F = 9$ and $N_B = 8$. Apart from a clear change in the shape, the peak of the

FIG. 13: The values of the root mean square radii (in Å) referred to the center-of-mass, for bosons (filled squares) and fermions (filled circles for normal clusters and open circles for polarized clusters) as a function of the number of fermions N_F in the cluster. The number of bosons is fixed to $N_B = 8$ (left panel) and to 20 (right panel).



normal case is close to 4 Å, while for the polarized cluster it is close to 7 Å.

VIII. SUMMARY AND CONCLUSIONS

This work has been devoted to a detailed analysis of clusters made of ${}^{4}\text{He}$ and ${}^{3}\text{He}$ atoms. This new analysis

FIG. 14: The fermion-fermion distributions (in Å⁻³), for $N_B = 8$ and unpolarized (top) and fully polarized (bottom) clusters.





FIG. 15: Comparison of the fermion-fermion distributions (lower panel), for clusters with $N_B = 8$ and $N_F = 9$. The dashed line corresponds to the polarized cluster and continuous line to the normal cluster.



was motivated by the improved knowledge of the ordering of single-particle orbits, obtained after a systematic study of the spectrum of a single ³He atom bound to a core of ⁴He atoms,¹⁸ which is expected to provide the optimal importance sampling trial function for the DMC calculation. The study has concentrated on clusters having a sufficiently large number of ⁴He atoms so as to offer a simplified pattern: the bosonic constituents arrange themselves as a quite rigid core whereas the fermionic atoms are distributed in the surface of the bosonic subcluster, producing a halo. This arrangement, previously obtained by means of density functional methods, has been confirmed and pushed down to systems with a small number of constituents.

One of the primary aims was to check the previously obtained stability map,⁸ after the optimization of the importance sampling function, as well as the improvement of the fermionic nodal surfaces. No significant change occurred, and large instability islands are still predicted for a small ($N_B \leq 3$) number of ⁴He atoms.

The determination of correlation functions, particularly the fermion-fermion distribution functions, as well as the analysis in terms of an effective interaction model suggests that the residual interactions between the fermions is very weak, of the order of the computational precision achieved (near 0.1 K). This fact is also reflected in the insensitivity of the energies to the (L, S) quantum numbers, the spectrum being essentially determined by the configuration.

In addition to the *normal* fermionic phase, with a small value of the spin, we have also analyzed the possibility of having a *ferromagnetic* phase, with all spins aligned. The normal phase is energetically favored but the ferromagnetic one gives rise to bound states, even for a moderately large number of ³He atoms. For example, a cluster with 20 bosons is able to bind up to six fermions, but beyond that number the system is above the dissociation limit. Nevertheless, because of the large values of the spin for the ferromagnetic phase, one may expect these unbound states to be long-lived, like in the case of polarized liquid ³He, and thus to be experimentally detectable. Perhaps by circulating bosonic clusters through a cold atmosphere of polarized ³He atoms one could create these spin-aligned states sticking ³He atoms one-by-one to the bosonic seed.

The mixed systems may have a very rich excitation spectrum, because of the gap between the normal and the polarized phase. For example, for $N_B = 20$ and $N_F = 6$ there is a difference of 1 K between the normal state (configuration $1s^21p^4$) and the polarized state (configuration $1s^{1}1p^{3}1d^2$). The analysis of the intermediate filled configurations, such as $1s^21p^{3}1d^1$ or $1s^21p^21d^2$ puts heavy demands on the DMC algorithm but it would be relatively simple in the density-functional method. Though it does not seem possible with the present experimental techniques to measure the spectrum, it is worth remembering that the excitation spectrum plays a relevant role in the production abundances of clusters.^{36,37}

Finally one should stress the power of the effective monopole interaction analysis that points to the basic simplicity of the DMC results and invites to take on the challenge of unearthing the deep reasons of such simplicity.

Acknowledgments

This work has been supported by MCyT/FEDER (Spain), grant number FIS2004-00912, GV (Spain), grant number GV2003-002 and MIUR (Italy), cofin-2001025498. Part of this study was done while RG was

- ¹ K.B. Whaley (editor), Special issue of J. Chem. Phys. **115** 10065-10281 (2001).
- ² E. Krotscheck and J. Navarro (editors), Advances in Quantum Many-Body Theory, Vol. 4 (2002), World Scientific (Singapore).
- ³ M. Barranco, J. Navarro and A. Poves, Phys. Rev. Lett **78**, 4729 (1997).
- ⁴ R. Guardiola and J. Navarro, Phys. Rev. A **71**, 035201 (2005).
- ⁵ R. Guardiola and J. Navarro, Phys. Rev. Lett. **89**, 193401 (2002).
- ⁶ D. Bressanini and G. Morosi, Phys. Rev. Lett. **90**, 133401 (2003).
- ⁷ D. Bressanini, G. Morosi, L. Bertini, and M. Mella, Few-Body Syst. **31**, 199 (2002).
- ⁸ R. Guardiola and J. Navarro, Phys. Rev. A 68, 055201 (2003).
- ⁹ W. Schöllkopf and J.P. Toennies, Science **266**, 1345 (1994).
- ¹⁰ W. Schöllkopf and J.P. Toennies, J. Chem. Phys. **104**, 1155 (1996).
- ¹¹ R.E. Grisenti, W. Schöllkopf, J.P. Toennies, G.C. Hegerfeldt, T. Köhler, and M. Stoll, Phys. Rev. Lett. 85, 2284 (2000).
- ¹² J.P. Toennies, private communication.
- ¹³ F. Dalfovo, Z. Phys. D **14**, 263 (1989).
- ¹⁴ M. Barranco, M. Pi, S.M. Gatica, E.S. Hernández, and J. Navarro, Phys. Rev. B 56, 8997 (1997).
- ¹⁵ A. Belić, F. Dalfovo, S. Fantoni, and S. Stringari, Phys. Rev. B **49**, 15253 (1994).
- ¹⁶ D. Bressanini, M. Zavaglia, M. Mella, and G. Morosi, J. Chem. Phys. **112**, 717 (2000).
- ¹⁷ E. Krotscheck and R. Zillich, J. Chem. Phys. **115**, 10161 (2001).
- ¹⁸ S. Fantoni, R. Guardiola, and J. Navarro, Phys. Rev. A 70, 023206 (2004).
- ¹⁹ M. Pi, R. Mayol, and M. Barranco, Phys. Rev. Lett. 82, 3093 (1999).

visiting professor at Universit Louis Pasteur in Strasbourg. One of us (RG) acknowledges financial support of the Secretaría de Estado de Educación y Universidades (Spain), Ref. PR2003-0374, as well as DEMOCRITOS by his hospitality.

- ²⁰ J. Navarro, A. Poves, M. Barranco, and M. Pi, Phys. Rev. A **69**, 023202 (2004)
- ²¹ S. Nakaichi, T.K. Lin, Y. Akaishi, and H. Tanaka, J. Chem. Phys. **71**, 4430 (1979).
- ²² R.A. Aziz, F.R. McCourt and C.C.K. Wong, Mol. Phys. 61, 1487 (1987).
- ²³ P.J. Reynolds, D.M. Ceperley, B.J. Alder and W.A. Lester Jr., J. Chem. Phys. **77**, 5593 (1982).
- ²⁴ J.W. Moskowitz, K.E. Schmidt, M.A. Lee and H.M. Kalos, J. Chem. Phys. **77**, 349 (1982).
- ²⁵ R.P. Feynman and M. Cohen, Phys. Rev. **102**, 1189 (1956).
- ²⁶ K.E. Schmidt, M.A. Lee, M.H. Kalos, and G.V. Chester, Phys. Rev. Lett. 47, 807 (1981).
- ²⁷ V.R. Pandharipande, S.C. Pieper, and R.B. Wiringa, Phys. Rev. B **34**, 4571 (1986).
- ²⁸ E.U. Condon and G.H. Shortley, *The Theory of Atomic Spectra*, (Cambridge University Press, Cambridge, 1951). See Table I, p. 208.
- ²⁹ J.W. Moskowitz, K.E. Schimdt, M.A. Lee, and H.M. Kalos, J. Chem. Phys. **77**, 349 (1982).
- ³⁰ P.J. Reynolds, D.M. Ceperley, B.J. Alder, and W.A. Lester Jr., J. Chem. Phys. **77**, 5593 (1982).
- ³¹ F.H.Zong, D.M.Ceperley, S.Moroni, and S. Fantoni, Mol. Phys. **201**, 1705 (2003).
- ³² J. B. French, in *Isospin in Nuclear Physics*, edited by D. H. Wilkinson (North Holland, Amsterdam, 1969).
- ³³ A. Abzouzi, E. Caurier, and A. P. Zuker, Phys. Rev. Lett. 66, 1134 (1991).
- ³⁴ J. Duflo and A. P. Zuker, Phys. Rev. C **59**, R2347 (1999).
- ³⁵ E. Caurier *et al.*, The Shell Model as Unified View of Nuclear Structure, 2003, nucl-th/0402046, to appear in RMP.
- ³⁶ L.W. Bruch, W. Schöllkopf, and J.P. Toennies, J. Chem. Phys. **177**, 1544 (2002).
- ³⁷ R. Brühl, R. Guardiola, A. Kalinin, O. Kornilov, J. Navarro, T. Savas, and J.P. Toennies, Phys. Rev. Lett. **92**, 185301 (2004).