

ON THE ORIGIN OF THE QUANTUM MECHANICS

JAUME GINÉ

ABSTRACT. Action at distance in Newtonian physics is replaced by finite propagation speeds in classical post-Newtonian physics. As a result, the differential equations of motion in Newtonian physics are replaced by functional differential equations, where the delay associated with the finite propagation speed is taken into account. Newtonian equations of motion, with post-Newtonian corrections, are often used to approximate the functional differential equations. Are the finite propagation speeds the origin of the quantum mechanics? In this work a simple atomic model based on a functional differential equation which reproduces the quantized Bohr atomic model is presented.

1. INTRODUCTION

Newtonian forces (for example, the inverse square law for gravitation) imply “action at distance”. This absurd, but outstandingly successful, premise of Newtonian theory predicts that signals propagate instantaneously. In classical physics, relativity theory postulates that signals propagate with a velocity that does not exceed the velocity of light. Thus, the forces of Newtonian physics must be replaced by force laws that take into account the finite propagation speed of the classical fields which determine the forces acting on a moving body. In turn, the ordinary or partial differential equations of Newtonian physics, which are derived from the second law of motion $m\mathbf{a} = \mathbf{F}$, must be replaced by corresponding functional differential equations where the force \mathbf{F} is no longer a function of just position, time, and velocity; rather, the classical force law must take into account the time delays due to the finite propagation speed of the classical fields.

The functional differential equations of motion for classical field theory are generally difficult, often impossible, to express in a form that is

1991 *Mathematics Subject Classification.* Primary 34C05. Secondary 58F14.

Key words and phrases. quantum theory, retarded systems, functional differential equations, limit cycle.

The author is partially supported by a DGICYT grant number BFM 2002-04236-C02-01 and by DURSI of Government of Catalonia “Distinció de la Generalitat de Catalunya per a la promoció de la recerca universitària”.

amenable to analysis. Thus, in order to obtain useful dynamical predictions from realistic models, it is frequently to replace the functional differential equations of motion by approximations that are ordinary or partial differential equations, see [3]. The purpose in these works is to discuss some of the mathematical issues that must be addressed to obtain a rigorous foundation for the post-Newtonian dynamics, that is, Newtonian dynamics with relativistic corrections, see for instance [3] and the references therein. For the electromagnetic classical field, in the ideal case of a point-charge particle, the resulting retarded potentials are the Liénard–Wiechert potentials. For the gravitational classical field we must use the Einstein’s field equation. Simple models of these equations are the subject of current research. The basic idea of post-Newtonian approximation, from a mathematical point of view, is the expansion of model equations in powers of $1/c$. From a physical point of view, the idea is to consider low velocity (compared with the speed of light). Note, for example, that the relativistic form of Newton’s second law, where the rate of change of the momentum is given by

$$\frac{d}{dt} \left(mv \left(1 - \frac{|v|^2}{c^2} \right)^{-1/2} \right),$$

reverts to Newton’s law in the low-velocity limit.

According to Maxwell’s field equations, a charged particle produces electromagnetic fields as it moves. Since, in this case, a particle radiates energy, it must slow down. In the theory of the electron point charge by considering motion along in a line, Dirac propose a self-force (the radiation reaction force) given by

$$F_{\text{self}} = \frac{2q^2}{3c^3} \ddot{x},$$

which is the half difference of the retarded and advanced forces, where q is the charge of the electron, see [11]. Therefore, a post-Newtonian model for the motion of an electron, confined to move on a line and with radiation reaction taken into account, is given by the Abraham–Lorentz equation

$$(1) \quad m\ddot{x} = \frac{2q^2}{3c^3} \ddot{x} + F,$$

where F is an external force. Since the electron radiates (produces fields that carry energy) the self force should cause the particle to lose energy and slow down. That’s why, the presence of the third derivative term in the first differential equation is called radiation damping. However, in these post-Newtonian models (where the differential equations are

not of second order) the “runaway” solutions appear, see [11]. For instance, in absence of external forces, equation (1) reduces to

$$m\ddot{x} = \frac{2q^2}{3c^3}\ddot{\ddot{x}},$$

and this equation has the solution $\dot{x} = C$ where C is an arbitrary constant, and other solutions where the acceleration is proportional to $\exp(3mc^3t/(2q^2))$. Hence, the acceleration grows indefinitely with time. This means that, a charge which goes out of a field, when leaving it, must self-accelerate indefinitely; which is an absurd. These runaway solutions are clearly not physical. What do they represent? How should they be eliminated? and What is the correct Newtonian equation with the radiation damping taken into account?

The mathematical answer to all these questions is approached in [3] and the subsequent works [5, 6], where these post-Newtonian models are recognized as singularly perturbed Newtonian equations. In order to recover the correct Newtonian equations with the post-Newtonian corrections, the Fenichel’s geometric singular perturbation theory is applied (in particular, the reduction to the slow–manifold). These Newtonian equations with the post-Newtonian corrections give physically reasonable dynamics; in particular, the runaway solutions are eliminated. Anyway, how can we justify using these models? Note, for instance, that the slow-manifolds in these models are unstable; nearby runaway solutions. In applied mathematics, we usually justify approximations by their stability. To validate the slow–manifolds reductions it must be shown that the resulting Newtonian model equations are stable with respect to the dynamics of the original functional differential equations, the true equations of motion in classical physics. Therefore further investigations are required in this direction for the study of the delay equations.

However, it is interesting to note that the presence of a small delay in a conservative system often results in damped long–term dynamics on an associated inertial manifold, see [3, 4]. For example, the Duffing–type model equation

$$\ddot{x} + \omega^2 x = -ax(t - \tau) + bx^3(t - \tau),$$

with small delay τ in the restoring force, reduce (by a formal computation to first order in τ) to the van der Pol–type model equation

$$\ddot{x} + \tau(3bx^2 - a)\dot{x} + (a + \omega^2)x - bx^3 = 0,$$

on its inertial manifold. As it has been noticed in [3] this example illustrates a phenomenon that is reminiscent of quantization: while most periodic solutions in one parameter families of periodic solutions

in a conservative system disappear in the presence of a small delay, some persist as limit cycles. The author of [3] asks himself whether this observation has a physical significance.

The solutions of the functional differential equations can, however, admit an infinite discrete spectrum. For example, we consider the retarded harmonic oscillator, given by the linear, second order, retarded functional differential equation

$$\ddot{x} + x(t - \tau) = 0,$$

with small delay τ . In [16] it is showed that this equation exhibits an infinite spectrum of discrete frequencies. Its general solution is a convergent linear combination of oscillations at an infinity of discrete (“quantized”) frequencies. As in quantum mechanics, in order to determine a unique solution an initial function needs to be known. The above consideration remains valid for any linear functional differential equation with constant coefficients and constant delay. Moreover, the locally linear approximation suggests that such “quantization” is also to be expected for non-linear functional differential equations.

At the beginning of the 20th century, Planck [12] initiated the quantum mechanics with his contribution to the black body radiation. Einstein [8], following the ideas of Planck [12], contributed to the development of the theory of quanta which is the embryonic step needed to arrive to the Quantum physics theory. It is interesting to note that Poincaré had been implicated in the discussion of the quantum theory, but the premature death of Poincaré deprives of its contributions in this theory, see for instance [14, 15]. In fact, Poincaré participated in the first congress of Solvay in October of 1911 and he died in July of 1912. One of the interesting known phenomenon studied by Poincaré is the concept of limit cycle, see [13]. This phenomenon does not occur in the Hamiltonian systems studied by standard physics theories. It only appears in systems with friction, i.e., systems with dissipative energy.

It is also interesting to note that one of the problems which originates the quantum theory was the problem that appears when the idea of the planetarium system is applied to the atomic model. This idea was proposed by Rutherford in 1911 [18] as a consequence of the experimental results obtained when bombing an atom with α particles. The problem in the model of Rutherford was that the charged electrons are accelerated in their movement around the nucleus and by the electromagnetic classical theory any accelerated charged body radiates energy. The problem of the atomic stability was initially solved by Bohr in 1913 [1] and it marks the success of the quantum theory and its posterior development. The atomic model of Bohr predicts the

radiation spectrum of certain atoms and the quantization of the energy in different energy levels is then obtained.

If you see the development of the quantum theory from the initial contributions, it is evident that each step is made with extra assumptions. For instance, the introduction of the quanta in the radiation of a black body and in the photo-electric effect, cf. [8], the quantization of the energy in the movement of an electron which moves as a harmonic oscillator under the influence of a harmonic restoring force, cf. [12]. Another example is the quantization of the angular orbital impulse of an electron in an atom, although the electron in an atom is accelerated in its movement around the nucleus. In this last case, it is assumed that this electron does not radiate energy, see [1]. However, we notice the difference between the Bohr quantization of the angular orbital impulse of an electron, which moves under the Coulomb force ($L = n\hbar$ for $n = 1, 2, 3, \dots$, where \hbar is a multiple of the Planck constant h divided by 2π), and the Planck quantization of the energy of a particle, as an electron, which moves as a harmonic oscillator ($E = nh\nu$ for $n = 1, 2, 3, \dots$). In fact the quantization of the angular orbital impulse of an electron leads to the quantization of the total energy but with an equation quite different than the Planck equation.

In [9] it is showed that the intrinsic phenomenon (the quantization of the energy) that appears in the first and simple systems initially studied by the quantum theory as the harmonic oscillator and the movement of a charged particle under the Coulomb force, can be obtained from the study of dissipative systems. In other words, it is showed that the phenomenon of the quantization of the energy of a particle which moves as a harmonic oscillator can be obtained via a classical system of equations. The same assertion also applies to the phenomenon of the quantization of the energy of a charged particle which moves under the Coulomb force and which loses and wins energy (for example if we consider the classical case where the electron radiates and absorbs energy from the electric field of the nucleus). Therefore, these phenomena are not intrinsic of the quantum theory, but also appear in classical systems. In fact, they appear in the qualitative theory of differential equations developed by Poincaré from 1881 [13].

Nevertheless, the most important problem is to find the exact form of the dissipative term and the interpretation of its physical meaning, see [9]. The retarded case, already explicitly incorporates certain subtle mathematical features of electrodynamics and relativity noticed by Poincaré, but overlooked by Einstein and subsequent researchers. Based on the study of the retarded systems, a simple atomic model

given by a functional differential equation which reproduces the quantized Bohr atomic model is presented in this paper.

The paper is not at all an alternative to the quantum theory, because the large development of the quantum theory in all the past century, the success in all its predictions, is outside of all doubt. On the other hand, the proposed model does not reflect the whole rich behavior of the quantum modern theories developed from 1925 by Schrödinger [19, 20, 21], Born [2], Heisenberg [10], Dirac [7], and others. The goal of the paper is to ask if the finite propagation speeds is the origin of the quantum mechanics.

To begin with, in [16], it is assumed that the two particles are rotating rigidly in circular orbits around a common center of masses. Moreover, a force which varies inversely as the square of the retarded distance is considered. The retarded distance is the distance from the current position of the electron to the "last seen" position of the proton. The simple expression for the force helps us to intuitively understand the consequences of a delay and under such circumstances, the angular momentum cannot be conserved. Thus, we have the astonishing situation that, purely under the action of internal forces, the system suffers a net torque. Now, the radiation term is introduced. But, the exact form of the radiation damping term is not clear. Finally, the simple heuristic case of the retarded inverse square force is used, to determine whether there can be a balance of forces between the delay torque and the 3rd order radiation damping. A total success does not take place and a value of r which is smaller than the Bohr radius is obtained. Nevertheless, the author, in [16], affirms that further investigations are required to determine the exact effects of radiative damping, and that it was prematurely concluded that radiative damping makes the classical hydrogen atom unstable. In the following section we present a simple atomic model based on a functional differential equation which reproduces the quantized Bohr atomic model.

2. THE RETARDED ELECTRODYNAMIC 2-BODY PROBLEM

We consider two particles interacting through the retarded inverse square force. The force on the electron exerted by the proton is given by

$$(2) \quad \mathbf{F} = K \frac{e^2}{r^3} \mathbf{r}.$$




FIGURE 1. The retarded electrodynamic 2-body problem.

The force acts in the direction of the 3-vector \mathbf{r} , along which the proton is "last seen" by the electron. The 3-vector \mathbf{r} may be represented by

$$\mathbf{r} = \mathbf{r}_p(t - \tau) - \mathbf{r}_e(t),$$

where $\mathbf{r}_p(t)$ and $\mathbf{r}_e(t)$ denote respectively the instantaneous position vectors of the proton and electron, respectively, at time t , and τ is the delay, so that $\mathbf{r}_p(t - \tau)$ is the "last seen" position of the proton. Assuming that the two particles are in rigid rotation with constant angular velocity ω , and referring back to Fig. 1, we have, in 3-vector notation,

$$\mathbf{r}_e = r_1[\cos \omega t \hat{\mathbf{i}} + \sin \omega t \hat{\mathbf{j}}],$$

and

$$\mathbf{r}_p = -r_2[\cos \omega(t - \tau) \hat{\mathbf{i}} + \sin \omega(t - \tau) \hat{\mathbf{j}}].$$

Hence, the 3-vector \mathbf{r} is given by

$$\mathbf{r} = [-r_2 \cos \omega(t - \tau) - r_1 \cos \omega t] \hat{\mathbf{i}} + [-r_2 \sin \omega(t - \tau) - r_1 \sin \omega t] \hat{\mathbf{j}},$$

Now, we introduce the polar coordinates (r, θ) and define the unitary vectors $\mathbf{l} = \cos \theta \hat{\mathbf{i}} + \sin \theta \hat{\mathbf{j}}$ and $\mathbf{n} = -\sin \theta \hat{\mathbf{i}} + \cos \theta \hat{\mathbf{j}}$. By straightforward calculations it is easy to see that the components of the force (2) in the polar coordinates are

$$F_r = K \frac{e^2}{r^3} \mathbf{r} \cdot \mathbf{l} = (-r_2 \cos(\omega\tau) - r_1) K \frac{e^2}{r^3}$$

and

$$F_\theta = K \frac{e^2}{r^3} \mathbf{r} \cdot \mathbf{n} = r_2 \sin(\omega\tau) K \frac{e^2}{r^3}$$

The equations of the movement are

$$(3) \quad m\ddot{r} - mr\dot{\theta}^2 = F_r,$$

$$(4) \quad mr\ddot{\theta} + 2m\dot{r}\dot{\theta} = F_\theta.$$

The second equation (4) can be written in the form

$$(5) \quad \frac{1}{r} \frac{dL}{dt} = \frac{1}{r} \frac{d}{dt}(mr^2\dot{\theta}) = F_\theta = r_2 \sin(\omega\tau) K \frac{e^2}{r^3}.$$

In 1913 Bohr [1] introduced the quantization of the angular momentum of the form $L = nh/(2\pi)$ where h is the Planck constant. If we accurately study equation (5) we see that the analytic function $\sin(\omega\tau)$ has a numerable number of zeros

$$(6) \quad \omega\tau = k\pi,$$

with $k \in \mathbb{Z}$, which are stationary orbits of the system of equations (3) and (4). When $\omega\tau \neq k\pi$ we have a torque which conduces the electron to the stationary orbits without torque with $\omega\tau = k\pi$. On the other hand in a first approximation the delay τ can be equal to r/c (the time that the field uses to goes from the proton to the electron at the speed of the light). From equation (6) we have

$$(7) \quad \tau = \frac{k\pi}{\omega} = \frac{r}{c}.$$

Taking into account that $\omega = v_\theta/r$, from (7) we have $v_\theta/c = k\pi$. But from the theory of relativity we know that $v_\theta/c < 1$, then we must introduce a new constant g in the delay. Hence, $\tau = gr/c$ and the new equation (7) is

$$(8) \quad \tau = \frac{k\pi}{\omega} = \frac{gr}{c},$$

and now $v_\theta/c = k\pi/g$, i.e. $v_\theta = k\pi c/g$ and from (8) we also have $r = k\pi c/(g\omega)$. In our model case of a classical rigid rotation we have $\theta = \omega t$ with $\omega > 0$. Therefore, $\dot{\theta} = \omega$ and $\ddot{\theta} = 0$. Hence, equation (4) for $\omega\tau = k\pi$ is

$$2m\dot{r}\omega = 0,$$

which implies $\dot{r} = 0$ and $r = r_k$ where r_k is a constant for each k . On the other hand, equation (3) for $\omega\tau = k\pi$ takes the form

$$(9) \quad -mr\dot{\theta}^2 = -m\frac{v_\theta^2}{r} = (-r_2(-1)^n - r_1)K\frac{e^2}{r^3} \approx -rK\frac{e^2}{r^3},$$

assuming that τ is small and then $r \sim r_1 + r_2$. From the definition of the angular momentum we have for $\omega\tau = k\pi$

$$(10) \quad L_n = mr_n^2 \dot{\theta} = mr_n^2 \omega = mr_n v_\theta = m \frac{k\pi c}{\omega g} \frac{k\pi c}{g} = m \frac{k^2 \pi^2 c^2}{\omega g^2}.$$

Now we compare with the quantization introduced by Bohr in 1913 and we obtain

$$m \frac{k^2 \pi^2 c^2}{\omega g^2} = \frac{nh}{2\pi}.$$

Therefore we have (identifying $k^2 = n$) that Planck constant satisfies the relation

$$(11) \quad h = \frac{2\pi^3 m c^2}{\omega g^2}.$$

The dimensional analysis gives $[J \cdot s] = [Kg][m/s]^2/[rad/s]$ which is correct. As the value of the constant angular speed ω is not known we express his value in function of k and the radius r_k

$$\omega = \frac{k\pi}{\tau} = \frac{k\pi c}{gr_k}.$$

To obtain the value of the constant g we substitute ω in (11) and evaluate at $n = 1$ (i.e. $k = 1$) and we obtain

$$(12) \quad h = \frac{2\pi^2 m c a_0}{g},$$

where a_0 is the Bohr radius at the fundamental level. Substituting into equation (12) the value of the reduced mass of the hydrogen atom m , the value of the speed of light c and the value of the Bohr radius of the fundamental level a_0 we obtain that the adimensional constant is $g = 429.868$. From equation (10) (using the quantification of Bohr, because we have identified it with the quantification that we have found) we obtain

$$(13) \quad v_\theta = \frac{nh}{2\pi m r_n},$$

and substituting into equation (9) we arrive to the Bohr radius

$$(14) \quad r_n = \frac{n^2 h^2 \epsilon}{\pi m e^2},$$

taking into account that $K = 1/(4\pi\epsilon)$. Finally, the energy levels are given by

$$E = \frac{mv_\theta^2}{2} - \frac{e^2}{4\pi\epsilon r_n} = -\frac{me^4}{8\epsilon^2 h^2 n^2}.$$

As a consequence, substituting into equation (12) the explicit expression of the Borh radius $a_0 = h^2\epsilon/(\pi e^2)$ we get the explicit value of the Planck constant

$$(15) \quad h = \frac{e^2 g}{2\pi \epsilon c}.$$

The introduction of a new fundamental constant is avoided (as it happens in the quantum mechanics with the Planck constant) because through the retard the speed of the light c appears. The appearance of this dimensional constant c , usually absent in the non-relativistic quantum mechanics, allows to give the expressions of the physical quantities with the correct dimensions.

I notice that this value of $g = 429.868$ is the inverse value of the fine structure constant $1/\alpha = 137.036$ multiplied by π , that is, $g = \pi/\alpha$. Moreover, the value of this fine structure constant α is

$$\alpha = \frac{e^2}{4\pi\epsilon\hbar c},$$

and when substituting this value of g in the explicit expression of the Planck constant (15) the equality is identically satisfied. Therefore we have found the value of the adimensional constant g and consequently the expression of the delay τ which is

$$(16) \quad \tau = \frac{\pi r}{\alpha c}.$$

Moreover, we have that $v_\theta/c = k\pi/g = k\alpha$ which is consequent with the definition of α . One of the interpretations of the fine structure constant α is that α relates the speed of the electron in the lowest energy level in the atom of hydrogen with the speed of the light. This is straightforward because if we substitute the expression (14) of r_n in the expression (13) of v_θ for the case $n = 1$ and we divide by the speed of the light c , we obtain that $v_\theta/c = \alpha$.

Summarizing, we could have begun our analysis with the founded retard definition (16), because our model reproduces the Borh atom faithfully. Quantum mechanics in the Borh atom is in fact the first approximation in the value v/c in the delay. It seems that with the second approximation in $(v/c)^2$ in the delay we will obtain the fine structure corrections of Sommerfeld [22] to the hydrogen atom.

The fact that the Planck constant is expressed in function of the parameters associated to the particular model system (see (15)) is a problem. If we consider another model system, usually another expression of the Planck constant would appear and another numerical value for the constant g would have to be chosen, and hence another expression for the delay.

From the fact that the atomic Bohr model can be completely described by means of functional differential equations, I believe that in the future, observing the physical reality at a deeper level, we will be able to interpret the laws of probability of the quantum physics as the statistical results of values of certain variables perfectly determined that at the moment are hidden for us...(I know that this idea it has been extremely discussed) the history of science shows us that the current state of the knowledge is always provisional and that it should exist, beyond what is known, immense regions to be discovered. However, functional differential equations are fundamentally different from ordinary differential equations, and their solutions can have qualitative features which are impossible for solutions of ordinary differential equations. The physical consequences of these differences are explained at length in [17]. I mean that the rich features that can appear in functional differential equations may explain quantum mechanics, from a deeper point of view.

Acknowledgements:

The author would like to thank Prof. H. Giacomini from Université de Tours for several useful conversations and remarks.

REFERENCES

- [1] N. BOHR, *On the Constitution of Atoms and Molecules*, Phil. Mag. **26** (1913), 1–25.
- [2] M. BORN, *Zur Quantenmechanik der Stoßvorgänge*, Z. Phys., **37** (1926), 863–867.
- [3] C. CHICONE, *What are the equations of motion of classical physics?*, Can. Appl. Math. Q. **10** (2002), no. 1, 15–32.
- [4] C. CHICONE, B. MASHHOON AND D. RETZLOFF, *Delay equations and radiation damping*, Phys. Letters A **285** (2000), 17–16.
- [5] C. CHICONE, *Inertial and slow manifolds for delay equations with small delays*, J. Differential Equations **190** (2003), no. 2, 364–406.
- [6] C. CHICONE, *Inertial flows, slow flows, and combinatorial identities for delay equations*, J. Dynam. Differential Equations **16** (2004), no. 3, 805–831.
- [7] P.A.M. DIRAC, *The quantum theory of the electron*, Proc. Roy. Soc. A **117** (1928), 610–624.
- [8] A. EINSTEIN, *Über einen die Erzeugung und Verwandlung des Lichts betreffenden heuristischen Gesichtspunkt*, Ann. Phys. **17** (1905), 132–148.
- [9] J. GINÉ, *On the classical descriptions of the quantum phenomena in the harmonic oscillator and in a charged particle under the coulomb force*, Chaos Solitons Fractals, to appear.
- [10] W. HEISENBERG, *Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik*, Z. Phys., **43** (1927), 172–198.
- [11] L.D. LANDAU AND E.M. LIFSHITZ, *The classical theory of fields*, Oxford: Pergamon Press, 1971.

- [12] M. PLANCK, *Über das Gesetz der Energieverteilung im Normalspectrum*, Ann. Phys. **4** (1901), 553-563.
- [13] H. POINCARÉ, *Mémoire sur les courbes définies par les équations différentielles*. Journal de Mathématiques **37** (1881), 375-422; **8** (1882), 251-296; Oeuvres de Henri Poincaré, vol. I, Gauthier-Villars, Paris, (1951), pp. 3-84.
- [14] H. POINCARÉ, *Sur la théorie des quanta*, C. R. Acad. Sci. Paris **153** (1911), 1103-1108.
- [15] H. POINCARÉ, *Sur la théorie des quanta*, J. Physique Théorique et Appliquée, 5 série **2** (1912), 5-34.
- [16] C.K. RAJU, *The electrodynamic 2-body problem and the origin of quantum mechanics*, Foundations of Physics **34** (2004), 937-962.
- [17] C.K. RAJU, *Time: towards a consistent theory*, Kluwer academic, Dordrecht, 1994.
- [18] E. RUTHERFORD, *The Scattering of α and β Particles by Matter and the Structure of the Atom*, Phil. Mag. **21** (1911) 669-668.
- [19] E. SCHRÖDINGER, *Quantisierung als Eigenwertproblem. (Erste Mitteilung.)*, Ann. Phys. (Leipzig) **79** (1926), 361-376.
- [20] E. SCHRÖDINGER, *Quantisierung als Eigenwertproblem. (Zweite Mitteilung.)*, Ann. Phys. (Leipzig) **79** (1926), 489-527.
- [21] E. SCHRÖDINGER, *Quantisierung als Eigenwertproblem. (Dritte Mitteilung.)*, Ann. Phys. (Leipzig) **80** (1926), 437-490.
- [22] A. SOMMERFELD, *Einführung in die Quantentheorie, Oscillator und Rotator*, Atombau und Spectrallinien, Chapter 2 §3, Friedr. Vieweg & Sohn, Braunschweig, 1924.
- [23] A. SOMMERFELD, *Die Bohrsche Theorie der Balmerserie*, Atombau und Spectrallinien, Chapter 2 §4, Friedr. Vieweg & Sohn, Braunschweig, 1924.

DEPARTAMENT DE MATEMÀTICA, UNIVERSITAT DE LLEIDA, AV. JAUME II,
69. 25001 LLEIDA, SPAIN
E-mail address: gine@eps.udl.es