

Derivation of Newton's law of gravitation based on a fluidic continuum model of vacuum and a sink flow model of particles

Xiao-Song Wang*

Institute of Mechanical and Power Engineering, Henan Polytechnic University, Jiaozuo, Henan Province, 454000, China

(Dated: August 22, 2018)

The main purpose of this paper is to seek a mechanical interpretation of gravitational phenomena. We suppose that the universe may be filled with a kind of fluid which may be called the $\Omega(0)$ substratum. Thus, the inverse-square law of gravitation is derived by methods of hydrodynamics based on a sink flow model of particles. The first feature of this theory of gravitation is that the gravitational interactions are transmitted by a kind of fluidic medium. The second feature is the time dependence of gravitational constant G and gravitational mass. Newton's law of gravitation is derived if we introduce an assumption that G and the masses of particles are changing so slowly that they can be treated as constants.

PACS numbers:

I. INTRODUCTION

The Newton's law of gravitation can be written as

$$\mathbf{F}_{21} = -G \frac{m_1 m_2}{r^2} \hat{\mathbf{r}}_{21}, \quad (1)$$

where m_1 and m_2 are the masses of two particles, r is the distance between the two particles, G is the gravitational constant, \mathbf{F}_{12} is the force exerted on the particle with mass m_2 by the particle with mass m_1 , $\hat{\mathbf{r}}_{21}$ denotes the unit vector directed outward along the line from the particle with mass m_1 to the particle with mass m_2 .

The main purpose of this paper is to derive the Newton's law of gravitation by means of fluid mechanics based on sink flow model of particles.

The motive of this paper is to seek a mechanism of gravitational phenomena. The reasons why new models of gravity are interesting may be summarized as follows.

Firstly, there exists some astronomical phenomena that could not be interpreted by the present theories of gravitation, for instance, the Titius-Bode law [1]. New theories of gravity may view these problems from new angles.

Secondly, whether the gravitational constant G depends on time and space is still unknown [2–8]. It is known that the gravitational constant G is a constant in the Newton's theory of gravitation and in theory of general relativity.

Thirdly, the mechanism of the action-at-a-distance gravitation remains an unsolved problem in physics for more than 300 years [9–11]. Although theory of general relativity is a field theory of gravity[12], the concept of field is different from that of continuum mechanics [13–16] because of the absence of a continuum in theory of general relativity. Thus, theory of general relativity can only be regarded as a phenomenological theory of gravity.

Fourthly, we do not have a satisfactory quantum theory of gravity presently [17–21]. One of the challenges in theoretical physics is to reconcile quantum theory and theory of general relativity [17, 22]. New theories of gravity may open new ways to solve this problem.

Fifthly, one of the puzzles in physics is the problem of dark matter and dark energy [23–31]. New theories of gravity may provide new methods to attack this problem[24, 25].

Finally, we do not have a successful unified field theory presently. Great progress has been made towards an unification of the four fundamental interactions in the universe in the 20th century. However, gravitation is still not unified successfully. New theories of gravity may shed some light on this puzzle.

To conclude, it seems that new considerations on gravitation is needed. It is worthy keeping an open mind with respect to all the theories of gravity before the above problems been solved.

Now let us briefly review the long history of mechanical interpretations of gravitational phenomena. Many philosophers and scientists, such as Laozi[32], Thales, Anaximenes, believed that everything in the universe is made of a kind of fundamental substance[9]. Descartes was the first to bring the concept of aether into science by suggesting that it has mechanical properties[9]. Since the Newton's law of gravitation was published in 1687[33], this action-at-a-distance theory was criticized by the French Cartesian[9]. Newton admitted that his law did not touch on the mechanism of gravitation[34]. He tried to obtain a derivation of his law based on Descartes' scientific research program [33]. Newton himself even suggested an explanation of gravity based on the action of an aetherial medium pervading the space[34, 35]. Euler attempted to explain gravity based on some hypotheses of a fluidic aether[9].

In a remarkable paper published in 1905, Einstein abandoned the concept of aether[36]. However, Einstein's assertion did not cease the explorations of aether [9, 37–46]. Einstein changed his view later and introduced his new concept of ether[47, 48]. I regret to admit that it is

*Electronic address: wangxs2002@163.com

impossible for me to mention all the works related to this field in history. Adolphe Martin and Roy Keys[49–51] proposed a gas model of vacuum to explain the physical phenomena such as electromagnetism, gravitation, quantum mechanics and the structure of elementary particles.

Inspired by the aforementioned thoughts and others[52–56], we show that the Newton’s law of gravitation is derived based on the assumption that all the particles are made of singularities of a kind of ideal fluid.

During the preparation of the manuscript, I noticed that John C. Taylor had proposed an idea that the inverse-square law of gravitation may be explained based on the concept of source or sink [57].

II. FORCES ACTING ON SOURCES AND SINKS IN IDEAL FLUIDS

The purpose of this section is to calculate the forces between sources and sinks in inviscid incompressible fluids which is called ideal fluids usually.

Suppose the velocity field \mathbf{u} of an ideal fluid is irrotational, then we have [16, 54–56, 58–60],

$$\mathbf{u} = \nabla\phi, \quad (2)$$

where ϕ is the velocity potential, $\nabla = \mathbf{i}\frac{\partial}{\partial x} + \mathbf{j}\frac{\partial}{\partial y} + \mathbf{k}\frac{\partial}{\partial z}$ is the Hamilton operator.

It is known that the equation of mass conservation of an ideal fluid becomes Laplace’s equation [54–56, 58–60],

$$\nabla^2\phi = 0, \quad (3)$$

where ϕ is velocity potential, $\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ is the Laplace operator.

Using spherical coordinates(r, θ, φ), a general form of solution of Laplace’s equation (3) can be obtained by separation of variables as[56]

$$\phi(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta), \quad (4)$$

where A_l and B_l are arbitrary constants, $P_l(x)$ are Legendre’s function of the first kind which is defined as

$$P_l(x) = \frac{1}{2^l l!} \frac{d^l}{dx^l} (x^2 - 1)^l. \quad (5)$$

If there exists a velocity field which is continuous and finite at all points of the space, with the exception of individual isolated points, then these isolated points are called singularities usually.

Definition 1 Suppose there exists a singularity at point $P_0 = (x_0, y_0, z_0)$. If the velocity field of the singularity at point $P = (x, y, z)$ is

$$\mathbf{u}(x, y, z, t) = \frac{Q}{4\pi r^2} \hat{\mathbf{r}}, \quad (6)$$

where $r = \sqrt{(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2}$, $\hat{\mathbf{r}}$ denotes the unit vector directed outward along the line from the singularity to the point $P = (x, y, z)$, then we call this singularity a source if $Q > 0$ or a sink if $Q < 0$. Q is called the strength of the source or sink.

Suppose a static point source with strength Q locates at the origin $(0, 0, 0)$. In order to calculate the volume leaving the source per unit time, we may enclose the source with an arbitrary spherical surface S with radius a . A calculation shows that

$$\oint_S \mathbf{u} \cdot \mathbf{n} dS = \oint_S \frac{Q}{4\pi a^2} \hat{\mathbf{r}} \cdot \mathbf{n} dS = Q, \quad (7)$$

where \mathbf{n} denotes the unit vector directed outward along the line from the origin of the coordinates to the field point (x, y, z) . Equation (7) shows that the strength Q of a source or sink evaluates the volume of the fluid leaving or entering a control surface per unit time.

From (4), we see that the velocity potential $\phi(r, \theta)$ of a source or sink is a solution of Laplace’s equation (3).

Theorem 2 Suppose (1) there exists an ideal fluid (2) the ideal fluid is irrotational and barotropic, (3) the density ρ is homogeneous, that is $\partial\rho/\partial x = \partial\rho/\partial y = \partial\rho/\partial z = \partial\rho/\partial t = 0$, (4) there are no external body forces exerted on the fluid, (5) the fluid is unbounded and the velocity of the fluid at the infinity is approaching to zero. Suppose a source or sink is stationary and is immersed in the ideal fluid. Then, there is a force

$$\mathbf{F}_Q = -\rho Q \mathbf{u}_0 \quad (8)$$

exerted on the source by the fluid, where ρ is the density of the fluid, Q is the strength of the source or the sink, \mathbf{u}_0 is the velocity of the fluid at the location of the source induced by all means other than the source itself.

Proof. Only the proof of the case of a source is needed. Let us select the coordinates that is attached to the static fluid at the infinity.

We set the origin of the coordinates at the location of the source. We surround the source by two arbitrary spherical surfaces S_ε and S centered at the origin of the coordinates. The radii of the surfaces S_ε and S are r_ε and r respectively. The radius r_ε of the surfaces S_ε is smaller than r and is arbitrarily small. The outward unit normal to the surface S is denoted by \mathbf{n} .

Let $\tau(t)$ denotes the mass system of fluid enclosed in the volume between the surface S_ε and S at time t . Let \mathbf{F}_Q denotes the hydrodynamic force exerted on the source by the mass system τ , then a reaction $-\mathbf{F}_Q$ of this force \mathbf{F}_Q must act on the the surfaces S_ε enclosing the source. Let \mathbf{F}_S denotes the hydrodynamic force exerted on the mass system τ due to the pressure distribution on the surface S .

As an application of the Newton's second law of motion to the mass system τ , we have

$$\frac{D\mathbf{K}}{Dt} = -\mathbf{F}_Q + \mathbf{F}_S, \quad (9)$$

where \mathbf{K} denotes momentum of the mass system τ , D/Dt represents the material derivative in the lagrangian system [16, 54–56, 58–60].

The expressions of the momentum \mathbf{K} and the force \mathbf{F}_S are

$$\mathbf{K} = \iiint_{\tau} \rho \mathbf{u} dV, \quad \mathbf{F}_S = \iint_S (-p)\mathbf{n} dS, \quad (10)$$

where the first integral is volume integral, the second integral is surface integral, \mathbf{n} denotes the unit vector directed outward along the line from the origin of the coordinates to the field point (x, y, z) .

Since the velocity field is irrotational, we have the following relation

$$\mathbf{u} = \nabla\phi, \quad (11)$$

where ϕ is the velocity potential.

According to Ostrogradsky–Gauss theorem (see, for instance, [54–56, 59, 60]), we have

$$\iiint_{\tau} \rho \mathbf{u} dV = \iiint_{\tau} \rho \nabla\phi dV = \iint_S \rho \phi \mathbf{n} dS. \quad (12)$$

Note that the mass system τ does not include the singularity at the origin. According to Reynolds' transport theorem [54–56, 59, 60], we have

$$\frac{D}{Dt} \iiint_{\tau} \rho \mathbf{u} dV = \frac{\partial}{\partial t} \iiint_V \rho \mathbf{u} dV + \iint_S \rho \mathbf{u}(\mathbf{u} \cdot \mathbf{n}) dS, \quad (13)$$

where V is the volume fixed in space which coincide with the mass system $\tau(t)$ at time t , that is $V = \tau(t)$.

Then, using (13), (10) and (12), we have

$$\frac{D\mathbf{K}}{Dt} = \iint_S \rho \frac{\partial\phi}{\partial t} \mathbf{n} dS + \iint_S \rho \mathbf{u}(\mathbf{u} \cdot \mathbf{n}) dS. \quad (14)$$

According to Lagrange–Cauchy integral [54–56, 59, 60], we have

$$\frac{\partial\phi}{\partial t} + \frac{(\nabla\phi)^2}{2} + \frac{p}{\rho} = f(t), \quad (15)$$

where $f(t)$ is an arbitrary function of time t .

According to the assumption that the velocity \mathbf{u} of the fluid at the infinity is approaching to zero, we have

$$|\mathbf{u}| \rightarrow 0, \quad r \rightarrow \infty. \quad (16)$$

Noticing (4), $\phi(t)$ must be of the following form

$$\phi(r, \theta, t) = \sum_{l=0}^{\infty} \frac{B_l(t)}{r^{l+1}} P_l(\cos \theta), \quad (17)$$

where $B_l(t), l \geq 0$ are functions of time t . Thus, we have the following estimations at the infinity of the velocity field

$$\phi = O\left(\frac{1}{r}\right), \quad \frac{\partial\phi}{\partial t} = O\left(\frac{1}{r}\right), \quad r \rightarrow \infty, \quad (18)$$

where $\varphi(x) = O(\psi(x)), x \rightarrow a$ stands for $\overline{\lim}_{x \rightarrow a} |\varphi(x)| / \psi(x) = k, (0 \leq k < +\infty)$.

Applying (15) at the infinity and using (18), we have

$$\frac{\partial\phi}{\partial t} \rightarrow 0, \quad p \rightarrow p_{\infty}, \quad r \rightarrow \infty, \quad (19)$$

where p_{∞} is a constant.

Putting (19) and (16) into (15), we have

$$f(t) = \frac{p_{\infty}}{\rho}. \quad (20)$$

Putting (20) into (15), we have

$$p = p_{\infty} - \rho \frac{\partial\phi}{\partial t} - \frac{\rho(\mathbf{u} \cdot \mathbf{u})}{2}. \quad (21)$$

Using (10) and (21), we have

$$\mathbf{F}_S = \iint_S \rho \frac{\partial\phi}{\partial t} \mathbf{n} dS + \iint_S \frac{\rho(\mathbf{u} \cdot \mathbf{u}) \mathbf{n}}{2} dS. \quad (22)$$

Using (9), (14), (22), we have

$$\mathbf{F}_Q = \iint_S \left[\frac{1}{2} \rho(\mathbf{u} \cdot \mathbf{u}) \mathbf{n} - \rho \mathbf{u}(\mathbf{u} \cdot \mathbf{n}) \right] dS. \quad (23)$$

Now let us calculate this velocity \mathbf{u} in order to obtain \mathbf{F}_Q . Since the velocity field induced by the source Q is (6), then according to the superposition principle of velocity field of ideal fluids, the velocity on the surface S is

$$\mathbf{u} = \frac{Q}{4\pi r^2} \mathbf{n} + \mathbf{u}_0, \quad (24)$$

where \mathbf{n} denotes the unit vector directed outward along the line from the origin of the coordinates to the field point (x, y, z) . Using (23) and (24), we have

$$\mathbf{F}_Q = \rho \iint_S \left[\frac{Q^2}{32\pi^2 r^4} \mathbf{n} + \frac{1}{2}(\mathbf{u}_0 \cdot \mathbf{u}_0) \mathbf{n} - \frac{Q}{4\pi r^2} \mathbf{u}_0 - (\mathbf{u}_0 \cdot \mathbf{n}) \mathbf{u}_0 \right] dS. \quad (25)$$

Since the radius r can be arbitrarily small, the velocity \mathbf{u}_0 can be treated as a constant in the integral of (25). Thus, (25) turns out to be

$$\mathbf{F}_Q = -\rho \iint_S \frac{Q}{4\pi r^2} \mathbf{u}_0 dS. \quad (26)$$

Since again \mathbf{u}_0 can be treated as a constant, (26) turns out to be (8). This completes the proof. \square

Remark. Lagally [52], Landweber and Yih [53, 54], Faber [55] and Currie [56] obtained the same result of Theorem 2 for the special case where the velocity field is steady.

Theorem 2 only considers the situation that the sources or sinks are at rest. Now let us consider the case that the sources or sinks are moving in the fluid.

Theorem 3 *Suppose the presuppositions (1), (2), (3), (4) and (5) in Theorem 2 are valid and a source or a sink is moving in the fluid with a velocity \mathbf{v}_s , then there is a force*

$$\mathbf{F}_Q = -\rho Q (\mathbf{u}_f - \mathbf{v}_s) \quad (27)$$

is exerted on the source by the fluid, where ρ is the density of the fluid, Q is the strength of the source or the sink, \mathbf{u}_f is the velocity of the fluid at the location of the source induced by all means other than the source itself.

Proof. The velocity of the fluid relative to the source at the location of the source is $\mathbf{u}_f - \mathbf{v}_s$. Let us select the coordinates that is attached to the source and set the origin of the coordinates at the location of the source. Then (27) can be arrived following the same procedures in the proof of Theorem 2. \square

Applying Theorem 3 to the situation that a source or sink is exposed to the velocity field of another source or sink, we have:

Corollary 4 *Suppose the presuppositions (1), (2), (3), (4) and (5) in Theorem 2 are valid and a source or a sink with strength Q_2 is exposed to the velocity field of another source or sink with strength Q_1 , then the force \mathbf{F}_{21} exerted on the singularity with strength Q_2 by the velocity field of the singularity with strength Q_1 is*

$$\mathbf{F}_{21} = -\rho Q_2 \frac{Q_1}{4\pi r^2} \hat{\mathbf{r}}_{21} + \rho Q_2 \mathbf{v}_2, \quad (28)$$

where $\hat{\mathbf{r}}_{21}$ denotes the unit vector directed outward along the line from the singularity with strength Q_1 to the singularity with strength Q_2 , r is the distance between the two singularities, \mathbf{v}_2 is the velocity of the source with strength Q_2 .

III. DERIVATION OF INVERSE-SQUARE-LAW OF GRAVITATION

Since quantum theory shows that vacuum is not empty and has physical effects, e.g., the Casimir effect[45, 61–63], it is valuable to probe vacuum by introducing the following hypotheses:

Assumption 5 *Suppose the universe is filled by an ideal fluid named $\Omega(0)$ substratum; the ideal fluid fulfil the conditions (2), (3), (4), (5) in Theorem 2.*

This fluid may be named $\Omega(0)$ substratum in order to distinguish with Cartesian aether.

The idea that all microscopic particles are source or sink flows in a fluidic substratum is not new. For instance, in order to compare fluid motions with electric fields, J. C. Maxwell introduced an analogy between source or sink flows and electric charges ([64], p243). H. Poincaré also speculates that matters are holes in fluidic aether ([65], p171). A. Einstein and L. Infeld said ([66], p256-257):”Matter is where the concentration of energy is great, field where the concentration of energy is small. . . . What impresses our senses as matter is really a great concentration of energy into a comparatively small space. We could regard matter as the regions in space where the field is extremely strong.” It seems that they are suggesting that particles are some kinds of singularities of field.

Following these researchers, we adopt the idea that particles may be looked as singularities in fields. Noticing (28), it is nature to introduce the following:

Assumption 6 *All the microscopic particles were made up of a kind of elementary sinks of $\Omega(0)$ substratum. These elementary sinks were created simultaneously. The initial masses and the strengths of the elementary sinks are the same.*

We may call these elementary sinks as monads. Suppose a particle with mass m is composed of N monads. Then, according to Assumption 6, we have:

$$m_0(t) = m_0(0) + \rho q_0 t, \quad (29)$$

$$Q = -Nq_0, \quad m(t) = Nm_0(t) = -\frac{Q}{q_0} m_0(t), \quad (30)$$

$$\frac{dm_0}{dt} = \rho q_0, \quad \frac{dm}{dt} = -\rho Q, \quad (31)$$

where $m_0(t)$ is the mass of monad at time t , $-q_0$ ($q_0 > 0$) is the strength of a monad, $m(t)$ is the mass of a particle at time t , Q is the strength of the particle, N is the number of monads that make up the particle, ρ is the density of the $\Omega(0)$ substratum, $t \geq 0$.

From (31), we see that the mass m_0 of a monad is increasing since q_0 evaluates the volume of the $\Omega(0)$ substratum fluid entering the monad per unit time. From (31), we also see that the mass of a monad or a particle is increasing linearly.

Based on Assumption 5 and Assumption 6, the motion of a particle is determined by:

Theorem 7 *The equation of motion of a particle is*

$$m(t) \frac{d\mathbf{v}}{dt} = \frac{\rho q_0}{m_0(t)} m(t) \mathbf{u} - \frac{\rho q_0}{m_0(t)} m(t) \mathbf{v} + \mathbf{F}, \quad (32)$$

where $m_0(t)$ is the mass of monad at time t , $-q_0$ is the strength of a monad, $m(t)$ is the mass of a particle at time t , \mathbf{v} is the velocity of the particle, \mathbf{u} is the velocity of the $\Omega(0)$ substratum at the location of the particle induced by all means other than the particle itself, \mathbf{F} denotes other forces.

Proof. Applying the Newton's second law and Theorem 3 to this particle, we have

$$m \frac{d\mathbf{v}}{dt} = -\rho Q(\mathbf{u} - \mathbf{v}) + \mathbf{F}. \quad (33)$$

Putting (30) into (33), we get (32). \square

Formula (32) shows that there exists a universal damping force

$$\mathbf{F}_d = -\frac{\rho q_0}{m_0} m \mathbf{v} \quad (34)$$

exerted on each particle.

Now let us consider a system consists of two particles. Based on Assumption 6, applying Theorem 7 to this system, we have:

Corollary 8 *Suppose there is a system consists of two particles and there are no other forces exerted on the particles, then the equations of motion of this system are*

$$m_1 \frac{d\mathbf{v}_1}{dt} = -\frac{\rho q_0}{m_0} m_1 \mathbf{v}_1 - \frac{\rho q_0^2}{4\pi m_0^2} \frac{m_1 m_2}{r^2} \hat{\mathbf{r}}_{12} \quad (35)$$

$$m_2 \frac{d\mathbf{v}_2}{dt} = -\frac{\rho q_0}{m_0} m_2 \mathbf{v}_2 - \frac{\rho q_0^2}{4\pi m_0^2} \frac{m_1 m_2}{r^2} \hat{\mathbf{r}}_{21}, \quad (36)$$

where $m_{i=1,2}$ is the mass of the particles, $\mathbf{v}_{i=1,2}$ is the velocity of the particles, m_0 is the mass of a monad, $-q_0$ is the strength of a monad, ρ is the density of the $\Omega(0)$ substratum, $\hat{\mathbf{r}}_{12}$ denotes the unit vector directed outward along the line from the particle with mass $m_2(t)$ to the particle with mass $m_1(t)$, $\hat{\mathbf{r}}_{21}$ denotes the unit vector directed outward along the line from the particle with mass $m_1(t)$ to the particle with mass $m_2(t)$.

Ignoring the damping forces in (36), we have:

Corollary 9 *Suppose (1) $\mathbf{v}_{i=1,2} \ll \mathbf{u}_{i=1,2}$, where \mathbf{v}_i is the velocity of the particle with mass m_i , \mathbf{u}_i is the velocity of the $\Omega(0)$ substratum at the location of the particle with mass m_i induced by the other particle, (2) there are no other forces exerted on the particles, then the force $\mathbf{F}_{21}(t)$ exerted on the particle with mass $m_2(t)$ by the velocity field of $\Omega(0)$ substratum induced by the particle with mass $m_1(t)$ is*

$$\mathbf{F}_{21}(t) = -G(t) \frac{m_1(t)m_2(t)}{r^2} \hat{\mathbf{r}}_{21}, \quad (37)$$

where

$$G(t) = \frac{\rho q_0^2}{4\pi m_0^2(t)}, \quad (38)$$

$\hat{\mathbf{r}}_{21}$ denotes the unit vector directed outward along the line from the particle with mass $m_1(t)$ to the particle with mass $m_2(t)$, r is the distance between the two particles.

Corollary 9 is similar to Newton's inverse-square-law of gravitation (1) except for two differences. The first difference is that $m_{i=1,2}$ are constants in the Newton's

law (1) while in Corollary 9 they are functions of time t . The second difference is that G is a constant in the Newton's theory while here G is a function of time t .

Let us now introduce an assumption that G and the masses of particles are changing so slowly relative to the time scale of human beings that they can be treated as constants approximately. Thus, the Newton's law of gravitation may be considered as a result of Corollary 9 based on this assumption.

IV. SUPERPOSITION PRINCIPLE OF GRAVITATIONAL FIELD

The definition of gravitational field \mathbf{g} of a particle with mass m is

$$\mathbf{g} = \frac{\mathbf{F}}{m_{test}}, \quad (39)$$

where m_{test} is the mass of a test point mass, \mathbf{F} is the gravitational force exerted on the test point mass by the gravitational field of the particle with mass m .

Based on (39), Theorem 7 and Corollary 9, we have

$$\mathbf{g} = \frac{\rho q_0}{m_0} \mathbf{u}, \quad (40)$$

where ρ is the density of the $\Omega(0)$ substratum, m_0 is the mass of a monad, q_0 is the strength of a monad, \mathbf{u} is the velocity of the $\Omega(0)$ substratum at the location of the test point mass induced by the particle mass m .

From (40), we see that the superposition principle of gravitational field is deduced from the superposition theorem of the velocity field of ideal fluids.

V. EQUATIONS OF GRAVITATIONAL FIELD OF CONTINUOUSLY DISTRIBUTED PARTICLES

The definition of the volume density of continuously distributed sink is

$$\rho_s = \lim_{\Delta V \rightarrow 0} \frac{\Delta Q}{\Delta V}, \quad (41)$$

where ΔV is a small volume, ΔQ is the sum of strengthes of all the sinks in the volume ΔV .

Now let us to derive the continuity equation of the taothe $\Omega(0)$ substratum from the principle of mass conservation. Consider an arbitrary volume V bounded by a closed surface S and fixed in space. Suppose there are some point sources or sinks continuously distributed in the volume V . The total mass in volume V is

$$M = \iiint_V \rho dV, \quad (42)$$

where ρ is the density of the $\Omega(0)$ substratum.

The rate of increase of the total mass in volume V is

$$\frac{\partial M}{\partial t} = \frac{\partial}{\partial t} \iiint_V \rho dV. \quad (43)$$

The rate of mass outflow through the surface S is

$$\oiint_S \rho(\mathbf{u} \cdot \mathbf{n}) dS, \quad (44)$$

where \mathbf{u} is the velocity field of the $\Omega(0)$ substratum.

The rate of mass created inside the volume V is

$$\iiint_V \rho \rho_s dV. \quad (45)$$

Now according to the principle of mass conservation, and making use of (43), (44) and (45), we have

$$\frac{\partial}{\partial t} \iiint_V \rho dV = \iiint_V \rho \rho_s dV - \oiint_S \rho(\mathbf{u} \cdot \mathbf{n}) dS \quad (46)$$

According to Ostrogradsky–Gauss theorem (refer to, for instance, [54–56, 59, 60]), we have

$$\oiint_S \rho(\mathbf{u} \cdot \mathbf{n}) dS = \iiint_V \nabla \cdot (\rho \mathbf{u}) dV, \quad (47)$$

Using (47), (46) becomes

$$\frac{\partial}{\partial t} \iiint_V \rho dV = \iiint_V \rho \rho_s dV - \iiint_V \nabla \cdot (\rho \mathbf{u}) dV \quad (48)$$

Since the volume V is arbitrary, from (48) we have

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = \rho \rho_s. \quad (49)$$

Since the $\Omega(0)$ substratum is homogeneous, i.e., $\partial \rho / \partial x = \partial \rho / \partial y = \partial \rho / \partial z = \partial \rho / \partial t = 0$, (49) becomes

$$\nabla \cdot \mathbf{u} = \rho_s. \quad (50)$$

Thus, from (40) and (50), we have

$$\nabla \cdot \mathbf{G} = \frac{\rho q_0 \rho_s}{m_0}. \quad (51)$$

Let ρ_m denotes the volume mass density of continuously distributed particles. According to Assumption 6, and using (30) and (41), we have

$$\rho_m = -\frac{m_0 \rho_s}{q_0}. \quad (52)$$

Thus, noticing that the velocity field \mathbf{u} of the $\Omega(0)$ substratum is irrotational and $G = \rho q_0^2 / (4\pi m_0^2)$, the equations of gravitational field of continuously distributed particles can be summarized as

$$\begin{cases} \nabla \times \mathbf{G} = \mathbf{0}, \\ \nabla \cdot \mathbf{G} = -4\pi G \rho_m. \end{cases} \quad (53)$$

VI. THE EQUIVALENCE OF INERTIAL MASS AND GRAVITATIONAL MASS

According to Assumption 6 and Corollary 9, we have

$$m_{inertial} = m_{gravitational}, \quad (54)$$

where $m_{inertial}$ is the inertial mass of a particle, $m_{gravitational}$ is the gravitational mass of the particle.

VII. TIME DEPENDENCE OF GRAVITATIONAL CONSTANT G AND MASS

The time dependence of the gravitational constant G can be seen from (38).

The time dependence of the gravitational mass can be seen from (31).

VIII. POSSIBLE SPACE DEPENDENCE OF GRAVITATIONAL CONSTANT G

If the density ρ of the $\Omega(0)$ substratum varies from place to place, i.e., $\rho = \rho(\mathbf{r})$, then the space dependence of the gravitational constant G can be seen from (38).

IX. DISCUSSION

Although the new formula of gravitation (37) is similar to Newton's inverse-square-law of gravitation (1), there exists the following five differences between this theory and Newton's theory.

1. The gravitational masses are constants in Newton's law, while in (37) they are functions of time t .
2. The gravitational constant G is a constant in Newton's theory, while in (38) G depends on time t .
3. In this theory, the parameter G depends on the density $\rho_{\Omega(0)}$ of the $\Omega(0)$ substratum. If $\rho_{\Omega(0)}$ varies from place to place, then the space dependence of the gravitational constant G can be seen from (38).
4. In Newton's theory, the gravity is action-at-a-distance [9]. In this theory, the gravity is transmitted by the $\Omega(0)$ substratum.
5. Newton's law of gravitation is an assumption. In this theory, (37) is derived by methods of classical fluid mechanics based on some assumptions.

X. CONCLUSION

We suppose that the universe may be filled with a kind of fluid which may be called the $\Omega(0)$ substratum. Thus, the inverse-square law of gravitation is derived by methods of hydrodynamics based on a sink flow model of particles. There are two features of this theory of gravitation. The first feature is that the gravitational interactions are

transmitted by a kind of fluidic medium. The second feature is the time dependence of gravitational constant and gravitational mass. Newton's law of gravitation is derived if we introduce an assumption that G and the masses of particles are changing so slowly that they can be treated as constants. As a byproduct, it is shown that there exists a universal damping force exerted on each particle.

Acknowledgments

I am grateful to Robert L. Oldershaw for informing me his researches in the field of Self-Similar Cosmolog-

ical Paradigm (SSCP) and discrete fractal cosmological models during the preparation of the manuscript. I wish to express my thanks to Dr. Roy Keys for providing me three interesting articles [49–51].

-
- [1] M. M. Nietro, *The Titius-Bode Law of planetary distances: its history and theory* (Pergamon Press, New York, 1972).
- [2] J. J. Gilvarry and P. M. Muller, Phys. Rev. Lett. **28**, 1665 (1972).
- [3] R. Tomaschitz, Astrophysics and Space Science **271**, 181 (2000).
- [4] E. Gaztanaga, E. Garcia-Berro, J. Isern, E. Bravo, and I. Dominguez, Phys. Rev. D **65**, 023506 (2001).
- [5] C. J. Copi, A. N. Davis, and L. M. Krauss, Phys. Rev. Lett. **92**, 171301 (2004).
- [6] J. Khoury and A. Weltman, Phys. Rev. Lett. **93**, 171104 (2004).
- [7] R. Nagata, T. Chiba, and N. Sugiyama, Phys. Rev. D **69**, 083512 (2004).
- [8] M. Biesiada and B. Malec, Mon. Not. R. Astron. Soc. **350**, 644 (2004).
- [9] E. Whittaker, *A History of the Theories of Aether and Electricity* (Thomas Nelson and Sons Ltd., London, 1953).
- [10] M. B. Hesse, *Forces and Fields: the Concept of Action at a Distance in the History of Physics* (Thomas Nelson and Sons Ltd, London, 1961).
- [11] T. Jaakkola, Apeiron **3**, 61 (1996).
- [12] V. Fock, *The Theory of Space, Time and Gravitation, translated by N. Kemmer* (Pergamon Press, London, 1959).
- [13] C. Truesdell, *The Elements of Continuum Mechanics* (Springer-Verlag, New York, 1966).
- [14] Y. C. Fung, *A First Course in Continuum Mechanics* (Prentice-Hall, London, 1977).
- [15] A. C. Eringen, *The Elements of Continuum Mechanics* (Robert E. Krieger Pub. Co., Huntington, 1980).
- [16] L. D. Landau and Lifshitz, *Fluid Mechanics, translated from the Russian by J.B. Sykes and W.H. Reid*. (Pergamon, New York, 1987).
- [17] S. Carlip, Rep. Prog. Phys. **64**, 885 (2001).
- [18] G. Amelino-Camelia, Modern Physics Letters A **17**, 899 (2002).
- [19] D. V. Ahluwalia, Modern Physics Letters A **17**, 1135 (2002).
- [20] I. G. Mitrofanov, Nature **426**, 139 (2003).
- [21] J. Christian, Phys. Rev. D **71**, 024012 (2005).
- [22] J. Collins, A. Perez, D. Sudarsky, L. Urrutia, and H. Vucetich, Phys. Rev. Lett. **93**, 191301 (2004).
- [23] J. Ellis, Philos. Trans. R. Soc. London, A **361**, 2607 (2003).
- [24] E. V. Linder, Phys. Rev. D **70**, 023511 (2004).
- [25] F. Bernardeau, Rep. Prog. Phys. **66**, 691 (2003).
- [26] L. Bergstrom, T. Bringmann, M. Eriksson, and et al., Phys. Rev. Lett. **94**, 131301 (2005).
- [27] J. F. Beacom, N. F. Bell, and G. Bertone, Phys. Rev. Lett. **94**, 171301 (2005).
- [28] D. S. Akerib, M. J. Attisha, C. N. Bailey, and et al., Phys. Rev. Lett. **96**, 011302 (2006).
- [29] J. L. Feng, S. F. Su, and F. Takayama, Phys. Rev. Lett. **96**, 151802 (2006).
- [30] G. S. P. Fayet, D. Hooper, Phys. Rev. Lett. **96**, 211302 (2006).
- [31] M. Carena, D. Hooper, and P. Skands, Phys. Rev. Lett. **97**, 051801 (2006).
- [32] Laozi, *Tao Te King : in seven languages* (Farkas Lorinc Imre Pub., transl. Stephen Mitchell et al., 1995).
- [33] I. Newton, *Mathematical Principles of Natural Philosophy and His System of the World* (Univ. of Calif. Press, Berkeley, 1962).
- [34] I. B. Cohen, *The Newtonian Revolution* (Cambridge University Press, 1980).
- [35] I. Newton, *Opticks* (Bell London, 1931).
- [36] A. Einstein, Ann. Phys. **17**, 891 (1905).
- [37] J. P. Vigièr, Lettere al Nuovo Cimento **29**, 467 (1980).
- [38] A. Barut, Found. Phys. **18**, 95 (1988).
- [39] R. L. Oldershaw, International Journal of Theoretical Physics **28**, 669 (1989).
- [40] R. L. Oldershaw, International Journal of Theoretical Physics **28**, 1503 (1989).
- [41] M. Carvalho and et al., Found. of Phys. Lett. **16**, 255 (2003).
- [42] M. Arminjon, Int. J. of Modern Phys. A **17**, 4203 (2002).
- [43] M. Carvalho and A. L. Oliveira, Found. Phys. Lett. **16**, 255 (2003).
- [44] T. Jacobson, Phys. Rev. D **70**, 024003 (2004).
- [45] P. C. W. Davies, J. Opt. B: Quantum Semiclass. Opt. **7**, S40 (2005).
- [46] M. Levin and X.-G. Wen, Phys. Rev. B **73**, 035122 (2006).
- [47] A. Einstein, *Aether and the Theory of Relativity, 1920, translated in 'Sidelights on Relativity', Dover* (1983).

- [48] L. Kostro, *Einstein and the Ether* (Apeiron, Montreal, 2000).
- [49] A. Martin, *Gravitation in a gaseous ether. In: Einstein and Poincaré: the physical vacuum, Valeri V. Dvoeglazov ed.* (Apeiron, Montreal, 2005).
- [50] A. Martin, *The electron as an extended structure in cosmic gas. In: What is the Electron? Volodimir Simulik ed.* (Apeiron, Montreal, 2005).
- [51] A. Martin and C. Keys, *The ether revisited. In: Frontiers of Fundamental Physics, Volodimir Simulik ed.* (Plenum, New York, 1994).
- [52] M. Lagally, *Z. Angew. Math. Mech.* **2**, 409 (1922).
- [53] L. Landweber and C.-S. Yih, *J. Fluid Mech.* **1**, 319 (1956).
- [54] C.-S. Yih, *Fluid Mechanics* (McGraw-Hill Book Company, New York, 1969).
- [55] T. E. Faber, *Fluid Dynamics for Physicists* (Cambridge University Press, Cambridge, 1995).
- [56] I. G. Currie, *Fundamental Mechanics of Fluids* (Cambridge University Press, 2003), 3rd ed.
- [57] J. Taylor, *Hidden unity in Nature's laws. Appendix A* (Cambridge University Press, 2001).
- [58] H. Lamb, *Hydrodynamics* (Cambridge University Press, 1932), 6th ed.
- [59] N. E. Kochin, I. A. Kibel, and N. V. Roze., *Theoretical hydrodynamics, Translated from the fifth Russian ed.* (Interscience Publishers, New York, 1964).
- [60] W.-Y. Wu, *Fluid Mechanics, vol. 1, in Chinese* (Beijing University Press, Beijing, 1982).
- [61] S. K. Lamoreaux, *Rep. Prog. Phys.* **68**, 201 (2005).
- [62] F. Intravaia and A. Lambrecht, *Phys. Rev. Lett.* **94**, 110404 (2005).
- [63] J. G. Guo and Y. P. Zhao, *Journal of Microelectromechanical Systems* **13**, 1027 (2004).
- [64] E. Whittaker, *A History of the Theories of Aether and Electricity, Revised and enlarged edition, vol. 1* (Thomas Nelson and Sons Ltd., London, 1951).
- [65] H. Poincare, *The science and hypothesis, translated from the French edition, in Chinese* (Shangwu Press, Beijing, 1997).
- [66] A. Einstein and L. Infeld, *The evolution of physics* (Cambridge University Press, London, 1938).