

Influence of Gravitation on Mass-Energy Equivalence Relation

R. V. R. Pandya*,
Department of Mechanical Engineering,
University of Puerto Rico at Mayaguez, Mayaguez, PR 00680, USA

November 20, 2018

Abstract

We study influence of gravitational field on the mass-energy equivalence relation by incorporating gravitation in the physical situation considered by Einstein (Ann. Physik, 17, 1905, English translation in ref. [1]) for his first derivation of mass-energy equivalence. In doing so, we also refine Einstein's expression (Ann. Physik, 35, 1911, English translation in ref. [3]) for increase in gravitational mass of the body when it absorbs E amount of radiation energy.

1 Introduction

Different forms of mass-energy equivalence relation existed even before Einstein's first derivation of the relation [1] and which have been reviewed along with other developments on the relation after the year 1905 (see Ref. [2] and references cited therein). These relations were obtained in the absence of gravitational field. In this paper, we show that the presence of gravitational field affects the mass-energy equivalence relation. In doing so, we include gravitational field in the physical situation considered by Einstein [1] and derive the general mass-energy equivalence relation which reduces to the well known Einstein's relation in the absence of gravitation. We also suggest refinement to the expression, provided by Einstein [3], for increase in gravitational mass due to the absorption of radiation energy E .

2 Analysis

Consider a 'stationary' reference frame S_s with coordinate axes x, y, z and another moving reference frame S_v having coordinate axes x', y', z' which are par-

*Email address: rvrpturb@uprm.edu

allel to axes x, y, z , respectively. Let S_v is moving along the x axis with a constant translational velocity v as measured in S_s . At time $t = 0$ measured in S_s , let the origins of S_v and S_s coincide. Also, consider these two reference frames in the homogeneous gravitational field whose potential is Φ_s and Φ_v as measured in S_s and S_v , respectively. For the homogeneous gravitational field with acceleration of gravity γ and lines of force of the gravitational field in the negative direction of the axes z and z' , we have $\Phi_s = \Phi_v = \gamma h$. Here h is distance along the z and z' axis from the origins of S_s and S_v , respectively.

Consider a body W of mass M_s kept at rest in S_s on the z axis at a distance h from the origin, i.e having coordinates $(0, 0, h)$. At some instance it emits in two opposite directions equal quantity of light having energy $L/2$ where M_s and L are measured in S_s . The conservation of energy principle for this situation in S_s can be written as

$$E_0 = E_1 + L \frac{\left(1 + \frac{\Phi_s}{c^2}\right)}{\sqrt{1 - \Phi_s^2/c^4}} \quad (1)$$

where E_0 and E_1 are, respectively, total energy of the body before and after the emission of the light as measured in S_s and c is speed of light in S_s . It should be noted that c depends on the gravitational field, written as [3]

$$c \cong c_0 \left(1 + \frac{\Phi_s}{c^2}\right) \quad (2)$$

where c_0 is speed of light in the absence of gravitational field, i.e. when $\Phi_s = 0$. The factor multiplying L in the last term takes into account gravitation of the emitted energy L [3] and which reduces to 1 in the absence of gravitational field i.e. $\Phi_s = 0$.

It should be noted that the last term can be approximated as

$$L \frac{\left(1 + \frac{\Phi_s}{c^2}\right)}{\sqrt{1 - \Phi_s^2/c^4}} \cong L + L \frac{\Phi_s}{c^2} \quad (3)$$

when $c^2 \gg \Phi_s$. Einstein [3] considered this approximate expression $L + L \frac{\Phi_s}{c^2}$, representing summation of emitted energy L and gravitation of energy $L \frac{\Phi_s}{c^2}$, for his derivation of gravitation of energy and showed increase in gravitational mass equal to increase in inertia mass for any body when the body absorbs radiation energy. Here we should mention that if, instead of approximate expression of

type $L + L \frac{\Phi_s}{c^2}$, we consider more accurate expression of type $L \frac{\left(1 + \frac{\Phi_s}{c^2}\right)}{\sqrt{1 - \Phi_s^2/c^4}}$ in the Einstein's analysis [3] on increase in gravitational mass of the body when it absorbs E amount of radiation energy, we would obtain this increase in mass equal to

$$\frac{E}{\Phi_s} \left[\frac{1 + \frac{\Phi_s}{c^2}}{\sqrt{1 - \Phi_s^2/c^4}} - 1 \right] \quad (4)$$

instead of E/c^2 as suggested by Einstein [3].

Now, the conservation of energy principle for the body W as observed from S_v can be written as

$$H_0 = H_1 + \frac{L}{\sqrt{1 - v^2/c^2}} \frac{\left(1 + \frac{\Phi_v}{c^2}\right)}{\sqrt{1 - \Phi_v^2/c^4}} \quad (5)$$

where H_0 and H_1 are, respectively, total energy of the body before and after the emission of the light as measured in S_v . While writing Eq. (5), we have used expression

$$\frac{L}{\sqrt{1 - v^2/c^2}} \quad (6)$$

for emitted energy as measured in S_v . Subtracting Eq. (1) from Eq. (5) and using $\Phi_v = \Phi_s$ yield

$$(H_0 - E_0) - (H_1 - E_1) = L \left[\frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right] \frac{\left(1 + \frac{\Phi_s}{c^2}\right)}{\sqrt{1 - \Phi_s^2/c^4}}. \quad (7)$$

Now, the total energy of the body is summation of potential energy P , kinetic energy K and energy related to internal state of the body which we refer here as internal energy I . We now write total energies (E_0, E_1, H_0 and H_1) before and after the emission in reference frames S_s and S_v in terms of potential, kinetic and internal energies.

In S_s ,

$$E_0 = M_s \Phi_s + M_s I_s \quad (8)$$

$$E_1 = (M_s - m_s) \Phi_s + (M_s - m_s) I'_s. \quad (9)$$

Here M_s is mass of the stationary body W before the emission, m_s is decrease in mass of the body due to the emission, I_s and I'_s are internal energy per unit mass of the body before and after the emission, respectively, and all are measured in S_s in which the body is stationary all the time.

As measured in S_v , we denote the mass of the moving body W before the emission by M_v , rest mass of the body before the emission when at rest in S_v by M_v^s , decrease in rest mass due to the emission by m_v^s , potential energy per unit mass by Φ_v , internal energy per unit mass of the body before and after the emission by I_v and I'_v , respectively. With these notations we can write total energy of the body before and after the emission as measured in S_v as

$$H_0 = K_0 + M_v^s \Phi_v + M_v^s I_v, \quad (10)$$

$$H_1 = K_1 + (M_v^s - m_v^s) \Phi_v + (M_v^s - m_v^s) I'_v, \quad (11)$$

where K_0 and K_1 represent kinetic energies before and after the emission, respectively. Substituting Eqs. (8)-(11) into Eq. (7), we obtain

$$K_0 - K_1 = L \left[\frac{1}{\sqrt{1 - v^2/c^2}} - 1 \right] \frac{\left(1 + \frac{\Phi_s}{c^2}\right)}{\sqrt{1 - \Phi_s^2/c^4}} \quad (12)$$

where we have used the fact that $M_s = M_v^s, m_s = m_v^s, \Phi_v = \Phi_s$ and internal energy per unit mass of the body is same in S_s and S_v , i.e. $I_v = I_s, I'_v = I'_s$. Further incorporation of expressions for kinetic energies K_0 and K_1 [4] yield

$$m_s c^2 = L \frac{\left(1 + \frac{\Phi_s}{c^2}\right)}{\sqrt{1 - \Phi_s^2/c^4}}, \quad (13)$$

exhibiting the effect of gravitation (through Φ_s) on the decrease in mass due to the emission of energy L . When $\Phi_s = 0$, this final Eq. (13) reduces to the well known mass-energy equivalence relation $L = m_s c_0^2$ derived by Einstein [1].

References

- [1] A. Einstein, in *The Principle of Relativity*, edited by W. Perrett and G. B. Jeffery (Dover Publications, Inc., New York, 1952), p. 69-71.
- [2] W. L. Fadner, *Am. J. Phys.* **56**, 114 (1988); J. Stachel and R. Torretti, *Am. J. Phys.* **50**, 760 (1982); *The EINSTEIN Myth and the IVES papers*, edited by D. Turner and R. Hazelett (The Devin-Adair Company, 1979)
- [3] A. Einstein, in *The Principle of Relativity*, edited by W. Perrett and G. B. Jeffery (Dover Publications, Inc., New York, 1952), p. 99-108.
- [4] A. Einstein, in *The Principle of Relativity*, edited by W. Perrett and G. B. Jeffery (Dover Publications, Inc., New York, 1952), p. 37-65.