

Stiff spinning torus of electromagnetic two-body motion

Jayme De Luca*

Universidade Federal de São Carlos,

Departamento de Física

Rodovia Washington Luis, km 235

Caixa Postal 676, São Carlos, São Paulo 13565-905

(Dated: November 13, 2018)

We study an orbit of the electromagnetic two-body problem that involves a fast (stiff) spinning motion about a circular orbit. We give a multiscale method of solution that solves for the fast timescale first. The solvability condition of the asymptotic expansion demands resonances for the fast dynamics. The stiff resonant tori are found precisely in the atomic magnitude and agree with many features of the Bohr atom in quantitative and qualitative detail; We calculate every first emission line of the first 13 observable spectroscopic series of hydrogen within a few percent deviation. The resonant orbits have angular momenta that are approximate multiples of Planck's constant and the emitted frequencies are given by a difference of two linear eigenvalues. This Lorentz-invariant two-body dynamics exhibits the phenomenon of resonant dissipation, i.e., the metastable dynamics radiates the center-of-mass energy while the particles perform fast spinning oscillations of small amplitude about a circular orbit, a collective radiative recoil.

We study a two-body orbit of classical electrodynamics involving a fast (stiff) spinning toroidal motion about a circular orbit. The nontrivial stiff dynamics is part of the full two-body motion with delay and it disappears in the Coulombian limit. Our equations of motion are prescribed by Dirac's electrodynamics of *point* charges with retarded-only fields [1], a fundamental physical theory. The existence of the fast delay-dynamics demands a multiscale asymptotic solution that solves for the fast timescale first. The multiscale method requires a resonance between two fast frequencies because of Fredholm's alternative [2], providing the natural coupling between the fast and the slow timescales. The resonant orbits turn out to be precisely in the atomic magnitude; These resonant orbits are labelled by an integer and have angular momenta that are multiples of a basic angular momentum, that agrees well with Planck's constant. The emitted frequencies agree with the sharp lines of quantum electrodynamics (QED) for the circular lines of hydrogen with a few percent average deviation. The angular momentum of the stiff spinning motion is of the order of the orbital angular momentum, again in qualitative agreement with QED.

Dirac's 1938 work [1] on the electrodynamics of *point* charges gave delay equations that were seldom studied. Among the few dynamics investigated, an early result of Eliezer [3, 4, 5] revealed a surprising result (henceforth called Eliezer's theorem); An electron moving in the Coulomb field of an infinitely massive proton can never fall into the proton by radiating energy. The result was generalized to tridimensional motions with self-interaction in a Coulomb field [3, 4], finding that only scattering states are possible. Since our model has the dynamics of Eliezer's theorem as the infinite-mass limit, a finite mass for the proton is essential for a physically meaningful dynamics; If the proton has a finite mass, there is no inertial frame where it rests at all times, and

this in turn causes delay because of the finite speed of light. A finite mass for the proton is what brings delay into the electromagnetic two-body dynamics, with its associated fast dynamics. We shall describe the two-body motion in terms of the familiar *center-of-mass coordinates* and *coordinates of relative separation*, defined as the familiar coordinate-transformation that maps the two-body Kepler problem onto the one-body problem with a reduced mass plus a free-moving center of mass. We stress that in the present relativistic motion the Cartesian center-of-mass vector is not ignorable, and it represents three extra coupled degrees-of-freedom. We introduce the concept of resonant dissipation to generalize the concept of non-ionizing orbits of Ref.[6]. Resonant dissipation is the condition that both particles recoil together, i.e., the center-of-mass vector decelerates, while the particles perform the almost-circular orbit illustrated in Fig. 1, despite of the energy losses of the metastable center-of-mass dynamics.

Here we use a circular orbit as the unperturbed solution of our perturbation scheme. This circular orbit is a solution of the low-velocity Coulombian-limit obtained by ignoring the delay and its associated fast dynamics. The quantities for this uniform circular motion are given now [7]; The electron has mass m_1 and travels a circle of radius r_1 while the proton has mass m_2 and travels a circle of radius r_2 . We henceforth use units where the electronic charge is $e = -1$ and the speed of light is $c = 1$. The circular orbit is defined by the retardation angle θ that one particle turns while the light emanating from the other particle reaches it (one light-cone distance away). The angular momentum l_z of the circular orbit is given to first order by $l_z^{-1} = \theta$, and for atomic orbits θ is a small parameter of the order of the fine structure constant $\alpha = 1/137.036$. For small values of θ the frequency of angular rotation Ω and the particle distance in light-cone r_b are given, to the leading order in θ , by

Kepler's law

$$\begin{aligned}\Omega &= \mu\theta^3 + \dots \\ r_b &= \frac{1}{\mu\theta^2} + \dots\end{aligned}\quad (1)$$

where $\mu \equiv m_1 m_2 / (m_1 + m_2)$ is the reduced mass. The circular orbit is not an exact solution of the full two-body equations of Dirac's theory[1] with a finite protonic mass and self-interaction; Substituting the circular orbit into the equations of motion of [1], we find for the electronic motion an offending force opposite to the electronic velocity of magnitude

$$r_b^2 F_1 = r_b^2 \left(\frac{2}{3} \ddot{x}_1 \right) \simeq -\frac{2}{3} \theta^3, \quad (2)$$

where we scaled the offending force by the magnitude of the attractive force between the particles, a scaling factor of $1/r_b^2$. Along the trajectory of the proton, the delayed Liénard-Wiechert attraction of the electron is the main offending force against the velocity, a force of exactly the same magnitude of Eq. (2) as calculated with the Page expansion of Ref.[8]. These offending forces cause dissipation along the unperturbed circular orbit. This radiative instability in the slow timescale is an incomplete picture of the dynamics though; it disregards the fast stiff motion that is present because of the delay [7]. Precisely because of the stiff delay dynamics, at very small deviations from circularity the nonlinear terms of the self-interaction force correct the leading term of Eq. (2), enough to balance the offending force of Eq. (2). In the following we explain in physical terms how the stiff torus of Fig.1 is formed.

We start by substituting the circular orbit plus a perturbation into the equations of motion of [1]. For the stability analysis we transform the Cartesian coordinates of the particles to gyroscopic coordinates

$$\begin{aligned}x_k + iy_k &\equiv r_b \exp(i\Omega t) [d_k + \xi_k^*], \\ x_k - iy_k &\equiv r_b \exp(-i\Omega t) [d_k + \xi_k], \\ z_k &\equiv r_b (Z_k + Z_k^*)\end{aligned}\quad (3)$$

where Z_k and ξ_k are complex numbers defining the perturbations to the circular orbit. In Eq. (3) the coordinates of the electron are defined by $k = 1$ and $d_1 \simeq (m_2 + m_1)/m_2$ while the coordinates of the proton are defined by $k = 2$ and $d_2 \simeq -(m_2 + m_1)/m_1$ [7]. Notice that the perturbations of Eq. (3) are scaled by the light-cone distance r_b . The linear stability analysis proceeds like in Refs. [7, 9], by substituting Eq. (3) into the equations of motion and expanding the equations of motion to linear order. Analogously to Ref. [7], the tangent dynamics perpendicular to the plane (the z -direction) is decoupled from the tangent dynamics along the orbital plane to linear order. As found in Ref [7], the

linearized equations of motion have normal-mode solutions with every coordinate proportional to $\exp(\lambda\Omega t/\theta)$. We shall see that $|\lambda|$ can be of the order of π or larger, such that $\lambda\Omega/\theta$ is a frequency much larger than the orbital frequency Ω (here the stiffness enters). Again a remarkable quasi-degeneracy is found between the planar and the perpendicular normal-mode conditions[7] in the stiff-limit. If $|\lambda|$ is large enough (the stiff-limit), both conditions reduce to

$$\left(\frac{\mu\theta^4}{M} \right) \exp(-2\lambda) = 1, \quad (4)$$

where μ is the reduced mass defined below Eq. (1) and $M \equiv m_1 + m_2$ is the total mass of the atom. For hydrogen (μ/M) is a small factor of about (1/1824). Atomic orbits have θ of the order of the order of the fine structure constant such that the small parameter $\mu\theta^4/M \sim 10^{-13}$ multiplying the exponential function in Eq.(4) determines that $\text{Re}(\lambda) \equiv -\sigma \simeq -\ln(\sqrt{M/\mu\theta^4})$. For the first 13 excited states of hydrogen this σ is in the interval $14.0 < |\sigma| < 18.0$. The imaginary part of λ can be an arbitrarily large multiple of π , such that the general solution to Eq. (4) is

$$\lambda = -\sigma + \pi q i \quad (5)$$

where $i \equiv \sqrt{-1}$ and q is an arbitrary integer. Notice that the real part of λ is always negative, such that the tangent dynamics about circular orbits is stable in the stiff-limit with delay-only[1], at variance with the dynamics with advance and delay of Ref. [7].

The full normal-mode condition in the stiff-limit is derived in the same way of Ref. [7] and it will be published elsewhere. The perpendicular normal-mode condition is

$$\begin{aligned}& \left(1 + \frac{2}{\lambda_z} - \frac{1}{\lambda_z^2} - \frac{1}{\lambda_z^3} + \frac{1}{\lambda_z^4} + \dots \right) \left(\frac{\mu\theta^4}{M} \right) \exp(-2\lambda_z) \\ &= 1 - \frac{2}{3} \theta^2 \lambda_z + \frac{4\mu}{M} \theta^4 \lambda_z^2 + \dots,\end{aligned}\quad (6)$$

and the planar normal-mode condition is

$$\begin{aligned}& \left(1 + \frac{2}{\lambda_{xy}} + \frac{7}{\lambda_{xy}^2} + \frac{10}{\lambda_{xy}^3} - \frac{5}{\lambda_{xy}^4} + \dots \right) \left(\frac{\mu\theta^4}{M} \right) \exp(-2\lambda_{xy}) \\ &= 1 - \frac{2}{3} \theta^2 \lambda_{xy} + \frac{1}{9} \theta^4 \lambda_{xy}^2 + \dots,\end{aligned}\quad (7)$$

The coefficient of the term of order $1/\lambda^2$ is the main difference between Eqs. (6) and (7) for θ in the atomic range. The term on the right-hand side of Eqs. (6) and (7) with the $2/3$ coefficient is due to the third derivative of the self-interaction force of Eq. (2). The exact roots of Eqs. (7) and (6) near the limiting-root (5) are defined respectively by

$$\begin{aligned}\lambda_{xy}(\theta) &\equiv -\sigma_{xy} + \pi q i + i\epsilon_1, \\ \lambda_z(\theta) &\equiv -\sigma_z + \pi q i + i\epsilon_2,\end{aligned}\quad (8)$$

where $\epsilon_1(\theta)$ and $\epsilon_2(\theta)$ are real and small numbers that depend on the orbit through θ . The stiff torus is formed from an initial circular orbit as follows; (i) the offending force against the velocity makes the electron loose radius by radiating energy on a slow timescale, and (ii) when the electron deviates enough from the circular orbit, the stiff nonlinear terms compensate the offending force of Eq. (2) and balance the real parts of Eq. (8), the σ 's. The motion continues to be stiff because of the large imaginary part of the λ 's of Eq. (8). In the following we outline a multiscale expansion about the balanced stiff torus to derive a resonance condition between the modes of tangent dynamics. The circular orbit recoils in a slow timescale to compensate the momentum of the radiated energy, such that the particle coordinates are described by a composition of a translation mode and a stiff mode, both with slowly varying amplitudes

$$\begin{aligned}\xi_k &= A_k(T) + u_k(T) \exp[(\pi q i + i\epsilon_1^a)\Omega t/\theta], \\ Z_k &= B_k(T) + R_k(T) \exp[(\pi q i + i\epsilon_2^a)\Omega t/\theta],\end{aligned}\quad (9)$$

where the amplitudes u_k and R_k must be near the size a where the nonlinearity balances the negative real part of the linear modes. Because of the large frequency of the stiff motion, this balancing is established at a very small radius, as we estimate in the following.

The correction to Eq. (2) is provided by the nonlinear terms of the Lorentz-Dirac force, as given in page 116 of Ref [10]. Keeping only terms along the velocity, the self-interaction force along the electronic velocity is

$$r_b^2 F_1 = \frac{2}{3} \gamma_1^3 r_b^2 [\ddot{x}_1 + \gamma_1^2 (v_1 \cdot \dot{x}_1) v_1 + \dots], \quad (10)$$

where we again multiplied by the convenient scaling factor, as in Eq. (2). Using the stiff term of Eq. (9) into Eq. (3) to estimate the coefficient of the velocity term on the right-hand side of Eq. (10) yields

$$r_b^2 F_1 \simeq \frac{2}{3} [-\theta^3 + (\pi q)^4 a^2 \theta] = 0, \quad (11)$$

where we used $|u_1| \simeq |R_1| \simeq a$ in Eq. (9), a being the radius of the stiff torus scaled by the light-cone radius r_b . Equation (11) estimates that along the stiff torus the stiff nonlinear term balances the offending force on the electron at the amplitude

$$a = \frac{\theta}{\pi^2 q^2}. \quad (12)$$

This distance is a small fraction of the orbital radius and it represent a length of

$$r_b a = \frac{1}{\theta \pi^2 q^2}, \quad (13)$$

which for θ in the atomic scale is less than 30 classical electronic radii. This is the radius of the stiff torus,

a very reasonable physical length. Notice also that the velocity associated with the stiff motion is given by

$$v = \frac{\pi q}{\theta} \Omega r_b a = \frac{\theta}{\pi q}, \quad (14)$$

which is much lesser than the speed of light for θ in the atomic scale.

Assuming that $|u_1| = |R_1| = a$, as estimated by Eq. (12), we can use (9) to evaluate the angular momentum of the stiff torus along the orbital plane. Disregarding fast oscillatory terms, this angular momentum along the orbital plane is

$$\begin{aligned}l_x + i l_y &= \mu r_b^2 a^2 \frac{\pi q \Omega}{\theta} b_1 \exp[i(\epsilon_2^a - \epsilon_1^a + \theta)\Omega t/\theta] \\ &= \frac{\pi q}{\theta^2} a^2 \exp[i(\epsilon_2^a - \epsilon_1^a + \theta)\Omega t/\theta],\end{aligned}\quad (15)$$

where we used formulas (1) for Ω and for r_b on the second line of Eq. (15). Notice that we chose the same q in Eq. (9) to cancel the fast oscillation in Eq. (15). As illustrated in Fig. 1, the fast spinning angular momentum should rotate with the circular orbit at the frequency Ω , such that its phase should be equal to Ωt . This phase-velocity condition imposed on Eq. (15) yields the following resonance condition

$$\epsilon_1^a - \epsilon_2^a = 0 \quad (16)$$

The calculation of ϵ_1^a and ϵ_2^a necessitates expanding the equations of motion to quadratic order to include the nonlinear stiff terms (the linear stability of the stiff torus). A purely harmonic solution to these *nonlinear* equations of tangent dynamics exists because the nonlinear stiff terms introduce a correction proportional to a/θ into Eqs. (6) and (7), and the complete calculation will be given elsewhere. We expect nevertheless the quasi-degeneracy to be preserved because the stiff limit depends basically on the time light takes to travel between the particles. Once the stiff torus is very close to the circular orbit, ϵ_1^a and ϵ_2^a should differ from ϵ_1 and ϵ_2 by a correction which is essentially the same because of the quasi-degeneracy property, plus a correction of order θ , such that the difference can be well approximated by

$$\epsilon_1 - \epsilon_2 \simeq \epsilon_1^a - \epsilon_2^a + b\theta = b\theta \quad (17)$$

Condition (16) determines an orbital frequency proportional to a difference of two eigenvalues, a Rydberg-Ritz-like formula

$$\Omega = \Omega(\epsilon_1 - \epsilon_2)/b\theta = \mu \frac{\theta^2}{b} (\epsilon_1 - \epsilon_2). \quad (18)$$

where we have used Kepler's law of Eq. (1). The exact calculation of b demands the inclusion of the nonlinear terms into Eqs. (6) and (7) and shall be given elsewhere.

Since b must be of order one, we henceforth set $b = -1$ into Eq. (17) as a qualitative approximation, yielding

$$\epsilon_1 - \epsilon_2 + \theta = 0, \quad (19)$$

which has the solutions listed in Table 1. Resonance (16) can also be derived from the solvability of the multiscale asymptotic solution with inclusion of the nonlinear terms by Freedholm's alternative[2], and shall be given elsewhere [11]. In Ref. [7] we postulated heuristically a similar type of resonance for an analogous electromagnetic-like dynamical system, a resonance involving the real parts of the λ 's. The resonant orbits of Ref. [7] were in the atomic scale and had the same qualitative QED-like dependence with the integer q , which seems to be a generic feature of these stiff electromagnetic resonances of the two-body problem. The root-searching problem of Eq. (19) is well posed and for each integer q it turns out that one can find a pair of the form (8) if one sacrifices θ in Eqs. (6) and (7), i.e., θ must be quantized! According to QED, circular Bohr orbits have maximal angular momenta and a radiative selection rule ($\Delta l = \pm 1$) restricts the decay from level $k + 1$ to level k only, i.e., circular orbits emit the first line of each spectroscopic series (Lyman, Balmer, Ritz-Paschen, Brackett, etc...), the third column of Table 1. We have solved Eq. (19) and Eqs. (6) and (7) with a Newton method in the complex λ -plane. Every angular momentum $1/\theta$ determined by Eq. (19) has the correct atomic magnitude for any q . The subset of resonant orbits listed in Table 1 has orbital frequencies near a circular line of QED with a few percent deviation.

$l_z = \theta^{-1}$	$137^3 \theta^2 (\epsilon_2 - \epsilon_1)$	w_{QED}	q
185.99	3.996×10^{-1}	3.750×10^{-1}	7
307.63	8.831×10^{-1}	6.944×10^{-2}	9
475.08	2.398×10^{-2}	2.430×10^{-2}	11
577.99	1.331×10^{-2}	1.125×10^{-2}	12
694.77	7.667×10^{-3}	6.111×10^{-3}	13
826.22	4.558×10^{-3}	3.685×10^{-3}	14
973.12	2.790×10^{-3}	2.406×10^{-3}	15
1136.27	1.752×10^{-3}	1.640×10^{-3}	16
1316.44	1.127×10^{-3}	1.173×10^{-3}	17
1514.40	7.403×10^{-3}	8.678×10^{-4}	18
1730.93	4.958×10^{-3}	6.600×10^{-4}	19
1966.77	3.379×10^{-4}	5.136×10^{-4}	20
2222.70	2.341×10^{-4}	4.076×10^{-4}	21

Caption to Table 1: Numerically calculated angular

momenta $l_z = \theta^{-1}$ in units of e^2/c , orbital frequencies in atomic units $137^3 \theta^2 (\epsilon_2 - \epsilon_1)$, circular lines of QED in atomic units $w_{QED} \equiv \frac{1}{2} \left(\frac{1}{k^2} - \frac{1}{(k+1)^2} \right)$, and the values of the integer q of Eq. (8).

We identified the necessary existence of a fast timescale in the electromagnetic two-body motion near circular orbits and explored the consequences of balancing the associated fast dynamics. The balancing of the fast dynamics predicts magnitudes in qualitative and quantitative agreement with QED. The stability analysis naturally involves an integer and the angular momenta are quantized. The emission lines are in reasonable quantitative agreement with those of QED and the angular momentum of the fast spinning motion is of the order of the lowest orbital angular momentum, in agreement with QED where the electron has an angular momentum of spin given by $\sqrt{3}\hbar/2$

Caption to Figure 1: Coulombian guiding-center circular orbit of each particle, dashed lines. The trajectory of each particle is illustrated as the fast spiral encircling each guiding-center circular orbit (solid lines). Arbitrary units, distances are not on scale, illustrative purposes only.

We thank Savio B. Rodrigues and Reginaldo Napolitano for discussions.

* author's email address: deluca@df.ufscar.br

- [1] P.A.M.Dirac, Proc. R. Soc. London, Ser. A **167**, 148 (1938).
- [2] J. De Luca, R. Napolitano and V. S. Bagnato, *Phys. Rev. A* **52** R1597 (1997).
- [3] A. Carati, *J. Phys. A: Math. Gen.* **34**, 5937 (2001).
- [4] M. Marino, *J. Phys. A: Math. Gen.* **36**, 11247 (2003).
- [5] C.J. Eliezer, *Proc. Cambridge Philos. Soc.* **39**, 173 (1943).
- [6] J. De Luca, *Phys. Rev. E* **62**, 2060 (2000).
- [7] J. De Luca, *Phys. Rev. E* **71**, 056210 (2005).
- [8] J. De Luca, *Phys. Rev. Lett.* **80**, 680 (1998).
- [9] A. Staruszkiewicz, *Acta Physica Polonica* **XXXIII**, 1007(1968), see eq. (15).
- [10] F. Rohrlich, *Classical Charged Particles*, Addison-Wesley, NY (1965).
- [11] J. De Luca, to be published.

