

A Note On Stability of Steady Motion of a Rolling Disk

Milan Batista

University of Ljubljana
Faculty of Maritime Studies and Transportation
Pot pomorscakov 4, 6320 Portoroz, Slovenia
milan.batista@fpp.edu

(July 15, 2005)

Abstract

This note adds some critical remarks on the discussion presented in the McDonald's paper ([1]) on stability of steady motion of the well known problem of a disk rolling on a rough horizontal plane.

In McDonald's paper ([1]) the general analysis of small oscillations about steady motion is based on equation (equation 54 in [1])

$$\begin{aligned} \frac{\varpi^2}{\Omega_0^2}(k+1) = & 3k \cos^2 \alpha_0 + \sin^2 \alpha_0 + \frac{b}{a} \left[(6k+1) \cos \alpha_0 - (2k+1) \frac{\sin^2 \alpha_0}{\cos \alpha_0} \right] \\ & + 2 \frac{b^2}{a^2} (2k+1) \end{aligned} \quad (1)$$

but it seems that there is some incorrectness in their analysis. To clarify this, equation (1) is rewritten in the form

$$\lambda^2 = Ax^2 + Bx + C \quad (2)$$

where

$$\begin{aligned} \lambda^2 &\equiv \frac{\varpi^2}{\Omega_0^2}(k+1) \\ x &\equiv \frac{b}{a} \\ A &\equiv 2(2k+1) \\ B &\equiv (6k+1) \cos \alpha_0 - (2k+1) \frac{\sin^2 \alpha_0}{\cos \alpha_0} \\ C &\equiv 3k \cos^2 \alpha_0 + \sin^2 \alpha_0 \end{aligned} \quad (3)$$

In the next step of the analysis the authors calculate the angle at which B becomes negative

$$B \leq 0 \Rightarrow \alpha_0 \geq \arctan \sqrt{\frac{6k+1}{2k+1}} \quad (4)$$

For a uniform disk with $k = \frac{1}{4}$ this angle is $\alpha_0 \geq \arctan \frac{\sqrt{15}}{3} = 52.2^\circ$ (which differs from the value of 60° reported by the authors). From this the authors conclude that »for positive b the motion is unstable for large α_0 and that the disk will appear fall over quickly into a rolling motion with $\alpha_0 \leq 60^\circ$ «. It seems that this is slightly incorrect for the following reason. Namely, the rolling will be stable, if $\lambda^2 > 0$. For fixed α_0 this means that (2) must have no real zeros which imply that its discriminant of (2) must satisfy the condition $B^2 - 4AC < 0$. The limit is at

$$B^2 - 4AC = 0 \Rightarrow \alpha_0^{(c)} = \pm \arctan \frac{\sqrt{(2k+1)(6k+5+2\sqrt{6(k+1)(2k+1)})}}{2k+1} \quad (5)$$

which has the value $\alpha_0^{(c)} = \arctan \frac{\sqrt{39+18\sqrt{5}}}{3} \simeq 1.24 (\simeq 71.4^\circ)$ for a disk ($k = 1/2$) and $\alpha_0^{(c)} \simeq 70.8^\circ$ for a hoop ($k = 1/4$). The same value for $\alpha_0^{(c)}$ for a disk (obtained by a different procedure) is reported by Kuleshov ([2]).




Figure 1. Stability regions for a disk.

The graph of $Ax^2 + Bx + C = 0$ and associated stability regions for a disk are shown on Figure 1. Note that the value $\alpha_0 = 1.24$ represents the turning point of the graph. Note also that for very small and as well very large values of b the motion is also stable for large α_0 .

References

- [1] A.J.McDonald, K.T.McDonald, The Rolling Motion of a Disk on a Horizontal Plane, http://xxx.lanl.gov/PS_cache/physics/pdf/0008/0008227.pdf
- [2] A.S.Kuleshov. The Steady Motions of a Disc on An Rough Plane. J.Appl.Maths.Mechs. Vol 65, No 1 pp 171-173, 2001.