

A Count of Classical Field Theory Graphs

Gordon Chalmers

e-mail: gordon@quartz.shango.com

Abstract

A generating function is derived that counts the number of diagrams in an arbitrary scalar field theory. The number of graphs containing any number n_j of j -point vertices is given. The count is also used to obtain the number of classical graphs in gauge theory and gravity.

A count of scalar field theory graphs in a an arbitrary scalar Lagrangian is exactly given. The count also produces the relevant number of tree graphs in gauge theory and general relativity. Some well known work in this direction is contained in the articles [1].

The count is produced by coupling a scalar field theory to an auxiliary field that couples vertices together through the internal lines of the tree diagrams. The number of m -point vertices is deduced by keeping track of the coupling constant of the scalar and auxiliary field interaction.

The initial Lagrangian is,

$$\mathcal{L} = \frac{1}{2}\xi^2 + \sum_{j=3}^M \sum_{a=1}^{j-1} \lambda_j \xi^a \phi^{j-a} \frac{1}{a!(j-a)!} , \quad (1)$$

with the interactions $\phi^n \xi^m$. The λ_j couplings enter the graphs, with external ϕ lines and internal ξ lines. Integrating out the auxiliary ξ field produces all of these classical contributions.

The final Lagrangian is a function $\mathcal{L}(\phi, \lambda_j)$; expanding it in terms of the scalar field and the couplings λ_j generates

$$\mathcal{L}_f = \sum a_{n, \{n_j\}} \phi^n \prod_{j=1}^{\infty} \lambda_j^{n_j} , \quad (2)$$

and the number n_j tells how many j -point vertices there are in the diagram with n external lines. An n -point graph with $\sum n_j$ vertices has the multiplicity of $a_{n, \{n_j\}} n!$; this number is derived by integrating out the ξ field and expanding the expression in ϕ and the vertex couplings.

The equations of motion of ξ after a shift of $\xi \rightarrow \xi - \phi/2$ is,

$$0 = \xi - \phi/2 + \sum_{j=3}^M \frac{\lambda_j}{(j-1)!} \left((\xi + \phi/2)^{j-1} - (\xi - \phi/2)^{j-1} \right) . \quad (3)$$

In the limit of large vertex number M , i.e. $M \rightarrow \infty$, and when the couplings are identified $\lambda_j = \lambda$, the form in (3) is,

$$0 = \xi - \phi/2 + \lambda \left(-\xi - \phi/2 + \xi - \phi/2 \right) + \lambda \left(e^{\xi + \phi/2} - e^{\xi - \phi/2} \right) , \quad (4)$$




Figure 1: Example diagram with internal auxiliary field ξ and external scalar ϕ fields.

or,

$$0 = \xi + (-1/2 - \lambda)\phi + 2\lambda e^{\xi} \sinh(\phi/2) . \quad (5)$$

Shifting back $\xi \rightarrow \xi + \phi/2$ generates,

$$0 = \xi - \lambda\phi + 2\lambda e^{\xi} e^{\phi/2} \sinh(\phi/2) . \quad (6)$$

This is transcendental equation in ξ which has a formal power series solution.

The solution to (6) is obtained with the use of the Lambert solution. As only the perturbative solution in ξ is required the branch cuts are ignored. Label the parameters as,

$$a = 1 \quad b = 2\lambda e^{\phi/2} \sinh(\phi/2) \quad c = 1 \quad d = -\lambda\phi . \quad (7)$$

The general solution to this equation is,

$$\xi = -\frac{1}{ac} (a \text{WL}(\frac{cb}{a} e^{cd/a}) + cd) , \quad (8)$$

which in the case of interest in (6) is,

$$\xi = -\text{WL}(be^d) + d . \quad (9)$$

The Lambert function $WL(x)$ has the power series expansion of

$$W(x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^{n-2}}{(n-1)!} x^n , \quad (10)$$

and leads to the form,

$$\xi = d - \sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^{n-2}}{(n-1)!} b^n e^{nd} \quad (11)$$

$$\xi = -\lambda\phi - \sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^{n-2}}{(n-1)!} 2^n \lambda^n \sinh^n(\phi/2) e^{n\phi(1/2-\lambda)} . \quad (12)$$

The solution to ξ is used to resubstitute into the initial Lagrangian to obtain the power series expansion in ϕ and λ .

The expansion in ϕ and λ is

$$\mathcal{L} = \sum a_{n,m} \phi^n \lambda^m , \quad (13)$$

and counts the diagrams with multiple vertices at $n-2$ -point. The count of diagrams can be obtained with any arbitrary vertex configuration, i.e. any numbers n_j , with the λ coupling only; that is, this is simpler than retaining all of the λ_j couplings. An n -point diagram with only 3-point vertices has $n_3 = n-2$ and the n -point diagram with only a 3-point and $n-2$ vertex has $n_3 = n_{n-2} = 1$. These are the vertex bounds without external ξ fields. Define $m = \sum n_j$ as the sum of all of the n_j vertices; then $a_{n,m}$ counts the diagrams for m at n point. A normalization factor of $n!$ is required to compensate for the symmetry factor in the final scalar Lagrangian.

As an example, integrating out the auxiliary field that contributes to four-point diagrams gives,

$$\frac{1}{2}\xi^2 + \frac{1}{2}\xi\phi^2 \rightarrow -\frac{1}{2^3}\phi^4 . \quad (14)$$

The symmetry factor of 3 comes from the three diagrams that contribute, those in the s , u , and t channels. Thus, $4!$ times 2^{-3} equals 3, and this is how many diagrams

there are at four-point which stem from the three-point vertices. The more general case is examined with the use of solving the transcendental equation and re-inserting the solution into the Lagrangian in (1).

The series expansion complicated by the multiple sums involved. A Taylor series expansion is used to extract the appropriate n and m .

$$\mathcal{L}_f^{(1)} = \frac{1}{2} \left(\lambda \phi + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^{n-2}}{(n-1)!} 2^n \lambda^n \sinh^n(\phi/2) e^{n\phi(1/2-\lambda)} \right)^2 \quad (15)$$

$$\mathcal{L}_f^{(2)} = \lambda \sum_{j=3}^{\infty} \sum_{a=1}^{j-1} (-1)^a \phi^{j-a} \frac{1}{a!(j-a)!} \sum_{b=0}^a \frac{a!}{b!(a-b)!} \lambda^{b-a} \phi^{b-a} \quad (16)$$

$$\times \left(\sum_{n=1}^{\infty} (-1)^{n-1} \frac{n^{n-2}}{(n-1)!} 2^n \lambda^n \sinh^n(\phi/2) e^{n\phi(1/2-\lambda)} \right)^b \quad (17)$$

The counting numbers $a_{n,m}$ are then derived from

$$a_{n,m} = \frac{1}{n!m!} \partial_{\phi}^n \partial_{\lambda}^m \left(\mathcal{L}_f^{(1)} + \mathcal{L}_f^{(2)} \right). \quad (18)$$

These derivatives are straightforward to find, but tedious.

The ∂_{ϕ}^n derivatives distribute in the manner, with $n = \alpha_1 + \sum^{\alpha_2} \beta_i$,

$$H^{\alpha_1, \alpha_2; p_i; \beta_i} = \partial_{\phi}^{\alpha_1} \phi^{b+j-2a} \prod_{i=1}^{\alpha_2} \partial_{\phi}^{\beta_i} \left(\sinh^{p_i}(\phi/2) e^{p_i \phi(1/2-\lambda)} \right) \quad (19)$$

$$= \frac{(b+j-2a)!}{(b+j-2a-\alpha_1)!} \delta_{b+j-2a, \alpha_1} \times \prod_{i=1}^{\alpha_2} 2^{-p_i} \sum_{q=0}^{p_i} \frac{p_i!}{q!(p_i-q)!} (-1)^{p_i-q} (q-p_i \lambda)^{\beta_i}, \quad (20)$$

with the b th power of the sum in the parenthesis changing to $b-\alpha_2$ and a factor present of $b!/(b-\alpha_2)!$. All possible combinations are required, including the α_2 factors from the sums of the object in the parenthesis. This results, after evaluating the expression at $\phi = 0$, in

$$\lambda \sum_{j=3}^{\infty} \sum_{a=1}^{j-1} (-1)^a \frac{1}{a!(j-a)!} \sum_{b=0}^a \frac{a!}{b!(a-b)!} \lambda^{b-a} \quad (21)$$

$$\times \sum_{\alpha_1, \alpha_2; \beta_i}^{n=\alpha_1+\sum \beta_i} \delta_{b, \alpha_2} \frac{b!}{(b-\alpha_2)!} \sum_{\tilde{p}=p_1+\dots+p_{\alpha_2}; p_i} (-1)^{\tilde{p}+\alpha_2} \lambda^{\tilde{p}} \prod_{i=1}^{\alpha_2} \frac{p_i^{p_i-2}}{(p_i-1)!} \quad (22)$$

$$\times H^{\alpha_1, \alpha_2; p_i; \beta_i} \quad (23)$$

The sums include $n = \alpha_1 + \sum \alpha_2 \beta_i$, and α_2 ranging from 1 to n . The β_i must each be at least 1. There appear to be indefinite sums on the p_i due to the nested sums in the parenthesis of (17).

The coupling derivatives act as

$$\partial_\lambda^m \lambda^{b-a+\tilde{p}} \prod_{i=1}^{\alpha_2} (q-p\lambda)^{\beta_i} \quad (24)$$

$$= \sum_{r=0}^m \frac{(b-a+\tilde{p})!}{(b-a+\tilde{p}-r)!} \delta_{b-a+\tilde{p}, m} \sum_{\gamma_i} \prod_{i=1}^{\alpha_2} \frac{\beta_i!}{(\beta_i-\gamma_i)!} q^{\beta_i-\gamma_i} (-p)^{\gamma_i} |_{\sum \gamma_i=b-a+\tilde{p}-r} \cdot \quad (25)$$

These derivatives are taken on the couplings in (23), and the factors must be placed together. The two results, in (23) and (25), generate the expression for $a_{m,n}$ after dividing by $m!n!$.

The total contribution to $a_{n,m}$ from \mathcal{L}_2 is the complicated expression,

$$\lambda \sum_{j=3}^{\infty} \sum_{a=1}^{j-1} (-1)^a \frac{1}{a!(j-a)!} \frac{a!}{\alpha_2!(a-\alpha_2)!} \quad (26)$$

$$\times \sum_{\alpha_1, \alpha_2; \beta_i}^{n=\alpha_1+\sum \beta_i} \sum_{\tilde{p}=m+a-\alpha_2; p_i} (-1)^{\tilde{p}+\alpha_2} \prod_{i=1}^{\alpha_2} \frac{p_i^{p_i-2}}{(p_i-1)!} \quad (27)$$

$$\frac{(\alpha_2+j-2a)!}{(\alpha_2+j-2a-\alpha_1)!} \delta_{\alpha_2+j-2a, \alpha_1} \times \prod_{i=1}^{\alpha_2} 2^{-p_i} \sum_{q=0}^{p_i} \frac{p_i!}{q!(p_i-q)!} (-1)^{p_i-q} \quad (28)$$

$$\sum_{r=0}^m \frac{m!}{(m-r)!} \sum_{\gamma_i} \prod_{i=1}^{\alpha_2} \frac{\beta_i!}{(\beta_i-\gamma_i)!} q^{\beta_i-\gamma_i} (-p)^{\gamma_i} |_{\sum \gamma_i=b-a+\tilde{p}-r} \cdot \quad (29)$$

The contribution from \mathcal{L}_1 is similar and can be determined from the same operations. α_1 and α_2 are defined as before. The sums also could be simplified more, and the large n expansion generates the known exponential dependence.

These combinatorical factors can also be used to find the multiplicity of gauge and gravity diagrams. In Gervais-Neveu gauge, the 3-point and 4-point vertices of a non-abelian gauge theory contain 6 and 12 terms in a diagram without color ordering. The individual scalar field contributions with n_3 and n_4 vertices (3 and 4-point) expand into an additional $3^{n_3}4^{n_4}$ combinations; the propagator contains only a single $\eta_{\mu\nu}$ and does not cause further multiplicity.

Closed form expressions for graph multiplicities in quantum field theory are given. The count pertains to scalar field theory with any number and type of vertices. The number of diagrams $a_{n,m}$ counts the classical field diagrams at n -point containing $\sum v_i = m$ vertices, with i being the number of lines at each vertex. The count is useful for determining the naive complexity of tree graph calculations in quantum field theory. Also, the count is useful in determining Kähler potentials in toric varieties, which requires the number of tree graphs at zero momentum in order to determine the D-terms.

References

- [1] C. M. Bender and T. T. Wuk, Statistical Analysis Of Feynman Diagrams, Phys. Rev. Lett. **37**, 117 (1976); Large Order Behavior Of Perturbation Theory, Phys. Rev. Lett. **27**, 461 (1971).
- [2] G. Chalmers, *Quantum Solution to Scalar Field Models*, physics/0505018.
- [3] G. Chalmers, *Quantum Gauge Amplitude Solutions*, physics/0505077.
- [4] G. Chalmers, *Tree Amplitudes in Gauge and Gravity Theories*, physics/0504219.
- [5] G. Chalmers, *Tree Amplitudes in Scalar Field Theories*, physics/0504173.
- [6] G. Chalmers, *Derivation of Quantum Field Dynamics*, physics/0503062.