

ON THE ORIGIN OF THE ANOMALOUS PRECESSION OF MERCURY'S PERIHELION

JAUME GINÉ

ABSTRACT. Action at distance in Newtonian physics is replaced by finite propagation speeds in classical post-Newtonian physics. As a result, the differential equations of motion in Newtonian physics are replaced by functional differential equations, where the delay associated with the finite propagation speed is taken into account. Newtonian equations of motion, with post-Newtonian corrections, are often used to approximate the functional differential equations. In [12] a simple atomic model based on a functional differential equation which reproduces the quantized Bohr atomic model was presented. The unique assumption was that the electrodynamic interaction has a finite propagation speed. In [13] a simple gravitational model based on a functional differential equation which gives a gravitational quantification and an explanation of the modified Titius-Bode law is described. Firstly, in this work, we recall the calculations made by Einstein to arrive at the explanation of the anomalous precession of Mercury's perihelion. Secondly, we recover an ancient work of Gerber in 1898 as a precursor of the retarded theories, see [9]. In this paper Gerber gave an explanation of the anomalous precession of the Mercury's perihelion in terms of a velocity-dependent potential. In this paper an explanation of the anomalous precession of Mercury's perihelion is given in terms of a simple retarded potential, which, at first order, coincides with Gerber's potential, and which agrees with the author's previous works [12, 13].

1. INTRODUCTION

In 1859 the French astronomer Le Verrier announced, based on many years of careful observations and calculations, that the perihelion of the planet Mercury evidently precesses at a slightly faster rate than can be accounted for by Newtonian mechanics given the known distribution of

1991 *Mathematics Subject Classification.* Primary 34C05. Secondary 58F14.

Key words and phrases. gravitation, retarded systems, functional differential equations, limit cycle.

The author is partially supported by a DGICYT grant number BFM 2002-04236-C02-01 and by DURSI of Government of Catalonia "Distinció de la Generalitat de Catalunya per a la promoció de la recerca universitària".

material in the solar system. This discovery led to two different avenues of research. Some people began to search for unknown material in the solar system (possibly even a new planet) that would account for the anomaly in Mercury's orbit within the context of Newton's laws. Others considered possible ways of modifying or re-interpreting Newton's law of gravitation so that it would give Mercury's precession for the currently accepted distribution of matter. For a complete description of the historical development of the problem see [16].

It should be noted that Mercury is not the only object in the solar system that exhibits anomalous precession. The effect is most noticeable for objects near the Sun with highly elliptical orbits, but it can be seen even in the nearly circular orbits of Venus and Earth, although the discrepancy isn't nearly so large as for Mercury. In addition, the asteroid Icarus is ideal for studying this effect, because it has an extremely elliptical orbit and periodically passes very close to the Sun.

Einstein found that the extra precession arises unavoidably from the fundamental principles of General Relativity. The general problem of the integration of the Einstein equations, given by

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu},$$

where $R_{\mu\nu}$ is the Ricci tensor, R is the Ricci scalar, $g_{\mu\nu}$ is the metric tensor and $T_{\mu\nu}$ is the stress-energy tensor, is extremely difficult and the determination of the explicit solutions is only possible in a restrict number of cases. One of the most important is the Schwarzschild solution for the case of a punctual mass or spherical and homogeneous and with the assumption that the limit values in the infinite of the $g_{\mu\nu}$ are the galilean values. To determine the relativistic prediction for the advance of an elliptical orbit, we take polar coordinates centered in the center of mass of the central body. Assuming that the orbits remain permanently in the equatorial plane (as in Newtonian theory), let's work in this single plane $S = 0$, so of course dS/dt and all higher derivatives also vanish. Then, the term involving S in the Schwarzschild metric drops out, leaving just

$$d\tau^2 = \left(\frac{r-2m}{r}\right) dt^2 - \left(\frac{r}{r-2m}\right) dr^2 - r^2 d\theta^2,$$

taking $c = 1$. The equations of geodesic motion are

$$(1) \quad \frac{d^2 t}{d\tau^2} = -\frac{2m}{r(r-2m)} \left(\frac{dr}{d\tau}\right) \left(\frac{dt}{d\tau}\right),$$

$$(2) \quad \frac{d^2 r}{d\tau^2} = -\frac{m(r-2m)}{r^3} \left(\frac{dt}{d\tau} \right)^2 + \frac{m}{r(r-2m)} \left(\frac{dr}{d\tau} \right)^2 + (r-2m) \left(\frac{d\theta}{d\tau} \right)^2,$$

$$(3) \quad \frac{d^2 \theta}{d\tau^2} = -\frac{2}{r} \left(\frac{dr}{d\tau} \right) \left(\frac{d\theta}{d\tau} \right).$$

We can immediately integrate equations (1) and (3) to give

$$\frac{dt}{d\tau} = \frac{kr}{r-2m}, \quad \frac{d\theta}{d\tau} = \frac{h}{r^2},$$

where k and h are constants of integration, determined by the initial conditions of the orbit. We can now substitute for these into the basic Schwarzschild metric divided by $d\tau^2$ to give

$$1 = \left(\frac{k^2 r}{r-2m} \right) - \left(\frac{r}{r-2m} \right) \left(\frac{dr}{d\tau} \right)^2 - \frac{h^2}{r^2}.$$

Solving for $(dr/d\tau)^2$, we have

$$(4) \quad \left(\frac{dr}{d\tau} \right)^2 = \frac{2mh^2}{r^3} - \frac{h^2}{r^2} + \frac{2m}{r} + k^2 - 1$$

If we insert the squared derivatives of coordinates into equation (2), we obtain

$$\frac{d^2 r}{d\tau^2} = -\frac{m}{r^2} + \frac{h^2}{r^3} - \frac{3mh^2}{r^4}.$$

We arrive at this same equation if we differentiate equation (4) with respect to τ and we divide by $2(dr/d\tau)$. Let $\omega = d\theta/d\tau$ be the proper angular speed, we have $h = \omega r^2$, and the above equation can be written as

$$\frac{d^2 r}{d\tau^2} = -\frac{m}{r^2} + \omega^2(r-3m).$$

Obviously if $\omega = 0$ this gives the Newton's inverse-square law for radial gravitational acceleration. With non-zero ω the term $\omega^2 r$ corresponds to the Newtonian centripetal acceleration which, if we defined the tangential velocity $v = \omega r$, would equal the classical v^2/r . This term serves to offset the inward pull of gravity, but in the relativistic version we find not $\omega^2 r$ but $\omega^2(r-3m)$. For values of r much greater than $3m$ this difference can be neglected, but clearly if r approaches $3m$ we can expect to see non-classical effects, and of course if r ever becomes less than $3m$ we would expect completely un-classical behavior. In fact, this corresponds to the cases when an orbiting particle spirals into the center, which never happens in classical theory.

In a similar way as in the resolution of the classical case, we introduce the change of variable $u(\theta) = 1/r(\tau)$. Differentiating u respect to θ we have $du/d\theta = -(1/r^2)(dr/d\tau)(d\tau/d\theta)$. Taking into account that $d\theta/d\tau = h/r^2$, we have $dr/d\tau = -h(du/d\theta)$. Substituting $dr/d\tau$ and $1/r$ into equation (4) we obtain the following differential equation

$$h^2 \left(\frac{du}{d\theta} \right)^2 = 2mh^2u^3 - h^2u^2 + 2mu + k^2 - 1.$$

Differentiating with respect to θ and dividing by $2h^2(du/d\theta)$, we get

$$\frac{d^2u}{d\theta^2} + u = \frac{m}{h^2} + 3mu^2.$$

Solving this quadratic equation with respect to u and taking the solution which has physical sense, we obtain

$$u = \frac{1}{6m} \left(1 - \sqrt{1 - 12 \left(\frac{m^2}{h^2} - m \frac{d^2u}{d\theta^2} \right)} \right).$$

The quantity inside the parentheses and under the square root is typically quite small compared with 1, therefore we can approximate the square root by the first few terms of its expansion

$$u \approx \frac{1}{6m} \left[6 \left(\frac{m^2}{h^2} - m \frac{d^2u}{d\theta^2} \right) + \frac{3}{2} \left(\frac{m^2}{h^2} - m \frac{d^2u}{d\theta^2} \right)^2 \right].$$

Expanding the right hand side and re-arranging terms, we have

$$\left(1 + 6 \left(\frac{m}{h} \right)^2 \right) \frac{d^2u}{d\theta^2} + u = \left(\frac{m}{h^2} + \frac{3m^3}{h^4} \right) + 3m \left(\frac{d^2u}{d\theta^2} \right)^2.$$

The value of $d^2u/d\theta^2$ in typical astronomical problems is numerically quite small (many orders of magnitude less than 1), so the quantity $3m(d^2u/d\theta^2)^2$ on the right hand side will be negligible for planetary motions. Therefore, we obtain a simple harmonic oscillator of the form

$$M \frac{d^2u}{d\theta^2} + u = F,$$

where M and F are constants. For some choice of initial θ the general solution of this equation can be expressed as $u(\theta) = F[1 + k \cos(\theta/\sqrt{M})]$, where k is a constant of integration. Therefore, reverting back to the parameter $r = 1/u$, we have

$$r(\theta) = \frac{1}{F[1 + k \cos(\Omega \theta)]}, \quad \text{where} \quad \Omega = \frac{1}{\sqrt{1 + 6 \left(\frac{m}{h} \right)^2}}.$$

To determine the relativistic prediction for the orbital precession of the planetary orbits, we can expand the expression for Ω as follows

$$\Omega = \frac{1}{\sqrt{1 + 6\left(\frac{m}{h}\right)^2}} = 1 - 3\left(\frac{m}{h}\right)^2 + \frac{27}{2}\left(\frac{m}{h}\right)^4 - \dots$$

Since m/h is so small, we can take just the first-order term, and noting that one cycle of the radial function will be completed when $\Omega\theta = 2\pi$, we see that θ must increase by $2\pi/\Omega$ for each radial cycle, so the precession per revolution is

$$\frac{2\pi}{\Omega} - 2\pi = 2\pi \left(\frac{1}{1 - 3\left(\frac{m}{h}\right)^2} - 1 \right) \approx 6\pi \left(\frac{m}{h}\right)^2.$$

We notice that the semi-latus L is approximately h^2/m , so the amount of precession per revolution (for slow moving objects in weak gravitational fields, such as the planets in our solar system) can be written as simply as $6\pi m/L$, where m is the gravitational radius of the central body. Taking into account the values for the Sun and the Mercury planet, we obtain the value of 42.9195 arc seconds per century.

The observed precession of 43.1 ± 0.5 arc seconds per century for the planet Mercury is in close agreement with the theory. In 1915, Einstein [6, 7] proudly concluded his presentation of the vacuum field equations on General Relativity by pointing out that they explained the anomalous precession of Mercury.

2. GERBER'S THEORY

At the end of the 19th century, theoretical physicists such as Gauss and Weber were investigating modifications of the Coulomb inverse-square law by introducing a velocity-dependent potential to represent the electromagnetic field, consistent with the finite propagation speed of changes in the field (i.e., the speed of electromagnetic waves). It was found that this same type of law, when applied to gravitation, predicted perihelion advance for the two-body problem on the same order of magnitude as actually observed for Mercury. It became a fairly popular activity in the 1890s for physicists to propose various gravitational potentials based on finite propagation speed in order to account for some or all of Mercury's orbital precession. Oppenheim published a review of these proposals in 1895, see [18, 19]. In fact, this line of investigation goes back to the works of Laplace [14] in 1805 where it is presented a correction of the Newton force produced by the

particle m_1 in m_2 , which moves with velocity v given by

$$\mathbf{F} = -Gm_1m_2 \left(\frac{\mathbf{r}}{r^3} + \frac{\mathbf{v}}{h} \right),$$

where h is the finite propagation speed. But this work didn't find echo practically until the surroundings of 1880, when a series of works to estimate the gravitational finite propagation speed began. A brief list of authors that used the hypothesis of the finite propagation speed is Th. von Oppolzer (1883), J. von Hepperger (1889), R. Lehmann-Filhes (1894), K. Schwarzschild (1900), H. Minkowski (1908), H. Poincaré (1908), W. Ritz (1909). In other works different forms for the gravitational potential were proposed; we can mention H. von Seeliger (1895) and C. Neumann (1896). Under the influence of the electrodynamical development made by F. Neumann (1845), W. Weber (1846) and B. Riemann (1861), some authors began to think in modifying Newton's law adding terms which depend on the speeds of the involved bodies, see for instance [24]. In 1870 F.G. Holzmüller [8] proposed a law of gravitation of the same form that the electrodynamic Weber's law, given by

$$F = \frac{Gm_1m_2}{r^2} \left(1 - \frac{\dot{r}^2}{h^2} + \frac{2r\ddot{r}}{h^2} \right).$$

Later, F. Tisserand [23] had used this law to study the anomalous precession of Mercury's perihelion and he explained only 14.1 arc seconds per century. In the same way O. Liman (1886) and M. Lévi (1890), proposed a law of gravitation of the same form that the electrodynamic Riemann's law, given by

$$F = \frac{Gm_1m_2}{r^2} \left(1 - \frac{(\dot{r}_1 - \dot{r}_2)^2}{h^2} \right),$$

where r_1 and r_2 are the position vectors of the particles m_1 and m_2 , respectively. The Riemann-Liman-Lévi law explained only 28 arc seconds per century of the anomalous precession of Mercury's perihelion. Finally, M. Lévi, by means of a purely formal development, found a force law that led to the observed exact value of the anomalous precession of Mercury's perihelion. The theories to explain the form of the proposed law forces are based, in general, in to do a parallelism between the electromagnetism and the gravitation and to propose what is known as gravitational field with a gravelectric component and with a gravomagnetic component, see [1, 15] and references therein. Below, in the next section, we will see that all these laws are, in fact, developments until certain order of a retarded potential.

These lines of research were abandoned when it was definitively implanted Einstein's Relativity theory. The typical result of such proposals is a predicted non-Newtonian advance of orbital perihelia per revolution of $C\pi m/(Lc^2)$ where c is the posited speed of propagation, m is the Sun's mass, L is the semi-latus rectum of the orbit, and C is a constant depending on the precise form of the assumed potential. Of course, by Kepler's law, we can express the mass m of the Sun in terms of the semi-major axis a and the angular speed ω of an orbiting planet as $m = a^3\omega^2$. The angular speed can be expressed in terms of the orbital period T (revolutions per second) as $\omega = 2\pi/T$. In addition, we know from elementary geometry that the semi-latus rectum of an ellipse is $L = a(1 - e^2)$ where e is the eccentricity. Making these substitutions, the angular precession per orbit can be written in the form

$$\frac{C\pi m}{Lc^2} = \frac{4\pi^3 a^2 C}{c^2 T^2 (1 - e^2)}.$$

As we have seen with $C = 6$ this formula gives the observed non-Newtonian precession for planetary orbits, most notably Mercury's 43 arc seconds per century, so the objective was to find a potential that gives $C = 6$. Of course, it's trivial to do this in an ad hoc manner, but scientists were attempting to find some rational basis for a potential with the required property. The most straightforward way of incorporating a finite propagation speed into the law of gravity is to simplistically substitute the retarded radial distance for the present distance, which leads to a velocity-dependent potential of the form

$$V(r, \dot{r}) = -\frac{m}{r} \frac{1}{\left(1 - \frac{\dot{r}}{c}\right)},$$

but this V results in a value of $C = 2$, so it predicts only 14.1 arc seconds per century for the anomalous precession of Mercury's perihelion, one third of the observed value.

In 1898 a German school teacher named Paul Gerber wrote a paper in which he proposed a velocity-dependent potential that gives $C = 6$, see [9]. Gerber prepared an expanded version of this paper in 1902, and this expanded paper was subsequently republished in 1917 in the *Annalen der Physik* [10] at the urging of Ernst Gehrcke, who wished to call attention to Gerber's approach as a possible alternative to Einstein's General Relativity. In [16] it is concluded with a speculative re-construction of a semi-classical line of reasoning by which it is actually possible to derive Gerber's potential, albeit in a way that evidently never occurred to Gerber.

The contribution to the Newtonian potential at a given location in space at a given instant due to a mass m is strictly a function of the distance of that mass from the given location at the same instant. In other words, if we let $r(t)$ denote the distance from the given location to the mass particle at the time t , then the contribution of that mass to the gravitational potential $V(t)$ at the given location at the time t is $V(t) = -m/r(t)$. Thus, Newtonian gravity represents instantaneous action at a distance, because the effect of changing the position of a gravitating mass is felt throughout the universe at the instant when the mass moves. Needless to say, for the simple two-body problem this potential gives stable elliptical orbits.

However, suppose the gravitational potential experienced by an object depends not only on the radial distance from the gravitating source but also on the derivative (with respect to time) of that distance. Specifically, suppose the potential has the form

$$(5) \quad V(r, \dot{r}) = -\frac{m}{r} \frac{1}{\left(1 - \frac{\dot{r}}{c}\right)^2}.$$

We are going to see that this velocity-dependent potential results in elliptical orbits that precess by the same amounts as predicted by General Relativity (to the lowest order of approximation), and of course this fact agrees with the observed precession rates for the perihelia of the planets, including Mercury (see below). The question, then, is whether we can justify the use of this particular velocity-dependent potential rather than the Newtonian potential $V = -m/r(t)$. The rational interpretation is totally discerned in [16]. We now review the steps by which the correct precession is given by Gerber's potential (5). We will see that Gerber's velocity-dependent potential gives the same prediction for the precession of orbital perihelia (to the first order for weak fields) as does General Relativity.

On this basis, the proposal of a velocity-dependent potential function (5) corresponds to the proposal of a certain gravitational force law, given by substituting the potential function (5) into the equation

$$\begin{aligned} f &= \frac{d}{dt} \left(\frac{\partial V}{\partial \dot{r}} \right) - \frac{\partial V}{\partial r} \\ &= -\frac{m}{r^2} \left(1 - \frac{\dot{r}}{c}\right)^{-4} \left(\frac{6r\ddot{r}}{c^2} - \frac{2\dot{r}}{c} \left(1 - \frac{\dot{r}}{c}\right) + \left(1 - \frac{\dot{r}}{c}\right)^2 \right). \end{aligned}$$

Expanding this expression in powers of $\dot{r}/c$, we get

$$(6) \quad f = -\frac{m}{r^2} \left(1 - \frac{3\dot{r}^2}{c^2} + \frac{6r\ddot{r}}{c^2} - \frac{8\dot{r}^3}{c^3} + \frac{24r\dot{r}\ddot{r}}{c^3} - \dots \right).$$

Since equation (6) represents the force per unit mass, directed toward the central gravitating mass m , it is equal to the radial acceleration. To determine the acceleration of a point in the plane, we consider polar coordinates (r, θ) , where $\omega = d\theta/dt$ is the angular velocity. Hence the radial and tangential components of the point's acceleration are:

$$a_r = \ddot{r} - r\omega^2, \quad a_t = r\dot{\omega} + 2\dot{r}\omega.$$

There is no tangential force on the orbiting particle (according to Gerber's model), so we have

$$r\dot{\omega} + 2\dot{r}\omega = \frac{1}{r} \frac{d}{dt}(r^2\omega) = 0.$$

It follows that the quantity inside the parentheses is a constant, which we will denote as usually by $h = r^2\omega$. Equating the radial acceleration with the radial specific force (up to second order), and reverting to units such that $c = 1$, we get

$$(7) \quad \ddot{r} - r\omega^2 = -\frac{m}{r^2}(1 - 3\dot{r}^2 + 6r\ddot{r}).$$

If we define, as before, $u(\theta) = 1/r(t)$, then we have

$$\begin{aligned} \dot{r} &= -\frac{1}{u^2} \frac{du}{dt} = -\frac{h}{\omega} \frac{du}{d\theta} \frac{d\theta}{dt} = -h \frac{du}{d\theta}, \\ \ddot{r} &= \frac{d\dot{r}}{dt} = -h \frac{d}{dt} \left(\frac{du}{d\theta} \right) = -h \frac{d^2u}{d\theta^2} \frac{d\theta}{dt} = -h \frac{d^2u}{d\theta^2} \omega = -h^2 u^2 \frac{d^2u}{d\theta^2}, \end{aligned}$$

Substituting these expressions in (7) and re-arranging terms, we obtain

$$(8) \quad (1 + 6mu) \frac{d^2u}{d\theta^2} + u = \frac{m}{h^2} \left(1 - 3h^2 \left(\frac{du}{d\theta} \right)^2 \right),$$

The quantities inside the parentheses are both nearly equal to unity, because the terms added to or subtracted from 1 are many orders of magnitude less than 1 (for astronomical orbits). Moreover, as before, the quantity $3h^2(du/d\theta)^2$ will be negligible for planetary motions. Therefore, this equation is essentially a harmonic oscillator of the form

$$A \frac{d^2u}{d\theta^2} + u = \frac{m}{h^2},$$

where A is taken to be nearly equal to unity. For a suitable choice of the reference angle θ , the general solution of this equation can be expressed as $u(\theta) = (m/h^2)[1 + k \cos(\theta/\sqrt{A})]$, where k is a constant of integration. Notice that, if the value of A is consistently greater than 1, even if only by a very small amount, there is a cumulative effect, because the angle θ must increase by more than 2π in order for

the system to return to the same radial position. On each revolution this extra angular displacement accumulates, so the spatial path of the orbit undergoes a steady precession.

Reverting back to the parameter $r = 1/u$, the orbital equation is

$$r(\theta) = \frac{h^2}{m[1 + k \cos(\sigma \theta)]}, \quad \text{where} \quad \sigma = \frac{1}{\sqrt{1 + 6\frac{m}{r}}}.$$

If σ were equal to unity, this would be the polar equation of an ellipse with the pole at one focus, and with the constant k signifying the eccentricity and h^2/m signifying the semi-latus rectum L . The actual value of σ depends on r , but for orbital motion the mean value of r can be expressed in terms of the orbital constants. This fact is more easily seen for the case of a circular orbit, so the derivatives of u with respect to θ are all zero, and equation (8) immediately gives $r = h^2/m$ (which of course is equivalent to Kepler's third law, $m = r^3 w^2$, since $h = mwr^2$). Substituting for r in the expression for σ , we get

$$\sigma = \frac{1}{\sqrt{1 + 6\left(\frac{m}{h}\right)^2}}.$$

This value is the same as Ω , as before, and therefore it gives the same as the first-order non-Newtonian precession predicted by General Relativity for weak fields. Gerber's potential also gives values for all other known bodies orbiting the Sun consistent with observation.

We should mention that although General Relativity and Gerber's potential predict the same first-order precession (for weak fields), the respective equations of motion are not identical, even at the first non-Newtonian level of approximation. In terms of the parameter $u = 1/r$ the equations of motion are

$$\begin{aligned} \frac{d^2 u}{d\theta^2} + u &= \frac{m}{h^2} + 3mu^2, \\ \frac{d^2 u}{d\theta^2} + u &= \frac{m}{h^2} - 3m \left(\frac{du}{d\theta} \right)^2 - 6mu \frac{d^2 u}{d\theta^2} + \dots, \end{aligned}$$

so the non-Newtonian terms are actually quite different. Of course, any non-Newtonian term will lead to orbits that fail to close, so there will be some cumulative precession for the two-body problem. It just happens that the term $3mu^2$ in the General Relativity equation of motion and the term $6mu(d^2 u/d\theta^2)$ in Gerber's equation of motion both result in a first-order precession of $6\pi m/L$ in the slow weak-field limit. Thus Gerber did not in any way anticipate the two-body equation of motion

predicted by General Relativity, let alone the field equations from which the relativistic equation of motion is derived.

The 1902 paper begins with 20 pages of text in which Gerber writes in vague and qualitative terms about the idea that gravity has a finite speed of propagation, and reviews the previous attempts to infer the speed of gravity from observations of the solar system. Following Oppenheim's review article closely, he discusses Laplace, Weber, Riemann, Neumann, Lehmann-Filhes and Hepperger, Clausius, and Lorentz. It is not until Section IV that he begins to present his idea in precise form, giving his derivation of the potential (5). In the introduction to the 1917 re-print of Gerber's paper, Ernst Gehrcke concludes:

Whether and how the theory of Gerber can be merged with the well-known electromagnetic equations into a new unified theory is a difficult problem, which still awaits a solution.

3. A SIMPLE RETARDED POTENTIAL

Action at distance in Newtonian physics is replaced by finite propagation speeds in classical post-Newtonian physics. As a result, the differential equations of motion in Newtonian physics are replaced by functional differential equations, where the delay associated with the finite propagation speed is taken into account. Newtonian equations of motion, with post-Newtonian corrections, are often used to approximate the functional differential equations, see, for instance, [2, 3, 4, 5, 11, 21, 22]. In [12] a simple atomic model based on a functional differential equation which reproduces the quantized Bohr atomic model was presented. The unique assumption was that the electrodynamic interaction has finite propagation speed, which is a consequence of the Relativity theory. An straightforward consequence of the theory developed in [12], and taking into account that gravitational interaction has also a finite propagation speed, is that the same model is applicable to the gravitational 2-body problem. In [13] a simple gravitational model based on a functional differential equation which gives a gravitational quantification and an explanation of the modified Titius-Bode law is described. In the following an explanation of the anomalous precession of Mercury's perihelion is given in terms of a simple retarded potential, which, at first order, coincides with the Gerber's potential.

The most straightforward way of incorporating a finite propagation speed into the law of gravity is to simplistically substitute the present

distance for the retarded radial distance, therefore, we consider the simplest retarded potential

$$(9) \quad V = -\frac{m}{r(t - \tau)},$$

where $r(t)$ denotes the instantaneous position vector of the test particle, at time t , and τ is the delay, so that $r(t - \tau)$ is the retarded position of the test particle. In fact this retarded potential depends on the position vector but also on the velocity vector $\dot{r}$, on the acceleration vector $\ddot{r}$ and so on. The appearance of a delay implies all these dependences in the potential. From the retarded potential (9) we will obtain, in a theoretical point of view, the equation of motion of the particle. This equation will be a functional differential equation. The functional differential equations of motion are generally difficult, often impossible, to express in a form that is amenable to analysis. Thus, in order to obtain useful dynamical predictions from realistic models, it is frequent to replace the functional differential equations of motion by approximations that are ordinary or partial differential equations, see [2]. In our case, if we develop the retarded potential (9) in powers of τ (up to second order in τ), we obtain

$$(10) \quad V \approx -\frac{m}{r} \left[1 + \frac{\dot{r}}{r} \tau + \left(\frac{\dot{r}^2}{r^2} - \frac{\ddot{r}}{2r} \right) \tau^2 \right],$$

To develop some easier calculations we can reject on the right hand side of expression (10) the term with $\ddot{r}$ (in fact this term is negligible and only gives terms of higher order). Hence, at this approximation, we obtain the velocity-dependent potential

$$(11) \quad V \approx -\frac{m}{r} \left[1 + \frac{\dot{r}}{r} \tau + \frac{\dot{r}^2}{r^2} \tau^2 \right],$$

In a first approximation, the delay τ must be equal to r/c (the time that the field uses to go from Mercury to the Sun at the speed of the light) and according with the theories developed in [12, 13], we introduce a new constant g in the delay and hence, $\tau = gr/c$. Introducing this expression of the delay in (12) we have

$$(12) \quad V \approx -\frac{m}{r} \left[1 + g \frac{r\dot{r}}{cr} + g^2 \frac{r^2 \dot{r}^2}{c^2 r^2} \right].$$

On this basis, of this velocity-dependent potential function (12), the gravitational force law is given by substituting the potential function

(12) into equation

$$f = \frac{d}{dt} \left(\frac{\partial V}{\partial \dot{r}} \right) - \frac{\partial V}{\partial r} = -\frac{m}{r^2} \left(1 - \frac{g^2 \dot{r}^2}{c^2} + \frac{2g^2 r \ddot{r}}{c^2} \right).$$

It is easy to see that if we fix $g = \sqrt{3}$, we obtain the same radial force, at first orders, that gives Gerber's potential, see (6). In fact, we have constructed a potential that, varying g , predicts $2g^2\pi m/(Lc^2)$ as non-Newtonian advance of orbital perihelia per revolution. Note that for $g = 1$, it results in a value of $C = 2$, so it predicts only 14.1 arc seconds per century for the precession of Mercury's perihelion, one third of the observed value. The problem of the retarded potential (9) is that it can account for the anomalous precession of the Mercury's perihelion precisely by adjusting a free parameter of the theory. In the following we give a retarded potential which gives an explanation of the anomalous precession of the Mercury's perihelion without adjusting any free parameter of the theory. We will see that this new retarded potential also coincides, at first order, with Gerber's one.

We now consider a small modification of the retarded potential (9), given by

$$(13) \quad V = -\frac{m}{r(t-\tau)} \frac{r(t)}{r(t-\tau)},$$

where the modification consists on dividing the retarded potential (9) by the quotient $r(t-\tau)/r(t)$. And this quotient represents the ratio of the distance between the masses when the potential was "emitted" to distance between the masses at the present instant. We can think that the retarded potential (9) was obtained from the Newtonian potential $V = -m/r(t)$ of the form

$$V = -\frac{m}{r(t) \frac{r(t-\tau)}{r(t)}} = -\frac{m}{r(t-\tau)},$$

and the quotient $r(t-\tau)/r(t)$ is the corrective factor to obtain the retarded distance. This corrective factor is applied because the potential must propagate from the source to the location particle in question. In the same way we can think that the retarded potential (13) is obtained from the Newtonian potential $V = -m/r(t)$ of the form

$$V = -\frac{m}{r(t) \frac{r(t-\tau)}{r(t)} \frac{r(t-\tau)}{r(t)}} = -\frac{m}{r(t-\tau)} \frac{r(t)}{r(t-\tau)},$$

In the same way that in the Neumann's theories [17] we conceive the potential essentially as information being transmitted from place

to place, and assumed a finite speed for the propagation of this information.




FIGURE 1. The retarded position of the test particle.

A particle sends forth a potential, the value of which depends not only on the emitting particle, but on the receiving particle. Therefore, the information must come back from the receiving particle to the emitting particle. Thus we ought to regard an elementary interaction not as a one-way exchange, but as a two-way round-trip exchange. Hence, we must apply the corrective factor twice in the initial potential.

In fact the correct expression of the retarded potential, taking into account that the information must do a two-way round-trip and that $\tau = r(t)/c$, is

$$(14) \quad V = -\frac{m}{r(t - \tau - \frac{r(t-\tau)}{c})}.$$

where $r(t - \tau - r(t - \tau)/c)$ is the distance between the masses when the potential was “emitted” to go from the emitting particle to the receiving particle and come back, see the second graphic of Fig. 1. To find the retarded potential (13) as approximation of the retarded potential (14) we take into account that for a small τ we have that

$$r(t) r(t - \tau - \frac{r(t-\tau)}{c}) \approx (r(t - \tau))^2.$$

Therefore, for a small τ we obtain

$$V = -\frac{m}{r(t - \tau - \frac{r(t-\tau)}{c})} \approx -\frac{m}{r(t - \tau)} \frac{r(t)}{r(t - \tau)}.$$

Hence, the correct retarded potential is (14), but is a functional potential which is difficult to express in a form that is amenable to analysis. Therefore, we use the approximation (13) whose physical interpretation and use is totally justified. In fact, the retarded potential (13) is a generalization of the Gerber's potential. The Gerber's potential is the particular case when the velocity of the test particle is constant, i.e., when $\ddot{r} = 0$. In [16] a physical explanation (albeit in a way that evidently never occurred to Gerber) of the form of the Gerber's potential is given.

Now we are going to see that the retarded potential (13) gives an explanation of the anomalous precession of Mercury's perihelion because coincides, at first order, with the force law associated to Gerber's one. If we develop the retarded potential (13) in powers of τ (up to second order in τ), we obtain

$$(15) \quad V \approx -\frac{m}{r} \left[1 + \frac{2\dot{r}}{r} \tau + \left(\frac{3\dot{r}^2}{r^2} - \frac{\ddot{r}}{r} \right) \tau^2 \right],$$

To develop some easier calculations we can reject, as before, on the right hand side of expression (15) the term with $\ddot{r}$ (in fact this term is negligible and only gives terms of higher order). Hence, at this approximation, we obtain the velocity-dependent potential

$$(16) \quad V \approx -\frac{m}{r} \left[1 + \frac{2\dot{r}}{r} \tau + \frac{3\dot{r}^2}{r^2} \tau^2 \right],$$

In a first approximation, the delay τ must be equal to r/c (the time that the field uses to go from Mercury to the Sun at the speed of the light) according with the theories developed in [12, 13]. Introducing this expression of the delay in (16) we have:

$$(17) \quad V \approx -\frac{m}{r} \left[1 + \frac{2r\dot{r}}{cr} + \frac{3r^2\dot{r}^2}{c^2r^2} \right].$$

On this basis, of this velocity-dependent potential function (17), the gravitational force law is given by substituting the potential function (17) into the equation:

$$f = \frac{d}{dt} \left(\frac{\partial V}{\partial \dot{r}} \right) - \frac{\partial V}{\partial r} = -\frac{m}{r^2} \left(1 - \frac{3\dot{r}^2}{c^2} + \frac{6r\ddot{r}}{c^2} \right).$$

Hence, we obtain (without fixing any parameters) the same radial force, at first orders, that gives Gerber's potential, see (6).

In fact, it is straightforward to see that, at first order, the retarded potential (13) and Gerber's potential coincide. If we develop the retarded potential (13) we have

$$\begin{aligned} V &= -\frac{m}{r(t-\tau)} \frac{r(t)}{r(t-\tau)} = -\frac{m}{r(t) - \dot{r}(t)\tau + \dots} \cdot \frac{r(t)}{r(t) - \dot{r}(t)\tau + \dots} \\ &= \frac{m}{r(t)(1 - \frac{\dot{r}(t)}{r(t)}\tau + \dots)} \cdot \frac{1}{1 - \frac{\dot{r}(t)}{r(t)}\tau + \dots}. \end{aligned}$$

Now substituting the delay $\tau = r/c$ we obtain

$$V = -\frac{m}{r(t)(1 - \frac{\dot{r}(t)}{c} + \dots)} \cdot \frac{1}{1 - \frac{\dot{r}(t)}{c} + \dots}.$$

Therefore, at first order, the retarded potential (13) has the form

$$V = -\frac{m}{r(t) \left(\left(1 - \frac{\dot{r}(t)}{c}\right)^2 + \dots \right)}.$$

4. CONCLUDING REMARKS

Therefore the anomalous precession of the Mercury's perihelion is, in fact, to take into account the second order in the delay of the retarded potential (13) which is an approximation of the correct retarded potential (14). It lacks to see if the prediction for the deflection of electromagnetic waves grazing the Sun using this potential coincide with value given by General Relativity, assuming a plausible application of such potential to the propagation of electromagnetic waves. We hope to give an answer in a future work.

Acknowledgements:

The author would like to thank Prof. M. Grau from Universitat de Lleida for several useful conversations and remarks.

REFERENCES

- [1] H. BEHERA AND P.C. NAIK, *A flat space-time relativistic explanation for the perihelion advance of Mercury*, astro-ph/0306611, preprint, 2003.
- [2] C. CHICONE, *What are the equations of motion of classical physics?*, Can. Appl. Math. Q. **10** (2002), no. 1, 15–32.
- [3] C. CHICONE, S.M. KOPEIKIN, B. MASHHOON AND D. RETZLOFF, *Delay equations and radiation damping*, Phys. Letters **A 285** (2000), 17–16.
- [4] C. CHICONE, *Inertial and slow manifolds for delay equations with small delays*, J. Differential Equations **190** (2003), no. 2, 364–406.
- [5] C. CHICONE, *Inertial flows, slow flows, and combinatorial identities for delay equations*, J. Dynam. Differential Equations **16** (2004), no. 3, 805–831.

- [6] A. EINSTEIN, *Erklärung der Perihelbewegung des Merkur aus der allgemeinen Relativitätstheorie (Explanation of the perihelion motion of mercury from the general theory of relativity)*. Königlich Preuáische Akademie der Wissenschaften, Sitzungsberichte (1915), 831–839.
- [7] A. EINSTEIN, *Die Grundlage der allgemeinen Relativitätstheorie (The foundation of the general theory of relativity)*. Ann. Phys. **49** (1916), 769–822.
- [8] F.G. HOLZMULLER, Z. Math. Phys. **15**, 69 (1870).
- [9] P. GERBER, *Die räumliche und zeitliche Ausbreitung der Gravitation (Space and temporary propagation of gravitation)*. Z. Math. Phys. **43** (1898), 93–104.
- [10] P. GERBER, *Die Fortpflanzungsgeschwindigkeit der Gravitation (The propagation-velocity of gravitation)*. Ann. Phys. **52** (1917), 415–444.
- [11] J. GINÉ, *On the classical descriptions of the quantum phenomena in the harmonic oscillator and in a charged particle under the coulomb force*, Chaos Solitons Fractals **26** (2005), 1259–1266.
- [12] J. GINÉ, *On the origin of quantum mechanics*, physics/0505181, preprint, Universitat de Lleida, 2005.
- [13] J. GINÉ, *On the origin of gravitational quantization: the Titius–Bode law*, physics/0507072, preprint, Universitat de Lleida, 2005.
- [14] P.S. DE LAPLACE, *Traite de Mécanique Céleste* tome chapitre vii, §22, (1805).
- [15] B. MASHHOON, *Gravitoelectromagnetism: A Brief Review*, gr-qc/0311030, preprint, 2003.
- [16] MATHPAGES www.mathpages.com/home/kmath527/kmath527.htm
- [17] C. NEUMANN, *Principles of Electrodynamics*, Tübingen, 1868.
- [18] S. OPPENHEIM Ann. Phys. **52**, 415 (1917); **53**, 31 & 163 (1917); **54**, 38 (1917).
- [19] S. OPPENHEIM, *Kritik des Newtonschen Gravitationsgesetzes*, Encyklopädie der mathematischen Wissenschaften, vol. 6.2.2 pag. 80–158, Teuner, Leipzig.
- [20] H. POINCARÉ, *Mémoire sur les courbes définies par les équations différentielles*. Journal de Mathématiques **37** (1881), 375–422; **8** (1882), 251–296; Oeuvres de Henri Poincaré, vol. I, Gauthier-Villars, Paris, (1951), pp. 3–84.
- [21] C.K. RAJU, *The electrodynamic 2-body problem and the origin of quantum mechanics*, Foundations of Physics **34** (2004), 937–962.
- [22] C.K. RAJU, *Time: towards a consistent theory*, Kluwer academic, Dordrecht, 1994.
- [23] F. TISSERAND, Compt. Rend. Acad. Sci. (Paris) **75**, 760 (1872); **110**, 313 (1890).
- [24] E. WHITTAKER, *A History of the theories of ether and electricity*, vol. 2, Thomas Nelson, Edinburgh, 1953.

DEPARTAMENT DE MATEMÀTICA, UNIVERSITAT DE LLEIDA, AV. JAUME II,
69. 25001 LLEIDA, SPAIN

E-mail address: gine@eps.udl.es