

Moody's Correlated Binomial Default Distributions for Inhomogeneous Portfolios

S.Mori ††, K Kitsukawa ‡‡ and M Hisakado ¶¶§

† Department of Physics, School of Science, Kitasato University, Kitasato 1-15-1 , Sagamihara, Kanagawa 228-8555, Japan

¶ Graduate School of Media and Governance, Keio University, Endo 5322, Fujisawa, Kanagawa 252-8520, Japan

‡ Standard & Poor's, Marunouchi 1-6-5, Chiyoda-ku, Tokyo 100-0005, Japan

Abstract. This paper generalizes Moody's correlated binomial default distribution for homogeneous (exchangeable) credit portfolio, which is introduced by Witt, to the case of inhomogeneous portfolios. As inhomogeneous portfolios, we consider two cases. In the first case, we treat a portfolio whose assets have uniform default correlation and non-uniform default probabilities. We obtain the default probability distribution and study the effect of the inhomogeneity on it. The second case corresponds to a portfolio with inhomogeneous default correlation. Assets are categorized in several different sectors and the inter-sector and intra-sector correlations are not the same. We construct the joint default probabilities and obtain the default probability distribution. We show that as the number of assets in each sector decreases, inter-sector correlation becomes more important than the intra-sector one. We study the maximum values of the inter-sector default correlation. In the modeling of portfolio credit risk, it is possible to incorporate realistic magnitude of inter-sector default correlations. Our generalization method can be applied to any correlated binomial default distribution model which has explicit relations on the conditional default probabilities or conditional default correlations, e.g. Credit Risk⁺. We also compare some popular CDO pricing models from the viewpoint of the range of the implied tranche correlation.

Submitted to: *Quantitative Finance: Revised*

† mori@sci.kitasato-u.ac.jp

‡ kj198276@sfc.keio.ac.jp

§ masato_hisakado@standardpoors.com

1. Introduction

The modeling of portfolio credit risk and default correlation are hot topics and pose entirely new problems [1, 2, 3, 4, 5]. CDOs are financial innovations to securitize portfolios of defaultable assets. Many probabilistic models have been studied in order to price CDO tranches [6, 7, 8, 9, 10, 11, 12, 3, 13, 14, 15]. Most of them are implemented with Monte Carlo simulations and as the number of names in portfolio increases, the computational time increases. The Factor approach uses small number of latent factors that induce the default dependency [11]. Conditionally on the latent variables values, default events are independent. It becomes easy to calculate the loss (default) distribution function. Along this line, some semi-explicit expressions of most relevant quantities were obtained [16].

On the other hand, correlated binomial models were also proposed to describe the default dependency structures. The first one is a one-factor exchangeable version of CreditRisk⁺ [17, 18, 19]. The aggregate loss distribution function is given by the beta-binomial distribution (BBD). The second one is the Moody's correlated binomial default distribution model, which was introduced by Witt [20] and we call it as MCB model for abbreviation. The authors also consider the applicability of the long-range Ising model [21, 22]. They use Bernoulli random variables $X_i (i = 1, \dots, N)$ which take values 0, 1 depending on whether the i -th asset described by X_i is not defaulted or defaulted. The default probability p is defined by the average number of defaulted assets n per an asset as $p = \frac{\langle n \rangle}{N}$. Here $\langle \ \rangle$ means the expectation value. In the MCB model, the default correlation between the assets is set to be constant ρ irrespective of the number of defaults [20]. From these conditions, the joint default probabilities and default distribution function $P_N(n)$ were calculated explicitly. One-factor exchangeable version of CreditRisk⁺, which we call BBD model hereafter, the default correlation behaves as $\rho_n = \frac{\rho}{1+n\rho}$ under n defaults [23]. As default number increases, it decreases. $P_N(n)$ s of the MCB and BBD look similar to that of the Gaussian copula model [20, 23].

The advantage of these correlated binomial models comes from the fact that they are easier to evaluate than other more refined models. If a probabilistic model is implemented by Monte Carlo simulation, the evaluation of the price of these derivatives consumes much computer time and the inverse process to obtain the model parameters becomes tedious work. With the above correlated binomial models, one can estimate the model parameters from the derivative premiums more easily. However, these models are formulated only for homogeneous portfolio, where the assets are exchangeable and they have the same default probability p and default correlation ρ . Generalization to more realistic inhomogeneous portfolios where assets have different default probabilities and different default correlations should be done.

In this paper, we show how to generalize the Moody's correlated binomial default distribution (MCB) model to two types of inhomogeneous portfolios. Our generalization method can be applied to other correlated binomial models by changing the condition on ρ_n . We obtain the default probability function $P_N(n)$ and examine the dependence

of the expected loss rates of the tranches on the inhomogeneities. With the proposed model, we also estimate the implied values of the default correlation. Comparison of the range of the correlation skews with those of the Gaussian copula model and BBD model are also performed.

About the organization of the paper, we start with a short review of the Moody's correlated binomial default distribution (MCB) model in Section 2. The dependence of ρ on the number of defaults n is compared with the BBD and Gaussian copula. Section 3 is the main part of the paper. We show how to couple multiple MCB models as a portfolio credit risk model for inhomogeneous portfolio. In the first subsection, we couple two Bernoulli random variables X, Y and recall on the limit of the correlation ρ_{xy} between them. In the next subsection, we couple a MCB model of N assets X_1, X_2, \dots, X_N with a random variable Y and we study the maximum value of the correlation between X_i and Y . Then we couple two MCB models with N and M assets. Choosing the model parameters properly, we construct a MCB model for an inhomogeneous default probability case and obtain the default distribution function. The last subsection is devoted to an inhomogeneous default correlation case. Assets are categorized in different sectors and inter-sector and intra-sector default correlations are not the same. We consider a portfolio with K sectors and k -th sector contains N_k assets. Within each sector, the portfolio is homogeneous and it has parameters as p_k and ρ_k . The inter-sector default correlations are not the same and they depends on the choice of sector pairs. We construct the joint default probabilities and the default probability function $P_N(n)$ for the portfolio explicitly. In Section 4, using the above results, we estimate the implied default correlation for each tranche from a CDO's market quotes. We compare the range of the correlation skews of MCB, BBD and the Gaussian copula models. We conclude with some remarks and future problems.

2. Moody's correlated binomial default distribution




Figure 1. Homogeneous portfolio with N assets. The assets are exchangeable and the default probability is p and the default correlation is ρ . The state of i -th asset is described by a Bernoulli random variable X_i .

We review the definitions and some properties of Moody's correlated binomial default distribution (MCB) model. We consider a homogeneous portfolio, which

is composed of exchangeable N assets. Here the term “homogeneous” means that the constituent assets are exchangeable and their default probabilities and default correlations are uniform. We denote them as p and ρ . Bernoulli random variables X_i show the states of the i -th assets. $X_i = 1$ means that the asset is defaulted and non default state is represented as $X_i = 0$. The joint default probabilities is denoted as

$$P(x_1, x_2, \dots, x_N) = \text{Prob}(X_1 = x_1, X_2 = x_2, \dots, X_N = x_N). \quad (1)$$

In order to determine $P(x_1, x_2, \dots, x_N)$, we need $2^N - 1$ conditions for them. Here 2^N correspond to the number of possible configurations and -1 comes from the overall normalization condition for the joint probabilities. From the assumption of the homogeneity for the portfolio, the number of degrees of freedom of the joint probabilities are reduced. The probability for n defaults and $N - n$ non-defaults is the same for any configuration (x_1, x_2, \dots, x_N) with $\sum_{i=1}^N x_i = n$. The number of defaults n is ranged from 0 to N and considering the overall normalization condition, remaining degrees of freedom are N .

In MCB model, the conditional default probabilities are introduced. We denote p_n as the default probability for any assets under the condition that any other n assets of the portfolio are defaulted. To exemplify the situation concretely, we take the n assets as the first n of N assets, we denote them with n' as $n' = 1, 2, \dots, n$. The condition that they are defaulted is written concisely as $\prod_{n'=1}^n X_{n'} = 1$. The conditional default probability for $n + 1$ -th assets under the condition of n defaults can then be written as

$$p_n = \langle X_{n+1} | \prod_{n'=1}^n X_{n'} = 1 \rangle. \quad (2)$$

Here, $\langle A|C \rangle$ means the expectation value of random variable A under condition C is satisfied. X_n takes 1 for n -th asset default, $\langle X_n|C \rangle$ corresponds to its default probability with condition C . Of course, any asset from $k = n + 1, \dots, N$ can be chosen in the evaluation of the expectation value for p_n with the condition that $\prod_{n'=1}^n X_{n'} = 1$. p_0 is nothing but the default probability p .

N independent conditional default probabilities $p_n (i = 0, \dots, N - 1)$ are determined by the following condition on the default correlations.

$$\text{Cor}(X_{n+1}, X_{n+2} | \prod_{n'=1}^n X_{n'} = 1) = \rho. \quad (3)$$

Here, $\text{Cor}(X, Y|C)$ is defined as

$$\text{Cor}(X, Y|C) = \frac{\langle XY|C \rangle - \langle X|C \rangle \langle Y|C \rangle}{\sqrt{\langle X|C \rangle (1 - \langle X|C \rangle) \langle Y|C \rangle (1 - \langle Y|C \rangle)}}. \quad (4)$$

The conditions on the default correlations give us the following recursion relations for p_n as

$$p_{n+1} = p_n + (1 - p_n)\rho. \quad (5)$$

These recursion relations can be solved to give p_n as

$$p_n = 1 - (1 - p)(1 - \rho)^n. \quad (6)$$

p_n increases with n and $p_n \rightarrow 1$ as $n \rightarrow \infty$ for $\rho > 0$.

From these conditional default probabilities $p_n (n = 0, \dots, N-1)$, the joint default probabilities for the configuration $\vec{x} = (x_1, x_2, \dots, x_N)$ are given as

$$P(\vec{x}) = P(x_1, x_2, \dots, x_N) = \langle \prod_{n=1}^N X_n^{x_n} (1 - X_n)^{1-x_n} \rangle. \quad (7)$$

The normalization condition for $P(\vec{x})$ is guaranteed by the following decomposition of unity.

$$1 = \langle 1 \rangle = \langle \prod_{n=1}^N \{X_n + (1 - X_n)\} \rangle = \prod_{j=1}^N \left[\sum_{x_j=0}^1 \right] \langle \prod_{n=1}^N X_n^{x_n} (1 - X_n)^{1-x_n} \rangle \quad (8)$$

The probability for n defaults is

$$\begin{aligned} P_N(n) &= {}_N C_n \times P(1, 1, \dots, 1, 0, \dots, 0) = {}_N C_n \langle \prod_{i=1}^n X_i \prod_{i=n+1}^N (1 - X_i) \rangle \\ &= {}_N C_n \sum_{k=0}^{N-n} {}_{N-n} C_k (-1)^k \left(\prod_{n'=0}^{n+k-1} p_{n'} \right). \end{aligned} \quad (9)$$

We modify the above MCB model as follows. In the MCB model, the default correlation is set to be constant irrespective of the number of default (see(3)). We change the condition as

$$\text{Cor}(X_{n+1}, X_{n+2} | \prod_{n'=1}^n X_{n'} = 1) = \rho \exp(-n\lambda). \quad (10)$$

Here, we introduce a parameter $\lambda > 0$ and the default correlation under n defaults decay as $\exp(-n\lambda)$. If we set $\lambda = 0$, the modified model reduces to the original MCB model.

There are two motivation for the modification. The first one is that it is mathematically necessary to couple multiple MCB models. About the mechanism, we discuss it in greater detail in the next section. Here, we only comment on the limit value of p_n as $n \rightarrow \infty$. The modification changes the recursive relation for p_n to

$$p_{n+1} = p_n + (1 - p_n) \rho \exp(-n\lambda). \quad (11)$$

p_n is calculated as

$$p_n = 1 - (1 - p) \prod_{n'=0}^{n-1} (1 - \rho_{n'}). \quad (12)$$

Here ρ_n is defined as $\rho_n = \rho \exp(-n\lambda)$.

p_n increases with n , however the increase is reduced by the decay of the correlation with n . About the limit value of p_n with $n \rightarrow \infty$ is roughly estimated as

$$p_\infty = \lim_{n \rightarrow \infty} p_n = 1 - (1 - p)(1 - \rho) \exp\left(-\rho \frac{e^{-\lambda}}{1 - e^{-\lambda}}\right). \quad (13)$$

For $\lambda = 0$, $p_\infty = 1$ and $p_\infty = p + (1 - p)\rho = p_1$ for $\lambda = \infty$. In Figure 2, we show the enumerated data for p_∞ and the results from eq.(13). As λ increases, p_∞ decreases and the λ dependence is well described by eq.(13).




Figure 2. p_∞ for $0 \leq \lambda \leq 0.3$ with $p = \rho = 0.1$. The solid line comes from eq.(13). The numerically enumerated data are shown with + symbol.

The second motivation is that popular CDO pricing models have decaying correlation with n . The BBD model's ρ_n is given as [23]

$$\rho_n = \frac{\rho}{1 + n\rho}. \quad (14)$$

The Gaussian copula model, we do not have the explicit form for ρ_n . From its aggregate loss distribution function, it is possible to estimate them. In Figure 3, we show ρ_n for the MCB, BBD and Gaussian copula model. We set $p = \rho = 0.1$ and $N = 30$. The Gaussian copula's ρ_n does not show monotonic dependence on n . After a small peak, it decays to zero. In order to mimic the Gaussian copula model within the framework of correlated binomial model, such a dependence should be incorporated in the assumption on ρ_n .

Figure 4 depicts p_n in the same setting. With the same p and ρ , all p_n curves pass through $p_0 = p$ at $n = 0$ and $p_1 = p + (1 - p)\rho$ at $n = 1$. After $n > 1$, the behaviors of p_n depend on the models' definitions on ρ_n . p_n saturates to about 0.4 for MCB($\lambda = 0.3$), which means that large scale avalanche does not occur and the loss distribution has a short tail. MCB with $\lambda = 0.0, -0.05$, Gaussian copula model, their p_n saturates to 1. The behaviors are reflected in the fat and long tails in their loss distribution.

Figure 5 shows the semi-log plot of $P_{30}(n)$ for the MCB, BBD and Gaussian copula model. We also plot the binomial distribution $\text{Bin}(30, 0.1)$. The default correlation shifts the peak of the binomial distribution to $n = 0$ and $P_{30}(n)$ comes to have a long tail. MCB, BBD and Gaussian copula have almost the same bulk shape. In particular, about MCB, even if we change λ , $P_{30}(n)$ have almost the same shape for $n \leq 15$. We think that the bulk shape of $P_N(n)$ is mainly determined by p_n with small n . p_n s with large n comes from very rare events $\prod_{n'=1}^n X_{n'} = 1$ and contain informations about the tails of the distributions. They do not affect the bulk part so much.




Figure 3. ρ_n for MCB (blue,green,red),BBD (violet) and Gaussian copula model (aqua). We set $p = \rho = 0.1$ and $N = 50$.




Figure 4. p_n for MCB (blue,green,red),BBD (violet) and Gaussian copula model (aqua). We set $p = \rho = 0.1$ and $N = 50$.

About their tails, there are differences. One sorts the models in the order of thinnest tail to fattest tail, we have

$$\text{MCB}(\lambda = 0.3) < \text{BBD} < \text{Gaussian copula} < \text{MCB}(0.0) < \text{MCB}(-0.05).$$

$\text{MCB}(\lambda = 0)$ has almost the same shape with the Gaussian copula. However, it has a small peak near $n = 30$. $\text{MCB}(\lambda = -0.05)$ has also a fat tail and a large peak at $n = 30$. The tail of $\text{MCB}(\lambda = 0.3)$ is short as compare with other models. We can understand these behaviors from the behaviors of p_n .




Figure 5. Loss Distribution $P_{30}(n)$ for $p = 0.1$ and $\rho = 0.1$. $\lambda = 0.0, 0.3$ and $p = \rho = 0.1$. We also plot the profile for the binomial distribution $b(N, p)$. $\Delta P_N(n)$ is the difference of $P_N(n)$ with $\lambda = 0.3$ and $\lambda = 0.0$.

We also note another role of the damping parameter λ . In the calculation $P_N(n)$, there are many cancellations $\sum_{k=0}^{N-n} \binom{N-n}{k} (-1)^k$ in (9) from the decomposition of $\prod_{i=n+1}^N (1 - X_i)$. This causes numerical error in the evaluation and it is difficult to get $P_N(n)$ for $N \geq 40$, even if we use long double precision variable in the numerical implementation. When we set $\lambda = 0.3$, the numerical error diminishes much and we can obtain $P_N(n)$ even for $N = 100$. This point is important, when one uses the MCB

model for the actual CDO analysis, which has at least 50 assets. In addition, with $\lambda > 0$, we can take ρ to be negatively large enough. In S&P's data, the negative default correlation of 0.1% or so has been reported [24]. We think that this point is also the advantage of the modified model.

Hereafter, we mainly focus on the generalization of the MCB model. However, the same method and reasoning should be applicable to other correlated binomial models with any assumption on ρ_n . If we use ρ_n as in (14), we have Beta-Binomial default distribution models for inhomogeneous portfolios.

3. Generalization to Inhomogeneous Portfolios

In this section we couple multiple MCB models and construct the joint default probabilities and $P_N(n)$ for inhomogeneous portfolios. In addition, we show that the inter-sector default correlation can be set to be large enough by choosing λ and other parameters. We think that it is possible to use the model as a model for portfolio credit risk.

3.1. Coupling of X and Y : 1+1 MCB model

Before proceeding to the coupling of multiple MCB models, we recall some results for the coupling of two random variables X and Y . The default probability is p_x (p_y) for X (respectively for Y) and the default correlation between them is ρ_{xy} ,

$$\text{Corr}(X, Y) = \rho_{xy}. \quad (15)$$

Like in the MCB model, we introduce the conditional default probabilities as

$$\begin{aligned} p_0 &= p_x = \langle X \rangle & \text{and} & & p_1 &= \langle X|Y = 1 \rangle. \\ q_0 &= p_y = \langle Y \rangle & \text{and} & & q_1 &= \langle Y|X = 1 \rangle. \end{aligned} \quad (16)$$

From the default correlation ρ_{xy} , p_1 and q_1 are calculated as

$$\begin{aligned} p_1 &= p_0 + (1 - p_0) \sqrt{\frac{p_0(1 - q_0)}{(1 - p_0)q_0}} \rho_{xy} \\ q_1 &= q_0 + (1 - q_0) \sqrt{\frac{q_0(1 - p_0)}{(1 - q_0)p_0}} \rho_{xy}. \end{aligned} \quad (17)$$

In the symmetric (homogeneous) case $p_x = p_y$, the equality $p_1 = q_1$ holds and they are given as

$$p_1 = q_1 = p_x + (1 - p_x) \rho_{xy}.$$

The correlation ρ_{xy} can be set to be 1 and in the limit $p_1 = q_1 \rightarrow 1$. The maximum value of ρ_{xy} is 1 in the symmetric case. On the contrary, in the asymmetric case ($p_x \neq p_y$), ρ_{xy} cannot be set to be 1. The maximum value of ρ_{xy} is determined by the condition that $p_1 \leq 1$ and $q_1 \leq 1$. From these conditions, we derive the following conditions on ρ_{xy} as

$$\rho_{xy} \leq \sqrt{\frac{q_0(1 - p_0)}{(1 - q_0)p_0}} \quad \text{and} \quad \rho_{xy} \leq \sqrt{\frac{p_0(1 - q_0)}{(1 - p_0)q_0}}. \quad (18)$$

We introduce a asymmetric parameter r as

$$r = \frac{p_y}{p_x} \quad (19)$$

and a function $f(r, p)$ as

$$f(r, p) = \sqrt{\frac{(1-p)r}{1-rm}}. \quad (20)$$

The maximum value of ρ_{xy} is then given as

$$\text{Max}(\rho_{xy}) = \text{Min}(f(r, p_0), f(r, p_0)^{-1}). \quad (21)$$

Here $\text{Max}(\rho_{xy})$ represents the maximum value of ρ_{xy} and $\text{Min}(A, B)$ means that the smaller value of A and B is taken. Figure 6 shows $\text{Max}(\rho_{xy})$ as a function of the




Figure 6. $\text{Max}(\rho_{xy})$ for $p_x = 0.1, 0.5$ and $0.2 \leq r \leq 5.0$.

asymmetry (inhomogeneity) parameter r . We show two curves, the red one for $p_x = 0.1$ and the green one for $p_x = 0.5$. As the inhomogeneity r increases, that is r departs from $r = 1$, $\text{Max}(\rho_{xy})$ decreases. For fixed r , as p_x becomes large $\text{Max}(\rho_{xy})$ becomes small. The reason is that the condition $p_1 \leq 1$ becomes more difficult to be satisfied as $p_x = p_0$ increase. p_1 is a monotonous increasing function of p_0 . In the previous section, the conditional default probability p_n becomes smaller as we set λ larger. When we set λ large, we show that it is possible to couple multiple MCB models with large default correlation.

3.2. $N + 1$ MCB model

As the second step, we couple a N assets MCB model with one two-valued random variable Y . We introduce N random variables $X_n (n = 1, \dots, N)$ and the default probability and the default correlation for them is $\langle X_n \rangle = p_x$ and ρ . The default probability for Y is $\langle Y \rangle = p_y$ and the default correlation between Y and X_n is written

as ρ_{xy} . We assume homogeneity for the N assets MCB model, the default correlation between X_n and Y is independent of the asset index i . As in the previous cases, we introduce conditional default probabilities as

$$\begin{aligned} p_{n,0} &= \langle X_{n+1} | \prod_{n'=1}^n X_{n'} = 1 \rangle \quad \text{and} \quad p_{n,1} = \langle X_{n+1} | \prod_{n'=1}^n X_{n'} \times Y = 1 \rangle \\ q_n &= \langle Y | \prod_{n'=1}^n X_{n'} = 1 \rangle. \end{aligned} \quad (22)$$

The joint default probabilities $P(\vec{x}, y) = P(\vec{X} = \vec{x}, y)$ are calculated by decomposing the following expression with these conditional default probabilities

$$P(\vec{x}, y) = \langle \prod_{n=1}^N X_n^{x_n} (1 - X_n)^{1-x_n} \times Y^y (1 - Y)^{1-y} \rangle \quad (23)$$

The joint default probabilities with the condition $Y = 1$ are

$$P(\vec{x} | Y = 1) = \langle \prod_{n=1}^N X_n^{x_n} (1 - X_n)^{1-x_n} \rangle / p_y. \quad (24)$$

The joint probabilities with the condition $Y = 0$ are obtained by the following relations.

$$P(\vec{x} | Y = 0)(1 - p_y) + P(\vec{x} | Y = 1)p_y = P(\vec{x}) = \langle \prod_{n=1}^N X_n^{x_n} (1 - X_n)^{1-x_n} \rangle \quad (25)$$

The explicit form for $P(\vec{x} | y = 0)$ is

$$P(\vec{x} | Y = 0) = \frac{P(\vec{x}) - P(\vec{x} | Y = 1)p_y}{1 - p_y}. \quad (26)$$

The probability for n defaults is given as in eq.(9)

$$P_N(n) = {}_N C_n \sum_{k=0}^{N-n} {}_{N-n} C_k (-1)^k \left(\prod_{n'=0}^{n+k-1} p_{n',0} \right). \quad (27)$$

The probability for n defaults with $Y = 1$ is

$$P_N(n | Y = 1) = {}_N C_n \sum_{k=0}^{N-n} {}_{N-n} C_k (-1)^k \left(\prod_{n'=0}^{n+k-1} p_{n',1} \right). \quad (28)$$

Using the same argument as for the joint probabilities with $Y = 0$, $P_N(n | Y = 0)$ is written as

$$P_N(n | Y = 0) = \frac{P_N(n) - P_N(n | 1)p_y}{1 - p_y}. \quad (29)$$

About the conditional default probabilities $p_{n,0}$, we impose the same conditions as with the homogeneous N assets MCB model.

$$\text{Cor}(X_{n+1}, X_{n+2} | \prod_{n'=1}^n X_{n'} = 1) = \rho \exp(-n\lambda) \quad (30)$$

The same recursive relation (11) for $p_{n,0}$ is obtained and the $p_{n,0}$ is given by

$$p_{n,0} = 1 - (1 - p_x) \prod_{n'=0}^{n-1} (1 - \rho_{n'}). \quad (31)$$

Here ρ_n is defined as before. About the condition on $p_{n,1}$ and q_n , there are two possible ways. The only way to realize a strong correlation ρ_{xy} between X_{n+1} and Y is

$$\text{Corr}(X_{n+1}, Y | \prod_{n'=1}^n X_{n'}) = \rho_{xy} e^{-n\lambda}. \quad (32)$$

In the case, the relations for $p_{n,1}$ and q_n are

$$p_{n,1}q_n = q_{n+1}p_{n,0} = p_{n,0}q_n + \rho_{xy} \sqrt{p_{n,0}(1-p_{n,0})q_n(1-q_n)} \exp(-n\lambda). \quad (33)$$

The recursive relations are

$$p_{n,1} = p_{n,0} + \rho_{xy} \exp(-n\lambda)(1-p_{n,0}) \sqrt{\frac{p_{n,0}(1-q_n)}{(1-p_{n,0})q_n}} \quad (34)$$

$$q_{n+1} = q_n + \rho_{xy} \exp(-n\lambda)(1-q_n) \sqrt{\frac{(1-p_{n,0})q_n}{p_{n,0}(1-q_n)}}. \quad (35)$$

If we set $p_y = p_x$ and $\rho_{xy} = \rho$, these relation reduces to $p_{n,1} = p_{n+1,0}$ and $q_n = p_{n,0}$ and this coupled model is nothing but the $(N+1)$ assets MCB model.




Figure 7. Plot of $\text{Max}(\rho_{xy})/\rho$ vs $r = \frac{p_y}{p_x}$. $N = 30, \rho = 0.1, p_x = 0.1$. The solid line $\lambda = 0.0$ and the dotted line $\lambda = 0.3$

We write the ratio $\frac{q_n}{p_{n,0}}$ as r_n and the condition that $p_{n,1} \leq 1$ and $q_{n+1} \leq 1$ are summarized as

$$\rho_{xy} \leq \exp(n\lambda) \times \text{Min}(f(r_n, p_{n,0}), f(r_n, p_{n,0})^{-1}). \quad (36)$$

$\text{Min}(f(r_n, p_n), f(r_n, p_n)^{-1})$ is nothing but the condition for $\text{Max}(\rho_{xy})$ of the two random variables X, Y with default probabilities $(p_x, p_y) = (p_{n,0}, r_n p_{n,0})$ (see eq.(21)). As we

have explained before, $\text{Min}(f(r_n, p_n), f(r_n, p_n)^{-1})$ takes maximum value 1 at $r_n = 1$ for any value of p_n . It also decrease with the increase of p_n for fixed r_n .

The necessary condition that the model to be self-consistent is that $p_{n,1} \leq 1$ and $q_n \leq 1$ for all n . We discuss $\lambda = 0$ and $\lambda > 0$ cases separately.

- $\lambda = 0.0$ case : $p_{n,0}$ increases with n and $p_{n,0} \rightarrow 1$ as $n \rightarrow \infty$. The range of $p_{n,0}$ is $[p_x, p_{N-1}] \simeq [p_x, 1]$ and it is difficult to choose p_y such that $r_n \simeq 1$ for all n . If r_n departs much from 1 for some n , $\text{Max}(\rho_{xy})$ decreases. The choice $p_x = p_y$ and $\rho_{xy} = \rho$ is possible, we anticipate that $\text{Max}(\rho_{xy})$ decreases from ρ as p_y departs from p_x .
- $\lambda > 0$ case : The limit value of p_n becomes small (see eq.(13)) and the range of p_n is narrow as compared with the $\lambda = 0.0$ case. It may be possible to choose p_y such that the asymmetric parameter r_n is small for small n . In addition, for large n , it is easy to satisfy the condition (36) because of the prefactor $\exp(n\lambda)$. We think that $\text{Max}(\rho_{xy})$ is large in this case.

We have checked numerically the value of the joint probabilities for all configurations (\vec{x}, y) with $N = 30$ and $p_x = \rho = 0.1$. In figure 7, we show the data of $\text{Max}(\rho_{xy})/\rho$ for the case $\lambda = 0.0$ (solid line) and $\lambda = 0.3$ (dotted line). $\lambda = 0.0$ case, $\text{Max}(\rho_{xy}) \simeq \rho$ near $r = 1$ and as r departs from 1, $\text{Max}(\rho_{xy})$ decreases from ρ . The data for $\lambda = 0.3$ case show that it is possible to set $\text{Max}(\rho_{xy})$ large if we use large r . We can choose ρ_{xy} as strong as several times of ρ .

We also point out that the above $N + 1$ MCB model can be used to describe a credit portfolio where one obligor has great exposure. Such an obligor is described by Y and other obligors are by X_n . If p_y is quite different from p_x , we can couple Y and X_n with strong $\rho_{xy} \simeq \rho_x$ by choosing λ large enough.

3.3. $N + M$ MCB Model : Coupled MCB model




Figure 8. Portfolio of $N + M$ MCB model. The first (second) sector has parameters $N(M), p_x (p_y)$ and $\rho_x (\rho_y)$. The inter-sector default correlation is ρ_{xy}

We consider a portfolio with two sectors. The first sector has N assets and the second has M assets. To construct the joint default probabilities for the portfolio, we try to couple two MCB models. The former model's N assets are described by $X_n (n = 1, \dots, N)$ and the states of the latter model's assets are described by $Y_m (m = 1, \dots, M)$. The default probability and the default correlation in each sector are (p_x, ρ_x) and (p_y, ρ_y) . The default correlation between the assets in different sectors is denoted as ρ_{xy} (see Figure 8).

Introducing the conditional default probabilities $p_{n,m}$ and $q_{n,m}$ as

$$p_{n,m} = \langle X_{n+1} | \prod_{n'=1}^n X_{n'} \prod_{m'=1}^m Y_{m'} = 1 \rangle, \quad p_{0,0} = p_x \quad (37)$$

$$q_{n,m} = \langle Y_{m+1} | \prod_{n'=1}^n X_{n'} \prod_{m'=1}^m Y_{m'} = 1 \rangle, \quad q_{0,0} = p_y, \quad (38)$$

we impose the following conditions on $p_{n,0}$ and $q_{0,m}$

$$\text{Cor}(X_{n+1}X_{n+2} | \prod_{n'=1}^n X_{n'} = 1) = \rho_x \exp(-n\lambda_x) \quad (39)$$

$$\text{Cor}(Y_{m+1}Y_{m+2} | \prod_{m'=1}^m Y_{m'} = 1) = \rho_y \exp(-m\lambda_y). \quad (40)$$

The recursive relations for $p_{n,0}$ and $q_{0,m}$ are

$$p_{n+1,0} = p_{n,0} + \rho_x \exp(-n\lambda_x)(1 - p_{n,0}) \quad (41)$$

$$q_{0,m+1} = q_{0,m} + \rho_y \exp(-m\lambda_y)(1 - q_{0,m}). \quad (42)$$

Their solutions are, by denoting $\rho_{x,n} = \rho_x \exp(-n\lambda_x)$ and $\rho_{y,m} = \rho_y \exp(-m\lambda_y)$,

$$p_{n,0} = 1 - (1 - p_x) \prod_{n'=0}^{n-1} (1 - \rho_{x,n'}) \quad (43)$$

$$q_{0,m} = 1 - (1 - p_y) \prod_{m'=0}^{m-1} (1 - \rho_{y,m'}). \quad (44)$$

About the inter-sector correlation, we impose the next conditions on $p_{n,m}$ and $q_{n,m}$, which is natural generalization of $N + 1$ case (see eq.(32)).

$$\text{Cor}(X_{n+1}Y_{m+1} | \prod_{n'=1}^n X_{n'} \prod_{m'=1}^m Y_{m'} = 1) = \rho_{xy} e^{-(n\lambda_x + m\lambda_y)}. \quad (45)$$

We obtain the following recursive relations,

$$\begin{aligned} p_{n,m+1}q_{n,m} &= q_{n+1,m}p_{n,m} \\ &= p_{n,m}q_{n,m} + \rho_{xy} e^{-(n\lambda_x + m\lambda_y)} \sqrt{p_{n,m}(1 - p_{n,m})q_{n,m}(1 - q_{n,m})}. \end{aligned} \quad (46)$$

Using these relations, we are able to calculate $p_{n,m}$ and $q_{n,m}$ iteratively starting from $p_{n,0}$ and $q_{0,m}$.

The joint default probability for the portfolio configuration (\vec{x}, \vec{y}) is calculated by decomposing the following expression with $p_{n,m}$ and $q_{n,m}$

$$\begin{aligned} P(\vec{x}, \vec{y}) &= P(x_1, x_2, \dots, x_N, y_1, y_2, \dots, y_M) \\ &= \langle \prod_{n=1}^N X_n^{x_n} (1 - X_n)^{1-x_n} \prod_{m=1}^M Y_m^{y_m} (1 - Y_m)^{1-y_m} \rangle. \end{aligned} \quad (47)$$

In particular, the probability for n, m defaults in each sector, which is denoted as $P_{N,M}(n, m)$, is

$$\begin{aligned}
P_{N,M}(n, m) &= {}_N C_m \times {}_M C_m \times P(1, 1, \dots, 1, 0, \dots, 0, 1, 1, \dots, 1, 0, \dots, 0) \\
&= {}_N C_m \times {}_M C_m \times \left\langle \prod_{n'=1}^n X_{n'} \prod_{k=n+1}^N (1 - X_k) \prod_{m'=1}^m Y_{m'} \prod_{l=m+1}^M (1 - Y_l) \right\rangle \\
&= {}_N C_m \times {}_M C_m \times \sum_{k=0}^{N-n} \sum_{l=0}^{M-m} (-1)^{k+l} {}_{N-n} C_k \times {}_{M-m} C_l \\
&\times \left\langle \prod_{n'=1}^{n+k} X_{n'} \prod_{m'=1}^{m+l} Y_{m'} \right\rangle. \tag{48}
\end{aligned}$$

$P_{N+M}(n)$ is easily calculated from $P_{N,M}(n, m)$ as

$$P_{N+M}(n) = \sum_{n'=0}^n P_{N,M}(n', n - n'). \tag{49}$$

In decomposing $\langle \prod_{n=1}^k X_n \prod_{m=1}^l Y_m \rangle$, one can do it in any order. The independence of the order of the decomposition of $\langle \prod_{n=1}^k X_n \prod_{m=1}^l Y_m \rangle$ is guaranteed by (45) and (46). We decompose it as

$$\left\langle \prod_{n=1}^k X_n \prod_{m=1}^l Y_m \right\rangle = \prod_{n=0}^{k-1} p_{n,0} \times \prod_{m=0}^{l-1} q_{k,m}. \tag{50}$$

About the maximum value of ρ_{xy} , it is necessary to check all values of the joint probabilities. However, $N + M$ model is reduced to $N + 1$ or $1 + M$ model by choosing $M = 1$ or $N = 1$ respectively. From the discussions and the results in the previous subsection for $N + 1$ model, we can anticipate as follows.

- (i) $\lambda_x = \lambda_y = 0$ Case : If $p_x = p_y$ and $\rho_x = \rho_y$, two MCB models are the same and we can set $\rho_{xy} = \rho_x = \rho_y$. Two models merge completely and we have a $(N + M)$ assets MCB model. As the asymmetry between the two models becomes large ($p_x \neq p_y$ or $\rho_x \neq \rho_y$), $\text{Max}(\rho_{xy})$ decreases from ρ_x, ρ_y .
- (ii) $\lambda_x, \lambda_y > 0$ Case : As λ_x, λ_y increase, $\text{Max}(\rho_{xy})$ becomes large. The asymmetry in p_x, p_y and ρ_x, ρ_y diminishes $\text{Max}(\rho_{xy})$.

Using the above coupled $N + M$ MCB model, we study the effect of the non-uniformity of the default probability on $P_N(n)$ and on the evaluation of tranches. We consider a portfolio with $N + N$ assets. The assets in each sector has default probabilities $p \pm \Delta P_d$ and the intra-sector default correlation ρ . We set the inter-sector default correlation ρ_{xy} also as ρ . The inhomogeneity in the default probability is controlled by ΔP_d . If we set $\Delta P_d = 0$, the two sector are completely merged to one sector and we have a homogeneous $2N$ MCB model with p and ρ . In figure 9, we shows the default probability difference $\Delta P(n)$ between the inhomogeneous case $P_{N+N}(n)$ with $\Delta P_d \neq 0$ and homogeneous case $P_{2N}(n)$. $\Delta P(n)$ is defined as

$$\Delta P(n) = P_{N+N}(n) - P_{2N}(n). \tag{51}$$




Figure 9. $\Delta P(n)$ vs n . We set $N = 20, p = 0.03, \rho = 0.03$ and $\lambda_x = \lambda_y = 0.3$. MCB in red, green lines and BBD in blue and violet symbols.

We set $N = 20, p = \rho = 0.03$ and $\lambda = 0.3$. The solid red curve represents the data for $\Delta P_d = 0.01$ and the dotted green curve stands for the case $\Delta P_d = 0.02$. We see that $\Delta P(n)$ is large only for small n . Only the tranches with lower seniority are affected by the inhomogeneity ΔP_d . We also plot the results for the $N + M$ BBD model. To construct the loss distribution function, it is necessary to change (39) and (40) to

$$\text{Cor}(X_{n+1}X_{n+2} | \prod_{n'=1}^n X_{n'} = 1) = \rho_x / (1 + n\rho_x) \quad (52)$$

$$\text{Cor}(Y_{m+1}Y_{m+2} | \prod_{m'=1}^m Y_{m'} = 1) = \rho_y / (1 + m\rho_y). \quad (53)$$

The recursive relations are also changed, the remaining procedures are the same with $N + M$ MCB model. As we have shown in the previous section, the bulk shapes of the loss distribution of MCB and BBD models are almost the same, the effects of ΔP_d on them are also similar.

To see the effect of ΔP_d on the evaluations of tranches, we study the change in the cumulative distribution functions $D(i)$. $D(i)$ are defined as

$$D(i) = \sum_{n=i}^N P_N(n). \quad (54)$$

$D(i)$ represents the expected loss rate of the i -th tranche, which is damaged if more




Figure 10. $\Delta D(i)$ vs ΔP_d . The parameters are the same with the previous figure. We plot $D(n)$ for $1 \leq n \leq 8$.

than i assets are defaulted [22]. One of the important properties of $D(i)$ is

$$\frac{1}{N} \sum_{i=1}^N D(i) = p. \quad (55)$$

This identity means that the tranches distribute the portfolio credit risk between them. Another important property of $D(i)$ is that the expected loss rate $D[i, j]$ of the layer protection with attachment point i and detachment point j can be built up from $D(i)$ as

$$D[i, j] = \frac{1}{j - i + 1} \sum_{n=i}^j D(n). \quad (56)$$

Figure 10 shows the difference $\Delta D(i)$ between the inhomogeneous case and homogeneous case. The parameters are the same with those of figure 9. We see that $|\Delta D(n)|$ increase with ΔP_d for any n . For fixed ΔP_d , $|\Delta D(n)|$ for $n \geq 5$ becomes smaller than $|\Delta D(4)|$ and the inhomogeneity does not affect the evaluation of tranche with $n \geq 5$ so much as $n = 4$. We conclude that the tranches with higher seniority are not affected by ΔP_d . For $N = 20 \times 2, p = \rho = 0.03$ case, the border line is $n = 4$.

3.4. Multi Sectors Case

We try to couple two or more MCB models. We consider a portfolio with N assets, which are categorized in K different sectors. Figure 11 sketches the situation of the portfolio. The k -th sector contains N_k assets and the relation $\sum_{k=1}^K N_k = N$ holds. The states of the assets in the k -th sector is described by $X_{n_k}^k (n_k = 1, \dots, N_k)$ and the default rate and default correlation are denoted as p_k and ρ_k . About the inter-sector default correlation, we denote them as ρ_{ij} for the default correlation between the i -th and j -th sector. The intra-sector default correlation and inter-sector default correlation are different and the former is larger than the latter in general [25].




Figure 11. Structure of the portfolio. There are K sectors and k -th sector contains N_k assets. The default probability and default correlation for the assets in the k -th sector is p_k, ρ_k . The inter-sector default correlation between the i -th and j -th sector is ρ_{ij} .

We have not yet succeeded in the coupling of three or more MCB models, by generalizing the result for the coupled $N + M$ MCB model. The reason is that the self-consistency relations are very rigid restrictions on the MCB model. These relations, two sectors version is given by (46), assure the independence of the order of the decomposition in the estimation of the expectation values of the product of random variables. It is difficult to impose any simple relations on the conditional default probabilities that satisfy the self-consistency relations.

In order to construct the joint default probabilities for the assets states $(\vec{x}^1, \vec{x}^2, \dots, \vec{x}^K)$, we do not glue together K MCB models directly. Instead, as depicted in figure 12, we glue multiple MCB models through one random variable Y . More

concretely, we prepare K sets of $N_k + 1$ MCB models. $N_k + 1$ MCB model is the N_k MCB model coupled with Y . The probability of $P(Y = 1)$ is written as p_y . We introduce the following conditional default probabilities

$$\begin{aligned} p_{n_k,0}^k &= \langle X_{n_k+1}^k | \prod_{n'_k=1}^{n_k} X_{n'_k}^k = 1 \rangle \quad \text{and} \quad p_{n_k,1}^k = \langle X_{n_k+1}^k | \prod_{n'_k=1}^{n_k} X_{n'_k}^k \times Y = 1 \rangle \\ q_{n_k}^k &= \langle Y | \prod_{n'_k=1}^{n_k} X_{n'_k}^k = 1 \rangle. \end{aligned} \quad (57)$$

We also impose the following conditions on $p_{n_k,0}^k, p_{n_k,1}^k$ and $q_{n_k}^k$ as

$$\text{Cor}(X_{n_k+1}^k, X_{n_k+2}^k | \prod_{n'_k=1}^{n_k} X_{n'_k}^k = 1) = \rho_k \exp(-n_k \lambda) \quad (58)$$

$$\text{Corr}(X_{n_k+1}^k, Y | \prod_{n'_k=1}^{n_k} X_{n'_k}^k) = \rho_{ky} \exp(-n_k \lambda). \quad (59)$$

The joint default probabilities $P(\vec{x}^k, y)$ and the conditional joint default probabilities $P^k(\vec{x}^k | y)$ are constructed as before.

$$P(\vec{x}^k, y) = \langle \prod_{n_k=1}^{N_k} (X_{n_k}^k)^{x_{n_k}^k} (1 - X_{n_k}^k)^{1-x_{n_k}^k} \times Y^y (1 - Y)^{1-y} \rangle \quad (60)$$

$$P(\vec{x}^k, y) = P^k(\vec{x}^k | y) P(y). \quad (61)$$

Packing these conditional default probabilities $P^k(\vec{x}^k | y)$ into a bundle, we construct the joint default probabilities for the total portfolio as

$$P(\vec{x}^1, \vec{x}^2, \dots, \vec{x}^K) = \sum_{y=0,1} p(y) \times \prod_{k=1}^K P(x_1^k, x_2^k, \dots, x_{N_k}^k | y). \quad (62)$$

We also obtain the default probability function $P_N(n_1, n_2, \dots, n_K)$ for n_k default in the k -th sector as

$$P_N(n_1, n_2, \dots, n_K) = \sum_{y=0,1} p(y) \times \prod_{k=1}^K P_{N_k}(n_k | y) \quad (63)$$

From the expression, it is easy to calculate the probability for n defaults and we write it as $P_N(n)$.

About the default correlation between the different sectors, we can show the next relations.

$$\rho_{ij} = \rho_{iy} \times \rho_{jy}. \quad (64)$$

More generally, the conditional inter-sector default correlations obey the following relations.

$$\text{Cor}(X_{n_i+1}^i, X_{n_j+1}^j | \prod_{n'_i=1}^{n_i} X_{n'_i}^i \prod_{n'_j=1}^{n_j} X_{n'_j}^j = 1) = \rho_{ij} \exp(-(n_i + n_j) \lambda) \quad (65)$$

These relations mean that our construction procedure is natural from the viewpoint of original MCB model. In particular, in the $K = 2$ case, these relations are completely




Figure 12. Gluing multiple MCB models with Y . The correlation between $X_{n_k}^k$ and Y is ρ_{ky} . The default correlation between $X_{n_i}^i$ and $X_{n_j}^j$ is given as $\rho_{ij} = \rho_{iy} \times \rho_{jy}$.

equivalent with those of the $N + M$ coupled MCB model. See (45) and (65). The conditional default probabilities obey the same conditions.

Furthermore, from the results on $\text{Max}(\rho_{xy})$ of the $N + 1$ MCB model, we see that the model can induce realistic magnitude of inter-sector default correlation. By choosing λ and p_y properly, it is possible to take ρ_{ky} as large as several times of ρ_k .

Figure 13 plots $\text{Max}(\rho_{xy}) = \text{Max}(\rho_{ky})$ as functions of $\rho = \rho_k$. We set parameters as depicted in the figure. The green line depicts the data for $r = \frac{p_y}{p_x} = 3.0$. Other two curves correspond to $r = 1.0$ (+) and $r = 15.0$ (\times). We see that by setting $r = 3.0$, it is possible to set ρ_{ky} as large as several times of ρ_k . If the intra-sector correlation $\rho_i = \rho_j$ is 10%, we can set $\rho_{iy} = \rho_{jy} = 20\%$. The inter-sector correlations is then $\rho_{ij} = \rho_{iy} \times \rho_{jy} = 4\%$. In general, ρ_{inter} is smaller than ρ_{intra} , we think that the present model can incorporate strong enough inter-sector default correlation.

We prove the relations (64) and (65). In order to calculate the correlation, we need to estimate the next expression.

$$\langle X_{n_i+1}^i X_{n_j+1}^j | \prod_{i'=1}^{n_i} X_{i'}^i \prod_{j'=1}^{n_j} X_{j'}^j = 1 \rangle. \quad (66)$$

If we fix the random variable Y , $X_{n_i}^i$ and $X_{n_j}^j$ are independent. They are coupled by Y




Figure 13. Plot of $\text{Max}(\rho_{xy})$ vs ρ . $N = 30, p_x = 0.03$ and $\lambda = 0.3$.

and the above equation is estimated by the average over $Y = 0$ and $Y = 1$ as

$$\begin{aligned}
& \langle X_{n_i+1}^i X_{n_j+1}^j \mid \prod_{i'=1}^{n_i} X_{i'}^i \prod_{j'=1}^{n_j} X_{j'}^j = 1 \rangle \\
&= \langle X_{n_i+1}^i \mid \prod_{i'=1}^{n_i} X_{i'}^i = 1, Y = 1 \rangle \langle X_{n_j+1}^j \mid \prod_{j'=1}^{n_j} X_{j'}^j = 1, Y = 1 \rangle p_y \\
&+ \langle X_{n_i+1}^i \mid \prod_{i'=1}^{n_i} X_{i'}^i = 1, Y = 0 \rangle \langle X_{n_j+1}^j \mid \prod_{j'=1}^{n_j} X_{j'}^j = 1, Y = 0 \rangle (1 - p_y) \\
&= p_{n_i,1}^i p_{n_j,1}^j p_y + \tilde{p}_{n_i,0}^i \tilde{p}_{n_j,0}^j (1 - p_y). \tag{67}
\end{aligned}$$

Here, we denote the conditional default probabilities with the condition $Y = 0$ as $\tilde{p}_{n_i,0}^i$,

$$\tilde{p}_{n_i,0}^i = \langle X_{n_i+1}^i \mid \prod_{n'_i=1}^{n_i} X_{n'_i}^i \times (1 - Y) = 1 \rangle. \tag{68}$$

Between $p_{n_i,0}^i$ and $\tilde{p}_{n_i,0}^i$, the next relation holds.

$$p_{n_i,0}^i = p_{n_i,1}^i p_y + \tilde{p}_{n_i,0}^i (1 - p_y). \tag{69}$$

In addition, from the correlation between $X_{n_i}^i$ and Y , we also have the next relations.

$$p_{n_i,1}^i p_y = p_{i,0} p_y + \rho_{xy} \sqrt{p_{i,0} (1 - p_{i,0}) p_y (1 - p_y)} \tag{70}$$

Putting these relations into (67), we can prove the next equations.

$$\begin{aligned}
& \langle X_{n_i+1}^i X_{n_j+1}^j \mid \prod_{i'=1}^{n_i} X_{i'}^i \prod_{j'=1}^{n_j} X_{j'}^j = 1 \rangle \\
&= p_{n_i,0}^i p_{n_j,0}^j + \rho_{iy} \times \rho_{jy} \sqrt{p_{n_i,0}^i (1 - p_{n_i,0}^i) p_{n_j,0}^j (1 - p_{n_j,0}^j)} \times e^{-\lambda(n_i+n_j)} \tag{71}
\end{aligned}$$

Using these relations, we calculate the inter-sector correlation and prove (65).




Figure 14. Comparison of Two sectors MCB model: Directly Coupled Model vs Coupled by Y Mode ($p_y = 0.5$). Plot of $\Delta P(n) = P_{2N}(n) - P_{N+N}(n)$ vs $n/(2N)$. $N = 5, 10, 20, p = \rho = 0.03$ and $\lambda = 0.3$. Upper three curves have $\rho_{xy} = 0.01$ and the bottom line has $\rho_{xy} = 0.02$.

We check the validity of the above glueing process. We consider the two sector case $K = 2$ and their intra-sector parameters are set to be the same as N , p and ρ in each sector. About the inter-sector default correlation ρ_{12} , we set $\rho_{1y}^2 = \rho_{2y}^2 = \rho_{12}$ in the above multi-sector model. If the glueing process of the multi-sector model works well, $P_{2N}(n)$ should coincide with $P_{N+N}(n)$ of the coupled $N + N$ MCB model in the previous subsection. Figure 14 shows $\Delta P(n) = P_{2N}(n) - P_{N+N}(n)$ with $N = 5, 10, 20$ and $\lambda = 0.3$. We set $p = \rho = 0.03$ and $\rho_{12} = 0.01$. In addition we also plot the data for $\rho_{12} = 0.02$ and $N = 20$. As the system size N becomes large, the discrepancy $\Delta P(n)$ increases. With the same system size $N = 20$, as the inter-sector correlation increases, the discrepancy also increases. As we have stated previously, these models obey the same conditions on the conditional default probabilities, however default probability profile does not coincide. The glueing process by the auxiliary random variable Y may cause changes to the joint probabilities. We have not yet understood this point well.

With the present model, we study the effects of the inhomogeneous default correlation $\rho_{inter} \neq \rho_{intra}$ on the default distribution function $P_N(n)$ and the loss rates $D(i)$. We consider ideal portfolios which have the same default probability $p_k = p$, default correlation $\rho_k = \rho$. The inter-sector default correlations are also set to be the same value as $\rho_{ij} = \rho_{inter}$. N_k are also set to be the same $N_k = N_s$. We fix the total number of assets as N and compare $P_N(n)$ and $D(i)$ between the portfolios with different sector number K . Of course, between K and N_s the relation $N = N_s \times K$ holds. We

set $\rho_{inter} < \rho$. As the number K increases, the average default correlation is governed by the inter-sector correlation ρ_{inter} and becomes weak.

At the extreme limit $K = N$ case, each sector contains only one asset. By the inter-sector default correlation, $P_N(n)$ is given by the super-position of $P_N(n|Y = 1)$ and $P_N(n|Y = 0)$. The conditional default probabilities $p_{1,0}^k$, $p_{1,1}^k$ and $\tilde{p}_{1,0}^k$ are given as

$$p_{1,0}^k = p \quad \text{and} \quad p_{1,1}^k = p + \sqrt{\rho_{inter}}(1-p) \sqrt{\frac{p}{1-p} \frac{(1-p_y)}{p_y}} \quad (72)$$

$$\tilde{p}_{1,0}^k = p - \sqrt{\rho_{inter}} p \sqrt{\frac{1-p}{p} \frac{p_y}{(1-p_y)}}. \quad (73)$$

$P_N(n|Y = 1)$ and $P_N(n|Y = 0)$ are the binomial distributions $\text{Bin}(N, p_{1,1}^k)$ and $\text{Bin}(N, \tilde{p}_{1,0}^k)$.




Figure 15. Semi-log plot of $P_N(n)$ vs n . $N = 100, K = 2, 10, 100, p = \rho = 0.03, \rho_{inter} = 0.01$ and $\lambda = 0.3$. We set $p_y = 0.5$. Solid line represents $K = 1$ and $\rho = 0.03$ case and the dotted line shows the curve for $K = 1$ and $\rho = 0.01$.

Figure 15 shows the semi-log plot of the default probability $P_N(n)$ for $K = 1, 2, 10, 50, 100$. We set the model parameters as $p = \rho = 0.03$, $N = 100$, $\rho_{inter} = 0.01$ and $p_y = 0.5$. The red solid curve represents the data for $K = 1$ and $\rho = 0.03$ and the yellow line shows the data for $K = 1$ and $\rho = 0.01$. As K increases, the data for each K departs from the red line. At $K = 10$, the data almost shrinks on the yellow line ($K = 1$ and $\rho = 0.01$). The data for $K = 50$ almost coincide with those of $K = 100$, whose $P_N(n)$ is given by the superposition of the two binomial distributions. This point

is also a drawback of the present model. If the glueing process works perfect, these data should coincide with the homogeneous portfolio case $K = 1$ and $\rho = 0.01$. However, this discrepancy is inherent property of the model. Contrary to the discrepancy of the two-sector case, the conditions on the conditional default probabilities $p_{n_k,1}^k$ are different from those of the $K = 1$ homogeneous portfolio. They could cause the difference in $P_N(n)$.




Figure 16. Plot of $|\Delta D(i)|$ and $D(i)$ vs i . $\Delta D(i)$ are between $\rho = 0.01, 0.05$ and $\rho = 0.03$. $N = 100, K = 2, 10, 100, p = 0.03, \rho_{inter} = 0.01$ and $\lambda = 0.3$. We set $p_y = 0.5$. The solid curve shows the data of $D(i)$ with $\rho = 0.03$.

From the above discussions on $P_N(n)$ with different K , we think that the inter-sector default correlation ρ_{inter} is more important than the intra-sector default correlation ρ for the large K case. In the $K = 10$ case, the $P_N(n)$ are roughly given by those of the homogeneous portfolio with $\rho = \rho_{inter}$. If one estimates the implied values of ρ and ρ_{inter} from the premium (or $D(i)$) of the portfolio with large K , this point is crucial. In Figure 16, we show $|\Delta D(i)|$ for $K = 10$ between $\rho = 0.01, 0.05$ and $\rho = 0.03$. We set the model parameters as in the previous figure. For comparison, we also plot $D(i)$ for $\rho = 0.03$. $D(i)$ represents the expected loss rate of the i -th tranche, the magnitude of $|\Delta D(i)|$ is important when one estimates the implied default correlation ρ from the premium of the tranche. We see that $|\Delta D(i)|$ with small i is small as compared with $D(i)$. If we change ρ from $\rho = 0.03$, $D(i)$ does not change much. It is difficult to obtain the implied values of ρ from the premium of the tranche with the lower seniority. On the other hands, with medium value of $i \simeq 13$, the magnitude of $D(i)$ and $\Delta D(i)$ are almost comparable. $D(i)$ is sensitive to the change in ρ and it is not difficult to get the implied value of ρ .

4. Implied Default Correlation and Concluding Remarks

In the last section, as a concrete example, we try to estimate the implied values of the default correlation from the premium of a synthetic CDO. We treat the iTraxx-CJ, which is an equally weighted portfolio of 50 CDSs on Japanese companies. The standard attachment points and detachment points are $\{0, 3\%$, $\{3\%, 6\%$, $\{6\%, 9\%$, $\{9\%, 12\%$ and $\{12\%, 22\%$. Table 1 shows quotes on July 5, 2005. The quotes for the $\{0\%, 3\%$ tranche shows the upfront payment (as a percent of principal) that must be paid in addition to 300 basis points per year. The other quotes for the other tranches are the annual payment rates in basis points per year. The index indicates the cost of entering into a CDS on all 50 companies underlying the index. The recovery rate R is 0.35.

Table 1. Quotes for iTraxx-CJ Tranches on July 5, 2005. Quotes for the $\{0, 3\%$ tranche are the percent of the principal that must be paid up front in addition to 300 basis points per year. Quotes for other tranches and the index are in basis points. Source: Morgan Stanley Japan Securities Co. and Bloomberg

	$\{0, 3\%$	$\{3\%, 6\%$	$\{6\%, 9\%$	$\{9\%, 12\%$	$\{12\%, 22\%$	Index
5-year Quotes	15.75	113.25	42.0	30.5	15.5	24.55

In order to get the implied default correlation for each, it is necessary to relate the loss distribution function $P_N(n)$ to the premiums. The premiums are the present value of the expected cash flows. The calculation of this present value involves three terms [27]. We denote by $P_k(a_L, a_H)$ the remaining notional for the $\{a_L, a_H\}$ tranche after the k -th default. It is given as

$$P_k(a_L, a_H) = \begin{cases} (a_H - a_L)N & k < \lceil a_L N / (1 - R) \rceil \\ a_H N - k(1 - R) & \lceil a_L N / (1 - R) \rceil \leq k \leq \lceil a_H N / (1 - R) \rceil \\ 0 & k \geq \lceil a_H N / (1 - R) \rceil \end{cases} \quad (74)$$

Here, $\lceil x \rceil$ means the smallest integer greater than x . For simplicity, we treat the 5-year as one period. The three terms are written as

$$\begin{aligned} A &= 5.0 \times \langle P_k(a_L, a_H) \rangle e^{-5.0r} \\ B &= 2.5 \times ((a_H - a_L)N - \langle P_k(a_L, a_H) \rangle) e^{-r \frac{5}{2}} \\ C &= ((a_H - a_L)N - \langle P_k(a_L, a_H) \rangle) e^{-r \frac{5}{2}} \end{aligned} \quad (75)$$

where r is the risk-free rate of interest and we set $r = 0.01$. By considering the total value of the contract, one see that the break even spread is given as $C/(A + B)$.

For the index, $a_H = 1.0$ and $a_L = 0.0$ and it is possible to estimate the average default probability p . Instead, we use the CDS data for each company and estimate the average default probability p and its volatility ΔP_d as

$$p = 1.8393\%/5\text{-year} \quad \Delta P_d = 1.131\%/5\text{-year}.$$

With these parameters, the tranche correlations can be implied from the spreads quoted in the market for particular tranches. These correlations are known as tranche

correlations or compound correlation. As a pricing model, we use MCB, BBD and Gaussian copula models. For MCB, we use the following candidates.

- Original MCB model (MCB1). $N = 50, K = 1$ and $\lambda = 0.0$.
- Short tail MCB model (MCB2). $N = 50, K = 1$ and $\lambda = 0.3$.
- Disordered MCB model (MCB3). $N = 25 + 25, \lambda = 0.3, \rho_{xy} = \rho_x = \rho_y = \rho$ and $p_x = p + \Delta P_d, p_y = p - \Delta P_d$. $N + M$ MCB model with inhomogeneous default probability.
- 2-sector MCB model (MCB4). $K = 2, \rho_x = \rho_y = \rho, p_x = p_y = p$ and $\rho_{xy} = 0.0$. Assets are categorized in 2 sectors and $\rho_{inter} = 0.0$.

Table 2. Implied tranche correlation (%) for 5-year iTraxx-CJ on July 5,2005.

Tranches	MCB 1	MCB 2	MCB 3	MCB 4	BBD	Gaussian
{0%, 3%}	11.79	10.8	15.75	21.2	11.4	13.8
{3%, 6%}	1.27	1.18	1.13	2.45	1.26	1.35
{6%, 9%}	3.16	3.08	0.42	6.32	3.15	3.23
{9%, 12%}	6.16	5.95	0.30	12.15	6.11	6.31
{12%, 22%}	9.78	9.67	0.15	19.97	9.73	9.46

Table 2 shows implied tranche correlations for the 5-year quotes in Table 1. We see ‘‘Correlation Smile’’, which is a typical behavior of implied correlations across portfolio tranches [13]. We also find that the implied correlations are different between the models. About MCB, the model with large λ has narrow range of the correlation skew. The correlation for {0%, 3%} decrease with λ and other correlations does not move much. If a probabilistic model describe the true default distribution, there should not exist the correlation skew. As a model approaches the true distribution, we can expect that the skew decreases. MCB with $\lambda = 0.3$ is more faithful to the true default distribution. MCB with $\lambda = 0.0$ and BB have almost the same skew range. Gaussian copula’s skew range is more larger than MCB 1, MCB 2 and BBD. We think that its shape is far from the true default distribution.

About the effect of ΔP_d , the implied correlations are considerably different between $\Delta P_d \neq 0$ (MCB 3) and $\Delta P_d = 0.0$ (MCB 2). In the estimation of the implied correlation, we cannot neglect the fluctuation ΔP_d . In particular, about the tranches {0%, 3%}, the implied value is affected much by ΔP_d . As we have seen in subsection 3.3, the cumulative function $D(1)$ of the equity increases much with ΔP_d . In order to suppress the increase, it is necessary to increase the correlation.

$K = 2$ sector case (MCB4), the correlation level and correlation skew are very large. This means that if assets are categorized in many sectors and inter-sector correlation is very weak, the loss distribution accumulates around the origin $n = 0$, as in the MCB 3. In order to realize the market quotes, large intra-sector correlation is necessary.

About the estimation of the model parameter λ , we think that the resulting loss distribution should look closer to the implied loss distribution. That is, the range of tranche correlation skew should be small. Figure 17 shows the tranche correlation vs λ . As we increase λ , the skew range becomes small. At $\lambda \simeq 0.6$, the range becomes minimum. We should calibrate λ to be $\lambda = 0.6$.




Figure 17. Range of implied tranche correlation for 5-year iTraxx-CJ on July 5, 2005 vs λ .

In this paper, we generalize the Moody's correlated binomial default distribution to the inhomogeneous portfolio cases. As the inhomogeneity, we consider the non-uniformity in the default probability p and in the default correlation ρ and ρ_{inter} . To treat the former case, we construct a coupled $N + M$ MCB model and obtain the default probability function $P_{N+M}(n)$. The inhomogeneity in p causes the changes in the expected loss rates of the tranches with lower seniority. The tranches with higher seniority are not affected. In order to treat the inhomogeneity in the default correlation, we construct a multi-sector MCB model, by glueing multiple MCB models by an auxiliary random variable Y . We cannot take out the lines of joint between the MCB models, for small portfolio and small ρ_{inter} , the construction works well. About the inhomogeneity in ρ , we divide a homogeneous portfolio in K sectors. We set $\rho_{inter} < \rho$ and see the effect of the increase in K on $P_N(n)$. As the sector number K increases, the inter-sector correlation ρ_{inter} becomes more important than the intra-sector default correlation ρ . With large K , the default correlation is governed by ρ_{inter} only. The CDOs, whose assets are categorized in many sectors, ρ_{inter} should be treated more carefully than ρ .

In order to check the validity of the MCB model and generalization method,

more careful treatment and calibration should be done. We assume that ρ_n decays exponentially with n . **With such a modification, the skew range of the tranche correlations diminishes, however, the skew remains much.** Other model for ρ_n should be considered. For the purpose, it is necessary to study the implied loss distribution directly. Recently, Hull and White [26, 27] developed a method to get the implied loss distribution and to obtain the implied copula correlation from the market quotes of CDOs. By analyzing the conditional correlation ρ_n or default probability p_n of the implied loss distribution, it may be possible to propose a good model for ρ_n . We may have a "Perfect" correlated binomial default distribution model which reflects the market quotes completely.

Acknowledgments

This work has received financial support from Kitasato University, project SCI:2005-1706.

References

- [1] Fabozzi, F. J. and L. S. Goodman, 2001 *Investing in Collateralized Debt Obligations*, U.S. John Wiley & Sons.
- [2] Schönbucher, P., 2003, *Credit Derivatives Pricing Models : Model, Pricing and Implementation*, U.S. John Wiley & Sons.
- [3] Duffie, D. and K.J.Singleton, 2003, *Credit Risk-Pricing, Measurement and Management* (Princeton : Princeton University Press).
- [4] Hull, J., M. Predescu and A. White, 2005, *The Valuation of Correlation-Dependent Credit Derivatives Using a Structural Model*, Working Paper, University of Toronto.
- [5] Finger, C. C., 2005, *Issues in the Pricing of Synthetic CDOs*, Journal of Credit Risk, **1(1)**.
- [6] Cifuettes, A. and G. O'Connor, 1996, *The Binomial Expansion Method Applied to CBO/CLO Analysis*, Working Paper (Moody's Investors Service).
- [7] Martin, R., Thompson, K. and C. Browne, 2001, *How dependent are defaults*, Risk Magazine, **14(7)** 87-90.
- [8] Finger, C. C., 2000, *A Comparison of stochastic default rate models*, Working Paper (The RiskMetrics Group).
- [9] Duffie, D. and N. Gârleau, 2001, *Risk and the Valuation of Collateralized Debt Obligation*, Financial Analyst Journal **57(1)** 41-59.
- [10] Li, D., 2000, *On Default Correlation: a Copula Approach*, The Journal of Fixed Income **9(4)** 43.
- [11] Vasicek, O., 1987, *Probability of Loss on Loan Portfolio*, Working Paper (KMV Corporation).
- [12] Schönbucher, P. and D. Schubert, 2001, *Copula Dependent Default Risk in Intensity Models*, Working paper (Bonn University).
- [13] Andersen, L., J.Sidenius and S.Basu, 2003, *All your Hedges in one Basket*, RISK, 67-72.
- [14] Davis, M. and V. Lo, *Infectious defaults*, Quantitative Finance, **1**, 382-387.
- [15] Zhou, C., 2001, *An analysis of default correlation and multiple defaults*, Review of Financial Studies **4**, 555-576.
- [16] Laurent J-P. and J. Gregory, 2003, *Basket Default Swaps, CDO's and Factor Copulas*, Journal of Risk **7(4)**, 103-122.
- [17] CREDIT-SUISSE-FINANCIAL-PRODUCTS, 1997, *CreditRisk+ a Credit Risk Management Framework*, Technical Document.
- [18] Frey, R., Mcneil, A. and M. Nyfeler, 2001, *Copulas and Credit Models*, Risk, October, 111-113.

- [19] Frey, R. and A. McNeil, 2002, *VaR and Expected Shortfall in Portfolios of Dependent Credit Risks: Conceptual and Practical Insights*, Journal of Banking and Finance, 1317-1344.
- [20] Witt, G. ,2004, *Moody's Correlated Binomial Default Distribution*, Working Paper (Moody's Investors Service) **August 10**.
- [21] Molins, J. and E.Vives, 2004,*Long range Ising Model for credit risk modeling in homogeneous portfolios*, Preprint arXiv:cond-mat/0401378.
- [22] Kitsukawa, K., Mori, S. and M. Hisakado,2006,*Evaluation of Tranche in Securitization and Long-range Ising Model*, Physica **A 368** 191-206.
- [23] Hisakado, M, Kitsukawa, K. and S. Mori ,2006,*Correlated Binomial Models and Correlation Structures*, Preprint arXiv:physics/0605189.
- [24] Standard & Poor's CreditPro 7.00, 2005.
- [25] Jobst, N. J. and A. de Servigny, 2005,*An Empirical Analysis of Equity Default Swaps (II): Multivariate insights*, Working Paper (S&P).
- [26] Hull, J. and A. White, 2005,*The Perfect Copula*, Working Paper,(University of Toronto).
- [27] Hull, J. and A. White, 2006,*Valuing Credit Derivatives Using an Implied Copula Approach*, Working Papper (University of Toronto).