

# Gravity and the Electroweak Theory

Kenneth Dalton

e-mail: kxdalton@yahoo.com

## Abstract

This work shows that gravity creates inertia for the electron. The Lagrangian directly couples the gravitational field with the electron spinor field. It does so via the covariant spinor derivative. The Lagrangian is invariant with respect to the electroweak gauge group  $U(1) \otimes SU(2)_L$ .

The field equations are solved for an electron in uniform motion. The solution is the Dirac spinor for a massive electron. In effect, gravitational coupling takes the place of the Dirac mass term.

## 1. Metrical Gravity.

The formalism of time-dependent, three-dimensional geometry was derived in [1] and reviewed in [2]. A brief summary follows. We introduce a scalar, 3-vector basis  $e_\mu = (e_0, \mathbf{e}_i)$  and define inner products

$$g_{\mu\nu} = e_\mu \cdot e_\nu = \begin{pmatrix} g_{00} & 0 & 0 & 0 \\ 0 & & & \\ 0 & & g_{ij} & \\ 0 & & & \end{pmatrix} \quad (1)$$

The basis  $e_\mu$  changes from point to point in the manifold according to the formula

$$\nabla_\nu e_\mu = e_\lambda Q_{\mu\nu}^\lambda \quad (2)$$

This separates into scalar and 3-vector parts

$$\nabla_\nu e_0 = e_0 Q_{0\nu}^0 \quad (3)$$

$$\nabla_\nu \mathbf{e}_i = \mathbf{e}_j Q_{i\nu}^j \quad (4)$$

By definition  $Q_{j\nu}^0 = Q_{0\nu}^i \equiv 0$ . The 18 coefficients  $Q_{jk}^i$  would suffice in the case of ordinary static, three-dimensional geometry. Here, the geometry will be time-dependent, in general, and clock rates may differ from point to point. Hence, the additional coefficients  $Q_{j0}^i$  and  $Q_{0\nu}^0$ , respectively. All 28 independent  $Q_{\nu\lambda}^\mu$  are derivable from the metric

$$Q_{0\nu}^0 = \Gamma_{0\nu}^0 = \frac{1}{2} g^{00} \partial_\nu g_{00} \quad (5)$$

$$Q_{j0}^i = \Gamma_{j0}^i = \frac{1}{2} g^{il} \partial_0 g_{lj} \quad (6)$$

$$Q_{jk}^i = \Gamma_{jk}^i = \frac{1}{2} g^{il} (\partial_k g_{jl} + \partial_j g_{lk} - \partial_l g_{jk}) \quad (7)$$

where

$$\Gamma_{\nu\lambda}^\mu = \frac{1}{2} g^{\mu\rho} (\partial_\lambda g_{\nu\rho} + \partial_\nu g_{\rho\lambda} - \partial_\rho g_{\nu\lambda}) \quad (8)$$

are the Christofel coefficients. The symbols  $\Gamma_{\nu\lambda}^\mu$  are symmetric in  $\nu\lambda$ , while the  $Q_{\nu\lambda}^\mu$  are not. The following formula holds good

$$Q_{\nu\lambda}^{\mu} = \Gamma_{\nu\lambda}^{\mu} + g^{\mu\rho} g_{\lambda\eta} Q_{[\nu\rho]}^{\eta} \quad (9)$$

where

$$Q_{[\nu\lambda]}^{\mu} \equiv Q_{\nu\lambda}^{\mu} - Q_{\lambda\nu}^{\mu} \quad (10)$$

The gravitational field is introduced by means of the Einstein-Hilbert action

$$\frac{\kappa}{2} \int R \sqrt{-g} d^4x = \frac{\kappa}{2} \int g^{\mu\nu} R_{\mu\nu} \sqrt{-g} d^4x \quad (11)$$

where  $\kappa = c^4/8\pi G$ , and the Ricci tensor is

$$R_{\mu\nu} = \partial_{\nu} \Gamma_{\mu\lambda}^{\lambda} - \partial_{\lambda} \Gamma_{\mu\nu}^{\lambda} + \Gamma_{\rho\nu}^{\lambda} \Gamma_{\mu\lambda}^{\rho} - \Gamma_{\lambda\rho}^{\lambda} \Gamma_{\mu\nu}^{\rho} \quad (12)$$

Variation of the gravitational action gives [3]

$$\delta \int \frac{\kappa}{2} g^{\mu\nu} R_{\mu\nu} \sqrt{-g} d^4x = \frac{\kappa}{2} \int \left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) \delta g^{\mu\nu} \sqrt{-g} d^4x \quad (13)$$

The source of gravitation is expressed in terms of a matter Lagrangian  $\mathcal{L}_m$ . Variation with respect to  $\delta g^{\mu\nu}$  defines the energy tensor

$$\delta \int \mathcal{L}_m d^4x = \frac{1}{2} \int T_{\mu\nu} \delta g^{\mu\nu} \sqrt{-g} d^4x \quad (14)$$

Setting the sum of (13) and (14) equal to zero, we obtain the gravitational field equations

$$\kappa \left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) + T_{\mu\nu} = 0 \quad (15)$$

There are seven field equations corresponding to the seven variations  $\delta g^{00}$  and  $\delta g^{ij}$ .

The Lagrangian for gravitation may be derived by partial integration of the action [3]

$$\begin{aligned} \frac{\kappa}{2} \int g^{\mu\nu} R_{\mu\nu} \sqrt{-g} d^4x &= \frac{\kappa}{2} \int \left\{ \partial_{\nu} \left( \sqrt{-g} g^{\mu\nu} \Gamma_{\mu\lambda}^{\lambda} \right) - \partial_{\lambda} \left( \sqrt{-g} g^{\mu\nu} \Gamma_{\mu\nu}^{\lambda} \right) \right. \\ &\quad + \partial_{\lambda} \left( \sqrt{-g} g^{\mu\nu} \right) \Gamma_{\mu\nu}^{\lambda} - \partial_{\nu} \left( \sqrt{-g} g^{\mu\nu} \right) \Gamma_{\mu\lambda}^{\lambda} \\ &\quad \left. + g^{\mu\nu} \left( \Gamma_{\rho\nu}^{\lambda} \Gamma_{\mu\lambda}^{\rho} - \Gamma_{\lambda\rho}^{\lambda} \Gamma_{\mu\nu}^{\rho} \right) \sqrt{-g} \right\} d^4x \quad (16) \end{aligned}$$

The first two terms, when converted to boundary integrals, do not contribute to the field equations. With some rearranging, this leaves

$$\frac{\kappa}{2} \int g^{\mu\nu} R_{\mu\nu} \sqrt{-g} d^4x = \frac{\kappa}{2} \int g^{\mu\nu} \left( \Gamma_{\mu\nu}^{\lambda} \Gamma_{\rho\lambda}^{\rho} - \Gamma_{\rho\nu}^{\lambda} \Gamma_{\mu\lambda}^{\rho} \right) \sqrt{-g} d^4x \quad (17)$$

The Lagrangian

$$\mathcal{L}_g = \frac{\kappa}{2} g^{\mu\nu} \left( \Gamma_{\mu\nu}^{\lambda} \Gamma_{\rho\lambda}^{\rho} - \Gamma_{\rho\nu}^{\lambda} \Gamma_{\mu\lambda}^{\rho} \right) \sqrt{-g} \quad (18)$$

depends upon the metric  $g^{\mu\nu}$  and its first derivatives  $\partial_{\lambda} g^{\mu\nu}$ . Therefore, variation (13) now takes the form

$$\begin{aligned} \delta \int \mathcal{L}_g d^4x &= \int \left[ \frac{\partial \mathcal{L}_g}{\partial g^{\mu\nu}} \delta g^{\mu\nu} + \frac{\partial \mathcal{L}_g}{\partial (\partial_{\lambda} g^{\mu\nu})} \delta (\partial_{\lambda} g^{\mu\nu}) \right] d^4x \\ &= \int \left[ \frac{\partial \mathcal{L}_g}{\partial g^{\mu\nu}} - \partial_{\lambda} \frac{\partial \mathcal{L}_g}{\partial (\partial_{\lambda} g^{\mu\nu})} \right] \delta g^{\mu\nu} d^4x \end{aligned} \quad (19)$$

Variation of the matter action (14) takes a similar form

$$\delta \int \mathcal{L}_m d^4x = \int \left[ \frac{\partial \mathcal{L}_m}{\partial g^{\mu\nu}} - \partial_{\lambda} \frac{\partial \mathcal{L}_m}{\partial (\partial_{\lambda} g^{\mu\nu})} \right] \delta g^{\mu\nu} d^4x \quad (20)$$

Defining the total Lagrangian

$$\mathcal{L} = \mathcal{L}_g + \mathcal{L}_m \quad (21)$$

and setting the sum of (19) and (20) equal to zero, we obtain the gravitational field equations

$$\frac{\partial \mathcal{L}}{\partial g^{\mu\nu}} - \partial_{\lambda} \frac{\partial \mathcal{L}}{\partial (\partial_{\lambda} g^{\mu\nu})} = 0 \quad (22)$$

The Lagrange equations have a direct bearing upon the question of energy, which is defined in terms of  $\mathcal{L}$  (appendix C).

## 2. Tetrads.

In order to accommodate spinors, the system  $e_\mu$  can be expanded in terms of tetrads on a Pauli basis

$$e_\mu = e^\alpha{}_\mu(x) \sigma_\alpha \quad (23)$$

where

$$e^\alpha{}_\mu = \begin{pmatrix} e^0{}_0 & 0 & 0 & 0 \\ 0 & & & \\ 0 & & e^a{}_i & \\ 0 & & & \end{pmatrix} \quad (24)$$

The tetrads can be chosen such that

$$e^1{}_2 = e^2{}_1 \quad e^2{}_3 = e^3{}_2 \quad e^3{}_1 = e^1{}_3 \quad (25)$$

leaving seven independent functions  $e^\alpha{}_\mu(x)$ . The metric now depends upon the underlying tetrad field

$$g_{\mu\nu} = \eta_{\alpha\beta} e^\alpha{}_\mu e^\beta{}_\nu \quad (26)$$

As shown in appendix A, the covariant spinor derivative

$$D_\mu \psi = \partial_\mu \psi + iqA_\mu \psi + \Gamma_\mu \psi \quad (27)$$

yields the Lagrangian (adding the kinetic term for  $A_\mu$ )

$$\begin{aligned} L_e = & \frac{i}{2} \hbar c [\bar{\psi} \gamma^\mu \partial_\mu \psi - (\partial_\mu \bar{\psi}) \gamma^\mu \psi] + \frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\alpha{}^\lambda e_\gamma{}^\nu \partial_\nu e_{\beta\lambda} \\ & - \hbar c q \bar{\psi} \gamma^\mu \psi A_\mu - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \end{aligned} \quad (28)$$

$\hat{\gamma}^\alpha$  are the constant Dirac matrices, while  $\gamma^\mu(x) = e_\alpha{}^\mu(x) \hat{\gamma}^\alpha$ . The electron field equation is found by varying  $\bar{\psi}$

$$\frac{\partial \mathcal{L}_e}{\partial \bar{\psi}} - \partial_\mu \frac{\partial \mathcal{L}_e}{\partial (\partial_\mu \bar{\psi})} = 0 \quad (29)$$

Substitution of

$$\frac{\partial \mathcal{L}_e}{\partial \bar{\psi}} = \hbar c \left\{ \frac{i}{2} \gamma^\mu \partial_\mu \psi - q \gamma^\mu \psi A_\mu + \frac{1}{4} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\alpha^\lambda e_\gamma^\nu \partial_\nu e_{\beta\lambda} \right\} \sqrt{-g} \quad (30)$$

$$\frac{\partial \mathcal{L}_e}{\partial (\partial_\mu \bar{\psi})} = -\frac{i}{2} \hbar c \sqrt{-g} \gamma^\mu \psi \quad (31)$$

gives

$$i \gamma^\mu \partial_\mu \psi + \frac{i}{2} \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} \gamma^\mu) \psi - q \gamma^\mu \psi A_\mu + \frac{1}{4} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\alpha^\lambda e_\gamma^\nu \partial_\nu e_{\beta\lambda} = 0 \quad (32)$$

A similar calculation with respect to  $\delta\psi$  yields the conjugate equation

$$-i (\partial_\mu \bar{\psi}) \gamma^\mu - \frac{i}{2} \bar{\psi} \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} \gamma^\mu) - q \bar{\psi} \gamma^\mu A_\mu + \frac{1}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \epsilon^{\delta\alpha\beta\gamma} e_\alpha^\lambda e_\gamma^\nu \partial_\nu e_{\beta\lambda} = 0 \quad (33)$$

Multiply the first equation by  $\bar{\psi}$  the second by  $\psi$  and subtract, to find conservation of charge

$$\partial_\mu (\sqrt{-g} \bar{\psi} \gamma^\mu \psi) = 0 \quad (34)$$

In the following sections, we will be concerned with oscillating gravitational fields of very small amplitude. The coordinate system is taken to be nearly rectangular

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (35)$$

$$g^{\mu\nu} = \eta^{\mu\nu} - h^{\mu\nu} \quad (36)$$

The indices of  $h_{\mu\nu}$  are raised by  $\eta^{\mu\nu}$ ; for example, in lowest order

$$\begin{aligned} g_{\mu\lambda} g^{\lambda\nu} &= (\eta_{\mu\lambda} + h_{\mu\lambda})(\eta^{\lambda\nu} - h^{\lambda\nu}) \\ &= \delta_{\mu}^{\nu} + h_{\mu}^{\nu} - h_{\mu}^{\nu} = \delta_{\mu}^{\nu} \end{aligned} \quad (37)$$

The tetrads are expressed in a similar fashion

$$e^{\alpha}_{\mu} = \delta^{\alpha}_{\mu} + \xi^{\alpha}_{\mu} \quad (38)$$

$$e_{\alpha}^{\mu} = \delta_{\alpha}^{\mu} - \xi_{\alpha}^{\mu} \quad (39)$$

so that

$$\begin{aligned} e^{\alpha}_{\mu} e_{\alpha}^{\nu} &= \delta_{\mu}^{\nu} = (\delta^{\alpha}_{\mu} + \xi^{\alpha}_{\mu})(\delta_{\alpha}^{\nu} - \xi_{\alpha}^{\nu}) \\ &= \delta_{\mu}^{\nu} + \delta_{\alpha}^{\nu} \xi^{\alpha}_{\mu} - \delta_{\mu}^{\alpha} \xi_{\alpha}^{\nu} \end{aligned} \quad (40)$$

In order to simplify the notation, we mix indices obtaining

$$\xi^{\nu}_{\mu} = \xi_{\mu}^{\nu} \quad (41)$$

In terms of tetrads, the metric is

$$\begin{aligned} g_{\mu\nu} &= \eta_{\alpha\beta} e^{\alpha}_{\mu} e^{\beta}_{\nu} \\ &= \eta_{\mu\nu} + \eta_{\alpha\beta} (\delta^{\alpha}_{\mu} \xi^{\beta}_{\nu} + \delta^{\alpha}_{\nu} \xi^{\beta}_{\mu}) \end{aligned} \quad (42)$$

so that

$$\begin{aligned} h_{\mu\nu} &= \eta_{\alpha\beta} (\delta^{\alpha}_{\mu} \xi^{\beta}_{\nu} + \delta^{\alpha}_{\nu} \xi^{\beta}_{\mu}) \\ &= 2\xi_{\mu\nu} \end{aligned} \quad (43)$$

### 3. $U(1) \otimes SU(2)_L$ Gauge Invariance. [4, 5, 6]

The gravitational coupling term in  $L_e$  (28) contains the factor  $\bar{\psi}\gamma_5\hat{\gamma}_\delta\psi$ . This factor does not mix right- and left-handed spinor components,  $\psi_R$  and  $\psi_L$ . In order to prove this, set

$$\psi = \psi_R + \psi_L = \frac{1 + \gamma_5}{2} \psi + \frac{1 - \gamma_5}{2} \psi \quad (44)$$

where  $(\gamma_5)^2 = 1$  and  $\gamma_5\gamma_\delta = -\gamma_\delta\gamma_5$ . Also,  $\bar{\psi}_R = \bar{\psi} \frac{1-\gamma_5}{2}$  and  $\bar{\psi}_L = \bar{\psi} \frac{1+\gamma_5}{2}$ . In the expansion

$$\begin{aligned} \bar{\psi}\gamma_5\hat{\gamma}_\delta\psi &= (\bar{\psi}_R + \bar{\psi}_L) \gamma_5\gamma_\delta (\psi_R + \psi_L) \\ &= \bar{\psi}_R\gamma_5\gamma_\delta\psi_R + \bar{\psi}_L\gamma_5\gamma_\delta\psi_L + \bar{\psi}_R\gamma_5\gamma_\delta\psi_L + \bar{\psi}_L\gamma_5\gamma_\delta\psi_R \end{aligned} \quad (45)$$

the mixed terms are identically zero. For example,

$$\bar{\psi}_R\gamma_5\gamma_\delta\psi_L = \bar{\psi} \frac{1 - \gamma_5}{2} \gamma_5\gamma_\delta \frac{1 - \gamma_5}{2} \psi = \bar{\psi} \gamma_5\gamma_\delta \frac{1 - (\gamma_5)^2}{4} \psi = 0 \quad (46)$$

Therefore,

$$\bar{\psi}\gamma_5\hat{\gamma}_\delta\psi = \bar{\psi}_R\gamma_5\gamma_\delta\psi_R + \bar{\psi}_L\gamma_5\gamma_\delta\psi_L \quad (47)$$

An expression of this type will be invariant under  $U(1) \otimes SU(2)_L$  gauge transformations. <sup>1</sup> Introduce the right-handed singlet  $\psi_R = e_R$  and left-handed doublet  $\psi_L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$  in order to form the Lagrangian

$$\begin{aligned} L_{e-w} &= \frac{i}{2} \hbar c [\bar{\psi}_R \gamma^\mu \partial_\mu \psi_R + \bar{\psi}_L \gamma^\mu \partial_\mu \psi_L] + \text{h.c.} \\ &+ \frac{\hbar c}{4} [\bar{\psi}_R \gamma_5 \hat{\gamma}_\delta \psi_R + \bar{\psi}_L \gamma_5 \hat{\gamma}_\delta \psi_L] \epsilon^{\delta\alpha\beta\gamma} e_\alpha^\lambda e_\gamma^\nu \partial_\nu e_{\beta\lambda} + L_{int} \end{aligned} \quad (48)$$

$L_{int}$  contains the electroweak interaction terms as well as kinetic terms for  $A_\mu$ ,  $W_\mu^\pm$ , and  $Z_\mu^0$ .

---

<sup>1</sup>The Dirac mass term  $m\bar{\psi}\psi = m(\bar{\psi}_R\psi_L + \bar{\psi}_L\psi_R)$  mixes right- and left-handed spinors and cannot appear in the electroweak Lagrangian.



#### 4. An Electron in Uniform Motion.

In the previous article [2], the gravitational field was found for a spin up electron at rest

$$\psi = \frac{1}{\sqrt{V}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \exp(-i\omega t) \quad (49)$$

which took the form

$$e^\alpha{}_\mu = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & e^1{}_1 & e^1{}_2 & 0 \\ 0 & e^2{}_1 & e^2{}_2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (50)$$

Here, we seek the solution for an electron in uniform motion along  $x^3$

$$\psi = \frac{N}{\sqrt{V}} \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix} \exp(-ik_\mu x^\mu) \quad k_\mu = (k_0, k_3) \quad (51)$$

The second term in the electron equation (32) is zero,<sup>2</sup> which leaves

$$i\gamma^\mu \partial_\mu \psi + \frac{1}{4} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\alpha{}^\lambda e_\gamma{}^\nu \partial_\nu e_{\beta\lambda} = 0 \quad (52)$$

In the present case,

$$\partial_0 \psi = -ik_0 \psi \quad \text{and} \quad \partial_3 \psi = -ik_3 \psi \quad (53)$$

so that

$$(k_0 \hat{\gamma}^0 + k_3 \hat{\gamma}^3) \psi + \frac{1}{4} \gamma_5 \hat{\gamma}_3 \psi \epsilon^{3ab0} e_a{}^l \partial_0 e_{bl} + \frac{1}{4} \gamma_5 \hat{\gamma}_0 \psi \epsilon^{0ab3} e_a{}^l \partial_3 e_{bl} = 0 \quad (54)$$

where the tetrad assumes the form (50). The Dirac matrix representation is

$$\hat{\gamma}^0 = \begin{pmatrix} \sigma_0 & 0 \\ 0 & -\sigma_0 \end{pmatrix} \quad \hat{\gamma}^a = \begin{pmatrix} 0 & \sigma_a \\ -\sigma_a & 0 \end{pmatrix} \quad \gamma_5 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (55)$$

---

<sup>2</sup>The trial solution (51) gives  $\partial_\mu \psi = -ik_\mu \psi$  and  $\partial_\mu \bar{\psi} = ik_\mu \bar{\psi}$ . From the conservation law (34), it follows that  $\bar{\psi} \partial_\mu (\sqrt{-g} \gamma^\mu) \psi = 0$ .

while  $\epsilon^{0123} = -1$ . Substitution yields four equations

$$k_0 u_1 + k_3 u_3 + \frac{1}{4} u_1 (e_1^n \partial_0 e_{2n} - e_2^n \partial_0 e_{1n}) + \frac{1}{4} u_3 (e_1^n \partial_3 e_{2n} - e_2^n \partial_3 e_{1n}) = 0 \quad (56)$$

$$k_0 u_2 - k_3 u_4 - \frac{1}{4} u_2 (e_1^n \partial_0 e_{2n} - e_2^n \partial_0 e_{1n}) + \frac{1}{4} u_4 (e_1^n \partial_3 e_{2n} - e_2^n \partial_3 e_{1n}) = 0 \quad (57)$$

$$k_3 u_1 + k_0 u_3 + \frac{1}{4} u_3 (e_1^n \partial_0 e_{2n} - e_2^n \partial_0 e_{1n}) + \frac{1}{4} u_1 (e_1^n \partial_3 e_{2n} - e_2^n \partial_3 e_{1n}) = 0 \quad (58)$$

$$k_3 u_2 - k_0 u_4 + \frac{1}{4} u_4 (e_1^n \partial_0 e_{2n} - e_2^n \partial_0 e_{1n}) - \frac{1}{4} u_2 (e_1^n \partial_3 e_{2n} - e_2^n \partial_3 e_{1n}) = 0 \quad (59)$$

These equations pair  $(u_1, u_3)$  or  $(u_2, u_4)$ . Consider the case  $u_2 = u_4 = 0$ : the equations will be satisfied for all values of  $u_1$  and  $u_3$ , if the coefficients are zero<sup>3</sup>

$$4k_0 + e_1^n \partial_0 e_{2n} - e_2^n \partial_0 e_{1n} = 0 \quad (60)$$

$$4k_3 + e_1^n \partial_3 e_{2n} - e_2^n \partial_3 e_{1n} = 0 \quad (61)$$

When expressed in terms of the  $\xi^\alpha_\mu$  (38), these equations read

$$4k_0 + (\xi^1_1 - \xi^2_2) \partial_0 \xi^1_2 - \xi^1_2 \partial_0 (\xi^1_1 - \xi^2_2) = 0 \quad (62)$$

$$4k_3 + (\xi^1_1 - \xi^2_2) \partial_3 \xi^1_2 - \xi^1_2 \partial_3 (\xi^1_1 - \xi^2_2) = 0 \quad (63)$$

The first order terms vanish, leaving only second order terms. A solution is given by

$$(\xi^1_1 - \xi^2_2) = 2a \cos(-k'_\mu x^\mu) \quad (64)$$

$$\xi^1_2 = a \sin(-k'_\mu x^\mu) \quad (65)$$

---

<sup>3</sup>For the case  $u_1 = u_3 = 0$ , the coefficients of  $u_2$  and  $u_4$  must be zero

$$4k_0 - e_1^n \partial_0 e_{2n} + e_2^n \partial_0 e_{1n} = 0$$

$$4k_3 - e_1^n \partial_3 e_{2n} + e_2^n \partial_3 e_{1n} = 0$$

Thus, the two cases are mutually exclusive.

where the amplitude  $a$  is small compared with 1. Substitution yields

$$k_0 = \frac{1}{2}a^2k'_0 \ll k'_0 \quad (66)$$

$$k_3 = \frac{1}{2}a^2k'_3 \ll k'_3 \quad (67)$$

Therefore, the electron's frequency and wave vector are much smaller than those of the gravitational field. Their phase velocities are equal,  $k_0/k_3 = k'_0/k'_3$ .

The spinor

$$\psi = \frac{N}{\sqrt{V}} \begin{pmatrix} u_1 \\ 0 \\ u_3 \\ 0 \end{pmatrix} \exp(-ik_\mu x^\mu) \quad k_\mu = (k_0, k_3) \quad (68)$$

has the form of a Dirac spinor for a positive energy electron of mass  $m$ , moving along  $x^3$ . Therefore, it will satisfy the Dirac equation

$$i\gamma^\mu \partial_\mu \psi - \frac{mc}{\hbar} \psi = 0 \quad (69)$$

By giving rise to such a solution, gravity creates inertia for the electron. Components  $u_1$  and  $u_3$  now satisfy

$$k_0 u_1 - k_3 u_3 - \frac{mc}{\hbar} u_1 = 0 \quad (70)$$

$$k_3 u_1 + k_0 u_3 + \frac{mc}{\hbar} u_3 = 0 \quad (71)$$

so that

$$u_3 = \frac{-\hbar k_3 u_1}{(\hbar k_0 + mc)} \quad \text{and} \quad \hbar^2 k_0^2 = \hbar^2 k_3^2 + m^2 c^2 \quad (72)$$

The final spinor is

$$\psi = \frac{N}{\sqrt{V}} \begin{pmatrix} 1 \\ 0 \\ \frac{-\hbar k_3}{(\hbar k_0 + mc)} \\ 0 \end{pmatrix} \exp(-ik_\mu x^\mu) \quad k_\mu = (k_0, k_3) \quad (73)$$

where

$$N^2 = \frac{\hbar k_0 + mc}{2\hbar k_0} \quad (74)$$

We turn now to the gravitational field equations (15), retaining only the largest terms in the Ricci tensor

$$R_{\mu\nu} \doteq \partial_\nu \Gamma_{\mu\lambda}^\lambda - \partial_\lambda \Gamma_{\mu\nu}^\lambda \quad (75)$$

$$\begin{aligned} \partial_\rho g_{\mu\nu} &= \eta_{\alpha\beta} \left( e^\alpha_\mu \partial_\rho e^\beta_\nu + e^\alpha_\nu \partial_\rho e^\beta_\mu \right) \\ &\doteq \eta_{\alpha\beta} \left( \delta^\alpha_\mu \partial_\rho \xi^\beta_\nu + \delta^\alpha_\nu \partial_\rho \xi^\beta_\mu \right) \\ &= 2 \partial_\rho \xi_{\mu\nu} \end{aligned} \quad (76)$$

It follows that

$$R_{\mu\nu} = \eta^{\lambda\rho} \{ \partial_\lambda \partial_\rho \xi_{\mu\nu} + \partial_\mu \partial_\nu \xi_{\lambda\rho} - \partial_\mu \partial_\lambda \xi_{\rho\nu} - \partial_\nu \partial_\lambda \xi_{\rho\mu} \} \quad (77)$$

The field equations are

$$\frac{\kappa\lambda^3}{V} \partial_3 \partial_3 (\xi^1_1 + \xi^2_2) + T_{00} = 0 \quad (78)$$

$$\frac{\kappa\lambda^3}{V} (\partial_0 \partial_0 - \partial_3 \partial_3) \xi^2_2 + T_{11} = 0 \quad (79)$$

$$\frac{\kappa\lambda^3}{V} (\partial_0 \partial_0 - \partial_3 \partial_3) \xi^1_1 + T_{22} = 0 \quad (80)$$

$$\frac{\kappa\lambda^3}{V} \partial_0 \partial_0 (\xi^1_1 + \xi^2_2) + T_{33} = 0 \quad (81)$$

$$\frac{\kappa\lambda^3}{V} (\partial_0 \partial_0 - \partial_3 \partial_3) \xi^1_2 - T_{12} = 0 \quad (82)$$

A length parameter  $\lambda$  is introduced together with the arbitrary volume of integration,  $V$ . The energy tensor is derived in appendix B (128). Make use of the spinor (73) and

$$(u_1^* u_1 + u_3^* u_3) = \frac{2\hbar k_0}{\hbar k_0 + mc} \quad (83)$$

$$(u_1^* u_3 + u_3^* u_1) = \frac{-2\hbar k_3}{\hbar k_0 + mc} \quad (84)$$

to find

$$T_{00} = \frac{\hbar ck_0}{V} + \frac{\hbar c}{4V}(e_1^n \partial_0 e_{2n} - e_2^n \partial_0 e_{1n}) \quad (85)$$

$$T_{11} = -\frac{\hbar c}{2V} \partial_0 e_{21} + \frac{\hbar ck_3}{2k_0 V} \partial_3 e_{21} \quad (86)$$

$$T_{22} = \frac{\hbar c}{2V} \partial_0 e_{12} - \frac{\hbar ck_3}{2k_0 V} \partial_3 e_{12} \quad (87)$$

$$T_{33} = \frac{\hbar ck_3^2}{k_0 V} + \frac{\hbar ck_3}{4k_0 V}(e_1^n \partial_3 e_{2n} - e_2^n \partial_3 e_{1n}) \quad (88)$$

$$T_{12} = -\frac{\hbar c}{4V} \partial_0(e_{11} - e_{22}) + \frac{\hbar ck_3}{4k_0 V} \partial_3(e_{11} - e_{22}) \quad (89)$$

Comparison with (60, 61) shows that  $T_{00} = T_{33} = 0$ ; also,  $T_{11} = -T_{22}$ . The field equations then give  $\xi^1_1 = -\xi^2_2$  so that two independent equations remain

$$\kappa\lambda^3(\partial_0\partial_0 - \partial_3\partial_3)\xi^1_1 - \frac{\hbar c}{2}\partial_0\xi^1_2 + \frac{\hbar ck_3}{2k_0}\partial_3\xi^1_2 = 0 \quad (90)$$

$$\kappa\lambda^3(\partial_0\partial_0 - \partial_3\partial_3)\xi^1_2 + \frac{\hbar c}{2}\partial_0\xi^1_1 - \frac{\hbar ck_3}{2k_0}\partial_3\xi^1_1 = 0 \quad (91)$$

These equations are satisfied by (64, 65), if

$$\kappa\lambda^3 = \frac{\hbar c}{2k'_0} \quad (92)$$

The gravitational field is given by

$$e^1_1 = 1 + a \cos(-k'_\mu x^\mu) \quad (93)$$

$$e^2_2 = 1 - a \cos(-k'_\mu x^\mu) \quad (94)$$

$$e^1_2 = e^2_1 = a \sin(-k'_\mu x^\mu) \quad (95)$$

where  $k'_\mu = (k'_0, k'_3)$ . The metrical determinant is constant  $\sqrt{-g} = 1 - a^2$  so that the condition  $\partial_\mu(\sqrt{-g}\gamma^\mu) = 0$  is satisfied (footnote 2).

Finally, we calculate the total energy (153, appendix C)

$$\begin{aligned} \mathcal{H} &= \frac{\kappa}{2} \left( \eta^{00} \Gamma_{m0}^l \Gamma_{l0}^m + \eta^{lm} \Gamma_{nl}^p \Gamma_{mp}^n \right) \\ &\quad - \frac{i}{2} \hbar c [\bar{\psi} \gamma^3 \partial_3 \psi - (\partial_3 \bar{\psi}) \gamma^3 \psi] - \frac{\hbar c}{4} \bar{\psi} \gamma_{\hat{5}} \hat{\gamma}_0 \psi \epsilon^{0ab3} e_3^3 e_{am} \partial_3 e_b^m \quad (96) \end{aligned}$$

The last two terms are equal to  $T_{33} = 0$ . Therefore, the energy is determined by the gravitational field

$$\begin{aligned} \mathcal{H} &= \frac{\kappa \lambda^3}{V} \left\{ (\partial_0 \xi_1^1)^2 + (\partial_3 \xi_1^1)^2 + (\partial_0 \xi_2^1)^2 + (\partial_3 \xi_2^1)^2 \right\} \\ &= \frac{\hbar c}{2k'_0 V} a^2 \left( k_0'^2 + k_3'^2 \right) \quad (97) \end{aligned}$$

Integrate over all of space to find

$$E = \int \mathcal{H} d^3x = \hbar \omega + \frac{\hbar c k_3^2}{k_0} \quad (98)$$

A constant energy must be subtracted in order to obtain  $E = \hbar \omega$ .

## Appendix A: Electron Lagrangian.

The scalar, 3-vector basis changes according to the formula

$$\nabla_\nu e_\mu = e_\lambda Q_{\mu\nu}^\lambda \quad (99)$$

Expanding  $e_\mu$  in terms of tetrads

$$e_\mu = e^\alpha{}_\mu \sigma_\alpha \quad (100)$$

we have

$$\begin{aligned} \nabla_\nu e_\mu &= \sigma_\alpha \partial_\nu e^\alpha{}_\mu + e^\alpha{}_\mu \nabla_\nu \sigma_\alpha \\ &= \sigma_\alpha \left( \partial_\nu e^\alpha{}_\mu + e^\beta{}_\mu \omega^\alpha{}_{\beta\nu} \right) \end{aligned} \quad (101)$$

where, by definition,<sup>4</sup>

$$\nabla_\nu \sigma_\alpha = \sigma_\beta \omega^\beta{}_{\alpha\nu} \quad (102)$$

Equate the two expressions (99) and (101) to find

$$e^\alpha{}_\lambda Q_{\mu\nu}^\lambda = \partial_\nu e^\alpha{}_\mu + e^\beta{}_\mu \omega^\alpha{}_{\beta\nu} \quad (103)$$

Contract this equation with  $e_{\alpha\rho}$  and form the tensor

$$\begin{aligned} g_{\rho\lambda} Q_{[\mu\nu]}^\lambda &= e_{\alpha\rho} (\partial_\nu e^\alpha{}_\mu - \partial_\mu e^\alpha{}_\nu) + e_{\alpha\rho} (e^\beta{}_\mu \omega^\alpha{}_{\beta\nu} - e^\beta{}_\nu \omega^\alpha{}_{\beta\mu}) \\ &= e^\alpha{}_\rho (\partial_\nu e_{\alpha\mu} - \partial_\mu e_{\alpha\nu}) + \omega_{\rho\mu\nu} - \omega_{\rho\nu\mu} \end{aligned} \quad (104)$$

From (9), it follows that the totally anti-symmetric tensor

$$Q_{[\mu\nu\lambda]} = g_{\mu\rho} Q_{[\nu\lambda]}^\rho + g_{\nu\rho} Q_{[\lambda\mu]}^\rho + g_{\lambda\rho} Q_{[\mu\nu]}^\rho = 0 \quad (105)$$

Therefore,

$$0 = e^\alpha{}_{[\rho} \partial_\nu e_{\alpha\mu]} + \omega_{[\rho\mu\nu]} \quad (106)$$

---

<sup>4</sup> The  $\omega^\alpha{}_{\beta\nu}$  give the change of orientation of the orthonormal basis  $\sigma_\alpha$  from point to point. Since  $\eta_{\alpha\beta} = \frac{1}{2}(\sigma_\alpha \tilde{\sigma}_\beta + \sigma_\beta \tilde{\sigma}_\alpha)$  where  $\tilde{\sigma}_\alpha = (\sigma_0, -\sigma_a)$  and  $\nabla_\nu \eta_{\alpha\beta} = \partial_\nu \eta_{\alpha\beta} = 0$ , we have  $\omega_{\alpha\beta\nu} = -\omega_{\beta\alpha\nu}$ . Moreover,  $\sigma_0 \sigma_a = \sigma_a$  implies that  $\nabla_\nu \sigma_0 = 0$  or  $\omega_{a0\nu} = 0$ . This leaves 12 parameters  $\omega_{ab\nu}$ . They comprise 3 rotation parameters along each of the four coordinates  $x^\nu$ .

or

$$\omega_{[\mu\nu\lambda]} = e_{[\mu}^{\alpha} \partial_{\nu} e_{\alpha\lambda]} \quad (107)$$

where

$$\omega_{[\mu\nu\lambda]} \equiv \frac{1}{6} \left( \omega_{\mu\nu\lambda} + \omega_{\nu\lambda\mu} + \omega_{\lambda\mu\nu} - \omega_{\nu\mu\lambda} - \omega_{\mu\lambda\nu} - \omega_{\lambda\nu\mu} \right) \quad (108)$$

The covariant spinor derivative is [7, 8]

$$D_{\mu} \psi = \partial_{\mu} \psi + iqA_{\mu} \psi + \Gamma_{\mu} \psi \quad (109)$$

where

$$\Gamma_{\mu} = \frac{1}{8} \left( \hat{\gamma}^{\alpha} \hat{\gamma}^{\beta} - \hat{\gamma}^{\beta} \hat{\gamma}^{\alpha} \right) \omega_{\alpha\beta\mu} = \frac{1}{4} \hat{\gamma}^{[\alpha} \hat{\gamma}^{\beta]} \omega_{\alpha\beta\mu} \quad (110)$$

We have included the U(1) term  $iqA_{\mu} \psi$ . The conjugate expression is

$$D_{\mu} \bar{\psi} = \partial_{\mu} \bar{\psi} - iqA_{\mu} \bar{\psi} - \bar{\psi} \Gamma_{\mu} \quad (111)$$

giving the electron Lagrangian

$$\begin{aligned} L_e &= \frac{i}{2} \hbar c [\bar{\psi} \gamma^{\mu} D_{\mu} \psi - (D_{\mu} \bar{\psi}) \gamma^{\mu} \psi] \\ &= \frac{i}{2} \hbar c [\bar{\psi} \gamma^{\mu} \partial_{\mu} \psi - (\partial_{\mu} \bar{\psi}) \gamma^{\mu} \psi] - \hbar c q \bar{\psi} \gamma^{\mu} \psi A_{\mu} \\ &\quad + \frac{i}{2} \hbar c \bar{\psi} (\gamma^{\mu} \Gamma_{\mu} + \Gamma_{\mu} \gamma^{\mu}) \psi \end{aligned} \quad (112)$$

The gravitational coupling term can be expressed in terms of the tetrad field as follows:

$$\begin{aligned} \gamma^{\mu} \Gamma_{\mu} + \Gamma_{\mu} \gamma^{\mu} &= \frac{1}{4} e_{\gamma}^{\mu} \left( \hat{\gamma}^{\gamma} \hat{\gamma}^{[\alpha} \hat{\gamma}^{\beta]} + \hat{\gamma}^{[\alpha} \hat{\gamma}^{\beta]} \hat{\gamma}^{\gamma} \right) \omega_{\alpha\beta\mu} \\ &= \frac{1}{2} \hat{\gamma}^{[\alpha} \hat{\gamma}^{\beta} \hat{\gamma}^{\gamma]} \omega_{\alpha\beta\gamma} \end{aligned} \quad (113)$$



where we have used the identity [8]

$$\hat{\gamma}^{[\gamma\hat{\gamma}^\alpha\hat{\gamma}^\beta]} \equiv \frac{1}{2} \left( \hat{\gamma}^\gamma \hat{\gamma}^{[\alpha\hat{\gamma}^\beta]} + \hat{\gamma}^{[\alpha\hat{\gamma}^\beta]} \hat{\gamma}^\gamma \right) \quad (114)$$

The identity [8]

$$\hat{\gamma}^{[\alpha\hat{\gamma}^\beta\hat{\gamma}^\gamma]} \equiv -i\gamma_5 \hat{\gamma}_\delta \epsilon^{\delta\alpha\beta\gamma} \quad (115)$$

yields

$$\begin{aligned} \gamma^\mu \Gamma_\mu + \Gamma_\mu \gamma^\mu &= -\frac{i}{2} \gamma_5 \hat{\gamma}_\delta \epsilon^{\delta\alpha\beta\gamma} \omega_{\alpha\beta\gamma} \\ &= -\frac{i}{2} \gamma_5 \hat{\gamma}_\delta \epsilon^{\delta\alpha\beta\gamma} \omega_{[\alpha\beta\gamma]} \end{aligned} \quad (116)$$

since  $\epsilon^{\delta\alpha\beta\gamma}$  is totally anti-symmetric. Substitute (107) in order to obtain

$$\gamma^\mu \Gamma_\mu + \Gamma_\mu \gamma^\mu = -\frac{i}{2} \gamma_5 \hat{\gamma}_\delta \epsilon^{\delta\alpha\beta\gamma} e_\alpha^\lambda e_\gamma^\nu \partial_\nu e_{\beta\lambda} \quad (117)$$

and the Lagrangian

$$\begin{aligned} L_e &= \frac{i}{2} \hbar c [\bar{\psi} \gamma^\mu \partial_\mu \psi - (\partial_\mu \bar{\psi}) \gamma^\mu \psi] - \hbar c q \bar{\psi} \gamma^\mu \psi A_\mu \\ &\quad + \frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\alpha^\lambda e_\gamma^\nu \partial_\nu e_{\beta\lambda} \end{aligned} \quad (118)$$

## Appendix B: Electron Energy Tensor.

The electron energy tensor is found by varying the tetrad field  $e_\alpha^\mu$

$$\begin{aligned}
\delta \int \mathcal{L}_e d^4x &= \int \left( \frac{\partial \mathcal{L}_e}{\partial e_\alpha^\mu} \delta e_\alpha^\mu + \frac{\partial \mathcal{L}_e}{\partial (\partial_\lambda e_\alpha^\mu)} \delta \partial_\lambda e_\alpha^\mu \right) d^4x \\
&= \int \left[ \frac{\partial \mathcal{L}_e}{\partial e_\alpha^\mu} - \partial_\lambda \frac{\partial \mathcal{L}_e}{\partial (\partial_\lambda e_\alpha^\mu)} \right] \delta e_\alpha^\mu d^4x \\
&= \int \sqrt{-g} A_{\mu\nu} e^{\beta\nu} \delta e_\beta^\mu d^4x
\end{aligned} \tag{119}$$

where

$$\sqrt{-g} A_{\mu\nu} \equiv e_{\alpha\nu} \left[ \frac{\partial \mathcal{L}_e}{\partial e_\alpha^\mu} - \partial_\lambda \frac{\partial \mathcal{L}_e}{\partial (\partial_\lambda e_\alpha^\mu)} \right] \tag{120}$$

The action is invariant under arbitrary rotations of the tetrad [7]. An infinitesimal rotation takes the form

$$\delta e_0^0 = 0 \quad \delta e_a^i = \epsilon_a^b e_b^i \quad \epsilon^{ab} = -\epsilon^{ba} \tag{121}$$

Thus,

$$\delta \int \mathcal{L}_e d^4x = \int \sqrt{-g} A_{ij} e_a^i e_b^j \epsilon^{ba} d^4x = 0 \tag{122}$$

Therefore, the antisymmetric part of  $A_{ij}$  must be zero

$$\frac{1}{2} (A_{ij} - A_{ji}) = 0 \tag{123}$$

and we define the symmetric part to be

$$T_{\mu\nu} = \frac{1}{2} (A_{\mu\nu} + A_{\nu\mu}) \tag{124}$$

It follows that, for an arbitrary variation,

$$\begin{aligned}
\delta \int \mathcal{L}_e d^4x &= \int \sqrt{-g} A_{\mu\nu} \frac{1}{2} (e^{\beta\mu} \delta e_\beta^\nu + e^{\beta\nu} \delta e_\beta^\mu) d^4x \\
&= \frac{1}{2} \int e_{\alpha\nu} \left[ \frac{\partial \mathcal{L}_e}{\partial e_\alpha^\mu} - \partial_\lambda \frac{\partial \mathcal{L}_e}{\partial (\partial_\lambda e_\alpha^\mu)} \right] \delta g^{\mu\nu} d^4x \\
&= \frac{1}{2} \int T_{\mu\nu} \delta g^{\mu\nu} \sqrt{-g} d^4x
\end{aligned} \tag{125}$$

Explicitly,

$$\begin{aligned} \frac{\partial \mathcal{L}_e}{\partial e_\alpha^\mu} &= \frac{i}{2} \hbar c [\bar{\psi} \gamma_\lambda \partial_\mu \psi - (\partial_\mu \bar{\psi}) \gamma_\lambda \psi] e^{\alpha\lambda} \sqrt{-g} - \hbar c q \bar{\psi} \gamma_\lambda \psi A_\mu e^{\alpha\lambda} \sqrt{-g} \\ &\quad + \frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\gamma^\lambda (\partial_\lambda e_{\beta\mu} - \partial_\mu e_{\beta\lambda}) \sqrt{-g} \end{aligned} \quad (126)$$

$$\frac{\partial \mathcal{L}_e}{\partial (\partial_\lambda e_\alpha^\mu)} = -\frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\gamma^\lambda e_{\beta\mu} \sqrt{-g} \quad (127)$$

which give

$$\begin{aligned} T_{\mu\nu} &= \frac{i}{4} \hbar c [\bar{\psi} \gamma_\mu \partial_\nu \psi + \bar{\psi} \gamma_\nu \partial_\mu \psi - (\partial_\mu \bar{\psi}) \gamma_\nu \psi - (\partial_\nu \bar{\psi}) \gamma_\mu \psi] \\ &\quad - \frac{1}{2} \hbar c q (\bar{\psi} \gamma_\mu \psi A_\nu + \bar{\psi} \gamma_\nu \psi A_\mu) \\ &\quad + \frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\gamma^\lambda [(e_{\alpha\mu} \partial_\lambda e_{\beta\nu} + e_{\alpha\nu} \partial_\lambda e_{\beta\mu}) \\ &\quad - \frac{1}{2} (e_{\alpha\mu} \partial_\nu e_{\beta\lambda} + e_{\alpha\nu} \partial_\mu e_{\beta\lambda})] \end{aligned} \quad (128)$$

The Lagrange form of the gravitational field equations can now be expressed in terms of tetrads. We first substitute  $g^{\mu\nu} = \eta^{\alpha\beta} e_\alpha^\mu e_\beta^\nu$  in  $\mathcal{L}_g$  to find [2]

$$\frac{\partial \mathcal{L}_g}{\partial g^{\mu\nu}} - \partial_\lambda \frac{\partial \mathcal{L}_g}{\partial (\partial_\lambda g^{\mu\nu})} = \frac{1}{2} e_{\alpha\nu} \left[ \frac{\partial \mathcal{L}_g}{\partial e_\alpha^\mu} - \partial_\lambda \frac{\partial \mathcal{L}_g}{\partial (\partial_\lambda e_\alpha^\mu)} \right] \quad (129)$$

This, together with the energy tensor for the electron,

$$T_{\mu\nu} = e_{\alpha\nu} \left[ \frac{\partial \mathcal{L}_e}{\partial e_\alpha^\mu} - \partial_\lambda \frac{\partial \mathcal{L}_e}{\partial (\partial_\lambda e_\alpha^\mu)} \right] \quad (130)$$

give the field equations

$$\frac{\partial \mathcal{L}}{\partial e_\alpha^\mu} - \partial_\lambda \frac{\partial \mathcal{L}}{\partial (\partial_\lambda e_\alpha^\mu)} = 0 \quad (131)$$

### Appendix C: Energy Conservation.

The principle of energy conservation derives from the Lagrange equations of motion, by means of the Hamilton function

$$\mathcal{H} = \sum_{\phi} c\pi \partial_0 \phi - \mathcal{L} \quad (132)$$

The fields  $\phi$  include  $e_{\alpha}^{\mu}$ ,  $\psi$ ,  $\bar{\psi}$ , and  $A_{\mu}$ , while the momenta are defined by

$$c\pi = \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi)} \quad (133)$$

The temporal change of  $\mathcal{H}$  is

$$\begin{aligned} \partial_0 \mathcal{H} &= \sum_{\phi} \partial_0 \left[ \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi)} \right] \partial_0 \phi + \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi)} \partial_0 \partial_0 \phi - \frac{\partial \mathcal{L}}{\partial \phi} \partial_0 \phi - \frac{\partial \mathcal{L}}{\partial(\partial_{\lambda} \phi)} \partial_0 \partial_{\lambda} \phi \\ &= \sum_{\phi} \partial_0 \left[ \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi)} \right] \partial_0 \phi - \frac{\partial \mathcal{L}}{\partial \phi} \partial_0 \phi - \frac{\partial \mathcal{L}}{\partial(\partial_n \phi)} \partial_0 \partial_n \phi \end{aligned} \quad (134)$$

Making use of

$$\partial_n \left[ \frac{\partial \mathcal{L}}{\partial(\partial_n \phi)} \partial_0 \phi \right] = \partial_n \left[ \frac{\partial \mathcal{L}}{\partial(\partial_n \phi)} \right] \partial_0 \phi + \frac{\partial \mathcal{L}}{\partial(\partial_n \phi)} \partial_n \partial_0 \phi \quad (135)$$

we find

$$\begin{aligned} \partial_0 \mathcal{H} &= \sum_{\phi} \left[ \partial_{\lambda} \frac{\partial \mathcal{L}}{\partial(\partial_{\lambda} \phi)} - \frac{\partial \mathcal{L}}{\partial \phi} \right] \partial_0 \phi - \partial_n \left[ \frac{\partial \mathcal{L}}{\partial(\partial_n \phi)} \partial_0 \phi \right] \\ &= \sum_{\phi} -\partial_n \left[ \frac{\partial \mathcal{L}}{\partial(\partial_n \phi)} \partial_0 \phi \right] \end{aligned} \quad (136)$$

The last step is by virtue of the Lagrange equations of motion. Integrate over all 3-dimensional space and discard surface terms, in order to obtain conservation of total energy

$$\frac{d}{dx^0} \int \mathcal{H} d^3x = 0 \quad (137)$$

We now derive an explicit expression for the energy density

$$\mathcal{H} = c\pi_{\mu}^{\alpha} \partial_0 e_{\alpha}^{\mu} + c\bar{\pi} \partial_0 \psi + \partial_0 \bar{\psi} c\pi + c\pi^{\mu} \partial_0 A_{\mu} - \mathcal{L} \quad (138)$$

The spinor momenta are

$$c\bar{\pi} = \frac{\partial \mathcal{L}}{\partial(\partial_0 \psi)} = \frac{i}{2} \hbar c \sqrt{-g} \bar{\psi} \gamma^0 \quad (139)$$

$$c\pi = \frac{\partial \mathcal{L}}{\partial(\partial_0 \bar{\psi})} = -\frac{i}{2} \hbar c \sqrt{-g} \gamma^0 \psi \quad (140)$$

The electromagnetic momenta follow from

$$L_{e-m} = -\frac{1}{4} \left( 2g^{00} g^{ij} F_{i0} F_{j0} + g^{il} g^{jm} F_{ij} F_{lm} \right) \quad (141)$$

$$\frac{\partial F_{j0}}{\partial(\partial_0 A_i)} = -\delta_j^i \quad (142)$$

Therefore,

$$c\pi^0 = \frac{\partial \mathcal{L}}{\partial(\partial_0 A_0)} = 0 \quad (143)$$

$$c\pi^i = \frac{\partial \mathcal{L}}{\partial(\partial_0 A_i)} = \sqrt{-g} F^{i0} \quad (144)$$

Turning now to the tetrad momenta, the gravitational Lagrangian

$$\begin{aligned} L_g &= \frac{\kappa}{2} g^{\mu\nu} \left( \Gamma_{\mu\nu}^\lambda \Gamma_{\rho\lambda}^\rho - \Gamma_{\rho\nu}^\lambda \Gamma_{\mu\lambda}^\rho \right) \\ &= \frac{\kappa}{2} \left\{ g^{00} \left( \Gamma_{m0}^l \Gamma_{l0}^m - \Gamma_{l0}^l \Gamma_{m0}^m \right) + g^{lm} \left( \Gamma_{lm}^n \Gamma_{0n}^0 - \Gamma_{0l}^0 \Gamma_{pm}^p \right) \right. \\ &\quad \left. + g^{lm} \left( \Gamma_{lm}^n \Gamma_{pn}^p - \Gamma_{nl}^p \Gamma_{mp}^n \right) \right\} \end{aligned} \quad (145)$$

contains time derivatives in the first term

$$\frac{\partial \Gamma_{m0}^l}{\partial(\partial_0 g^{ij})} = -\frac{1}{2} g_{im} \delta_j^l \quad \frac{\partial \Gamma_{l0}^l}{\partial(\partial_0 g^{ij})} = -\frac{1}{2} g_{ij} \quad (146)$$

Therefore,

$$\frac{\partial \mathcal{L}_g}{\partial(\partial_0 g^{ij})} = -\frac{\kappa}{2} \sqrt{-g} g^{00} \left( g_{il} \Gamma_{j0}^l - g_{ij} \Gamma_{l0}^l \right) \quad (147)$$

The electron Lagrangian (28) contains time derivatives in the coupling term

$$\begin{aligned}\frac{\partial \mathcal{L}_e}{\partial(\partial_0 e_\beta^\mu)} &= \frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta\gamma} e_\gamma^0 e_{\alpha\mu} \sqrt{-g} \\ &= \frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_\delta \psi \epsilon^{\delta\alpha\beta 0} e_0^0 e_{\alpha\mu} \sqrt{-g}\end{aligned}\quad (148)$$

which leaves

$$\frac{\partial \mathcal{L}_e}{\partial(\partial_0 e_a^i)} = -\frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_d \psi \epsilon^{dab0} e_0^0 e_{bi} \sqrt{-g}\quad (149)$$

The tetrad momenta are given by

$$\begin{aligned}c\pi_\mu^\alpha &= \frac{\partial \mathcal{L}}{\partial(\partial_0 e_\alpha^\mu)} \\ &= \frac{\partial \mathcal{L}_g}{\partial(\partial_0 g^{\nu\lambda})} \frac{\partial(\partial_0 g^{\nu\lambda})}{\partial(\partial_0 e_\alpha^\mu)} + \frac{\partial \mathcal{L}_e}{\partial(\partial_0 e_\alpha^\mu)} \\ &= \frac{\partial \mathcal{L}_g}{\partial(\partial_0 g^{\nu\lambda})} \left( \delta_\mu^\nu e^{\alpha\lambda} + \delta_\mu^\lambda e^{\alpha\nu} \right) + \frac{\partial \mathcal{L}_e}{\partial(\partial_0 e_\alpha^\mu)}\end{aligned}\quad (150)$$

It follows that

$$c\pi_0^0 = \frac{\partial \mathcal{L}}{\partial(\partial_0 e_0^0)} = 0\quad (151)$$

$$\begin{aligned}c\pi_i^a &= \frac{\partial \mathcal{L}}{\partial(\partial_0 e_a^i)} \\ &= -\kappa \sqrt{-g} g^{00} \left( e_i^a \Gamma_{i0}^l - e_i^a \Gamma_{l0}^i \right) - \frac{\hbar c}{4} \bar{\psi} \gamma_5 \hat{\gamma}_d \psi \epsilon^{dab0} e_0^0 e_{bi} \sqrt{-g}\end{aligned}\quad (152)$$

Substituting the momenta into (138), we find the total energy density to be

$$\begin{aligned}
\mathcal{H} = & \frac{\kappa}{2}\sqrt{-g}\left\{g^{00}\left(\Gamma_{m0}^l\Gamma_{l0}^m - \Gamma_{l0}^l\Gamma_{m0}^m\right) - g^{lm}\left(\Gamma_{lm}^n\Gamma_{0n}^0 - \Gamma_{0l}^0\Gamma_{pm}^p\right)\right. \\
& \left. - g^{lm}\left(\Gamma_{lm}^n\Gamma_{pn}^p - \Gamma_{nl}^p\Gamma_{mp}^n\right)\right\} \\
& + \frac{1}{2}\sqrt{-g}g^{00}g^{lm}\left(\partial_l A_0\partial_m A_0 - \partial_0 A_l\partial_0 A_m\right) + \frac{1}{4}\sqrt{-g}g^{il}g^{jm}F_{ij}F_{lm} \\
& - \frac{i}{2}\hbar c\left[\bar{\psi}\gamma^l\partial_l\psi - (\partial_l\bar{\psi})\gamma^l\psi\right]\sqrt{-g} + \hbar c q\bar{\psi}\gamma^\mu\psi A_\mu\sqrt{-g} \\
& - \frac{\hbar c}{4}\bar{\psi}\gamma_5\hat{\gamma}_0\psi\epsilon^{0abc}e_c{}^le_{am}\partial_l e_b{}^m\sqrt{-g}
\end{aligned} \tag{153}$$

## References.

1. K. Dalton, "Electromagnetism and Gravitation," *Hadronic Journal* **17** (1994) 483; [www.arxiv.org/gr-qc/9512027](http://www.arxiv.org/gr-qc/9512027).
2. K. Dalton, "Gravity and the Dirac Electron," [www.arxiv.org/physics/0409042](http://www.arxiv.org/physics/0409042).
3. L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields* (Pergamon, 1975).
4. T-P Cheng and L-F Li, *Gauge Theory of Elementary Particle Physics* (Oxford, 1984).
5. M. Guidry, *Gauge Field Theories* (Wiley, 1991).
6. W. Rolnick, *The Fundamental Particles and Their Interactions* (Addison-Wesley, 1994).
7. S. Weinberg, *Gravitation and Cosmology* (Wiley, 1972).
8. V. de Sabbata and M. Gasperini, *Introduction to Gravitation* (World Scientific, 1985).