

# Poincaré's relativistic Doppler-Fizeau formula

Dr. Yves Pierseaux  
ULB ypiersea@ulb.ac.be

August 28, 2018

## Abstract

We deduce from Poincaré's ellipsoidal wavefronts a relativistic Doppler-Fizeau formula that is not the same as 1905 Einstein's one. Longitudinally, Einstein's formula and Poincaré's formula are the same. Both formulas are compatible with relativistic time dilation. The question of an experimental test is directly connected with the possibility or the impossibility of directly measuring the transverse Doppler effect. Hasselkamp's experiment (1978) becomes a crucial experiment because Poincaré's relativistic kinematics predicts an expansion of space directly connected with the Doppler-Fizeau effect for the remote objects.

## 1 Lorentz transformation and Poincaré's elliptical wavefront

Let us consider a *couple* of inertial systems K and K' in uniform translation relative one another. A source of light is at rest O in K. What is the image by LT in K', of a circular wavefront in K, emitted in  $t' = t = 0$  by this source S? Poincaré writes LT with perfectly spacetime symmetry (in x and t) [Poincaré H. 1905]:

$$x' = k(x + \varepsilon t) \quad y' = y \quad t' = k(t + \varepsilon x) \quad (1)$$

Poincaré's notations where  $\varepsilon, k$  correspond to Einstein-Planck's notations  $\beta, \gamma$  given that, according to Poincaré in his 1905 work about the theory of relativity, "*I shall choose the units of length and of time in such a way that the velocity of light is equal to unity*" [Pierseaux Y (2004)]. For the easiness of present reader we adopt standard notations  $\beta, \gamma$  but we keep Poincaré's spacetime units  $c = 1$ :

$$x' = \gamma(x + \beta t) \quad y' = y \quad t' = \gamma(t + \beta x) \quad (2)$$

Let us consider the relativistic invariant:

$$x^2 + y^2 = r_0^2 = t_0^2 \quad x'^2 + y'^2 = r'^2 = t'^2 \quad (3)$$

The object time  $t = t_0$  is fixed in K (circular wavefront in K) but image time  $t'$  is not fixed (by LT) in K'. We obtain with (2)  $t' = \gamma^{-1}t_0 + \beta x'$ ,

$$x'^2 + y'^2 = (\gamma^{-1}t_0 + \beta x')^2 \quad \text{ou} \quad (\gamma^{-1}x' - \beta t_0)^2 + y'^2 = t_0^2 \quad (4)$$

the Cartesian equation of an elongated ellipse in K' (the observer O' is at the focus, **figure 1**). Poincaré's ellipse is engraved in LT and its physical meaning is the **relativity of simultaneity**: two simultaneous events (*different abscissa*) in K are not simultaneous in K' [Moreau W.] & [Pierseaux Y.(ellipse)]:

$$\Delta t = 0 \quad \rightarrow \quad \Delta t' = \frac{t'^+ - t'^-}{2} \neq 0 = \gamma \beta x \quad (5)$$

$\Delta t'$  is "the gap of simultaneity" between two opposite events on the front (**figure 1**). In order to have a circular image<sup>1</sup> of the wavefront we must fix the time  $t'$  :

$$(\Delta t' = 0)_{\text{Einstein}} \quad (5\text{bis})$$

---

<sup>1</sup>If according to Einstein, the object (fixed time t) and the image (fixed time t'), are both spherical within the two systems, then two simultaneous events in K must be always also simultaneous in K'. [Einstein A.(1905), paragraphe 3)]. This is Einstein's convention of synchronisation in spherical waves ( $\Delta t = \Delta t' = 0$ )[Pierseaux Y.(ellipse)].

Poincaré's ellipse in polar coordinates ( $x' = r' \cos \theta'$ ,  $y' = r' \sin \theta'$  with  $\theta'$  as polar angle and F as pole) is :

$$r' = \frac{r_0}{\gamma(1 - \beta \cos \theta')} \quad (6)$$

With relativistic transformation of angle

$$\cos \theta' = \frac{\cos \theta + \beta}{1 + \beta \cos \theta} \quad (7)$$

we rediscover polar LT for any point of the wavefront ( $r, \theta$  or  $t, \theta$ ) :

$$r' = \gamma r_0(1 + \beta \cos \theta) \quad t' = \gamma t_0(1 + \beta \cos \theta) \quad (8)$$

that are the radial or the temporal equation of an ellipse. The LT of a circular wavefront into a elliptical wavefront has a 4-vectorial interpretation:

$$(r \cos \theta, r \sin \theta, 0, t) \quad \rightarrow \quad (r' \cos \theta', r' \sin \theta', 0, t') \quad (9)$$

Light-like unprimed 4-vector (null interval) is transformed into light-like primed 4-vector (null interval) (6)

$$r' \cos \theta' = \gamma(r \cos \theta + \beta t) \quad r' \sin \theta' = r \sin \theta \quad t' = \gamma(t + \beta r \cos \theta) \quad (10)$$

with the invariance of the quadratic form (the interval of events, see below):

$$r^2 - t^2 = 0 \quad \rightarrow \quad r'^2 - t'^2 = 0 \quad (11)$$

The transformed (primed) 4-vector describes an *isotropic ellipse* in the meaning that the velocity of light ("one way") is identical in all directions within both systems because we have by construction **figure 1**):

$$\frac{r_0}{t_0} = \frac{r'^+}{t'^+} = \frac{r'^-}{t'^-} = c = 1 \quad (12)$$

What is a wavefront in a relativistic sense (a "spacetime" wavefront)? The invariant interval between the event "emission" ( $t = t' = 0$ ) and any event on the wavefront ( $t = t_0, r = r_0$  in K and  $t', r'$  in K') is null within both systems. So Minkowski's spacetime 4-vector (with null interval) is, in Poincaré's kinematics, a **wavefront 4-vector (9)**. *The elliptical image by LT is deduced from a non-transversal  $t'$  section in Minkowski's cone<sup>2</sup>.* The essential fact that Poincaré's ellipse is engraved in LT can be seen with the inverse LT:

$$x = \gamma(x' - \beta t') \quad y = y' \quad t = \gamma(t' - \beta x') \quad (13)$$

We obtain indeed immediately the temporal equation of the inverse ellipse (**figure 2**) (here polar equations, from 6  $r = t$  and  $r' = t'$ ):

$$\left[ t' = \frac{t}{\gamma(1 - \beta \cos \theta')} \right]_{\text{direct ellipse}} \quad \left[ t = \frac{t'}{\gamma(1 + \beta \cos \theta)} \right]_{\text{inverse ellipse}} \quad (14)$$

Inverse elongated ellipse (14bis), with the source at the focus and the observer O' at the centre, is Poincaré's *historical ellipse (1906-1912)*. This ellipse explains, according to Poincaré, the null result of Michelson experiment (the mean time back and forth is the same in any direction, see Poincaré's oscillator [Pierseaux Y (2004)]). Let us however here focus the attention on the fact that the relation between t and t' is exactly the same if we consider the "source at rest" (14) or the "observer at rest" (14bis). Suppose now that the source emits a light signal with a period of T seconds. Given that the relativistic kinematics is a classical theory, the period of the emitted wave is also T seconds. The periodical wave that consists of a series of elliptical wavefronts received by the observer defines the standard configuration of a Doppler-Fizeau effect (with the frequency  $\nu = \frac{1}{T}$  and  $\nu' = \frac{1}{T'}$ ). In prerelativistic point of view the two cases are not symmetrical. In relativistic (elliptical) point of view, the respective role of the source and the observer are completely interchangeable (see Poincaré's wavefront 4-vector). However, in order to have a rigorous deduction of a Doppler-Fizeau formula (emission in a *constant* direction), we have to consider a plane wave ([Pierseaux Y.(ellipse)]).

<sup>2</sup>Poincaré's spacetime metrics is not the same as Einstein-Minkowski's spacetime metrics ([Pierseaux Y (2004)]).

## 2 Tangent to the ellipse, plane wave and Poincaré's Doppler formula

Until now we considered one only source  $S_O$  which emits at  $t = t' = 0$  a spherical wave with fixed time  $t = t_0$  (figure 1)- Let us now consider a second source  $S_\infty$  at infinity at rest in K in the direction  $\theta$  (**figure 3**) which emits a plane wavefront.

Suppose that the considered plane wavefront (here a "wave straight line" at two space dimensions) be in O at the time  $t = t' = 0$  (when  $S_O$  emits a spherical wave). It will be **tangent** in  $t = t_0$  to the circular wavefront, emitted by  $S_1$  (normalized  $t_0 = r_0 = 1$  **figure 3**). We note that the simultaneous events  $P_1TP_2$  in K are no longer simultaneous ( $P'_1T'P'_2$ ) in K'. We deduce respectively (2) for the object front ("wave straight line" with angular coefficient  $a = -\cot g \theta$ ) and the image front both following relations:

$$x \cos \theta + y \sin \theta = t_0 \quad y' \sin \theta' + x' \cos \theta' = t' \quad (15)$$

Exactly like the circular wavefronts (équation 3), there are two possibilities:

**If  $t'$  is not fixed (Poincaré, spacetime image wavefront)**, we have by LT,  $t' = \gamma(t + \beta x) = \gamma^{-1}t + \beta x'$ , and therefore the primed relation 15 is the equation of the tangent to the ellipse at the point  $T'$  (on which  $P'_1$  and  $P'_2$  are situated) :

$$y' \sin \theta' + x'(\cos \theta' - \beta) = \gamma^{-1}t \quad (16)$$

The angular coefficient of Poincaré's wavefront is  $\theta'$ :

$$a'_{poincaré} = tg\alpha' = \frac{\beta - \cos \theta'}{\sin \theta'} \quad (17)$$

**If  $t'$  is fixed (Einstein, space image wavefront<sup>3</sup>)** the primed equation 15 is the equation of the tangent to the circle (with centre O' and radius  $r'_T$ ). The angular coefficient of Einstein's wavefront is:

$$a'_{einstein} = -\cot g\theta' \quad (18)$$

Both formulas are the same for  $\theta' = 0$ . **Einstein's double transversality<sup>4</sup>** (or **Einstein's double simultaneity**) involves for the image front:

$$a' = -\cot g\theta' \quad \Leftrightarrow \quad (\Delta t')_{front} = 0 \quad (19)$$

This is consistent with Einstein's synchronisation (5bis). The phase  $\Psi$  of a sinusoidal monochromatic plane wave ( $\mathbf{A}$  is the amplitude)  $\mathbf{A} = \mathbf{A}_0 \sin \Psi$  ( $\mathbf{A}' = \mathbf{A}'_0 \sin \Psi'$ ) is defined by the **3-scalar product  $\mathbf{k} \cdot \mathbf{r}$**  ( $\mathbf{k}' \cdot \mathbf{r}'$ ) with the frequency  $\nu = \frac{\omega}{2\pi}$ , the wave vector  $\mathbf{k} = \frac{2\pi}{\lambda} \mathbf{1}_n$  with  $\mathbf{1}_n$  the unit vector normal to the front ( $\mathbf{k}' = \frac{2\pi}{\lambda'} \mathbf{1}_{n'}$ ,  $\nu' = \frac{\omega'}{2\pi}$ ) :

$$\omega t - \mathbf{k} \cdot \mathbf{r} = \Psi = \omega' t' - \mathbf{k}' \cdot \mathbf{r}' = \Psi' \quad \Leftrightarrow \quad \mathbf{A} \cdot \mathbf{k} = \mathbf{A}' \cdot \mathbf{k}' = 0 = Ak \cos \phi = A'k' \cos \phi' \quad (20)$$

This is a **Galilean invariant** (t is fixed on the object front and t' is fixed on the image front) in the meaning where the angle (**figure 3**)  $\phi = \phi' = 90^\circ$  is not altered by Galilean transformation (GT).

$$\mathbf{k} \cdot \mathbf{r} \quad \underline{TG} \quad \mathbf{k}' \cdot \mathbf{r}' \quad \text{ou} \quad \mathbf{A} \perp \mathbf{k} \quad \underline{TG} \quad \mathbf{A}' \perp \mathbf{k}' \quad (20bis)$$

But LT changes (17) this angle  $\phi = 90^\circ$  into  $\phi'$  in the following way (**figure 3**):

$$\tan \phi' = \tan(\alpha' - \theta') = \frac{\beta \cos \theta' - 1}{\beta \sin \theta'} \quad (17bis)$$

The right angle  $\phi = \phi' = 90^\circ$  is conserved *if and only if* the propagation is purely longitudinal  $\sin \theta' = 0$  or "everything happens as if  $\beta = 0$ ", in other words exactly as in Einstein's synchronisation in spherical waves (5bis, note1). Einstein considers [Einstein A.(1905), paragraphe 7] that the Galilean invariant (20) is *by definition* a Lorentz invariant. Einstein's definition of the invariance of the phase is therefore based on a **4-scalar product** between the 4-vector position  $(t, \mathbf{r})$  and another 4-vector with null norm  $(\omega, \mathbf{k})$  :

$$(\omega, \mathbf{k}) \cdot (t, \mathbf{r}) = \omega t - \mathbf{k} \cdot \mathbf{r} = \Psi \quad \nu(t - x \cos \theta - y \sin \theta) = \nu'(t' - x' \cos \theta' - y' \sin \theta') \quad (21)$$

<sup>3</sup>The classical rigidity is incompatible with LT: a set of simultaneous events cannot be remain by LT a set of simultaneous events. Einstein's rigid rods are consistent with Einstein's rigid front wave.

<sup>4</sup>Einstein defined clearly the image of the wavefront in paragraph 7 of his 1905 paper [Einstein A.(1905)]: "If we call the angle  $\theta'$  the angle between **the wave-normal (direction of the ray)** and the "direction of movement"."

On the basis of Einstein's wave 4-vector (with two space dimensions) Einstein deduces from the covariance of scalar product  $\mathbf{k} \cdot \mathbf{r}$  a Doppler-Fizeau formula for a monochromatic sinusoidal wave plane:

$$\left(\frac{\nu'}{\nu}\right)_{\text{Einstein}} = \gamma(1 - \beta \cos \theta) \quad (22)$$

Einstein's extension of prerelativistic concept of wave **3-vector** (double transversality) into a wave **4-vector**  $(\omega, \mathbf{k})$  is a very serious problem ( $t'$  is fixed on the front but  $r'$  is not fixed) because the LT is not compatible with the double transversality (double simultaneity).

Suppose now that the source  $S_2$  (at the infinity, **figure 3**) emits a series of plane wavefronts (here a train of "wave straight line") at the same *rhythm* as the spherical wavefronts ( $\nu = \frac{1}{T} = \frac{1}{t}$ ). Let us construct a Lorentz invariant phase  $\Phi$  on the only basis of Lorentz invariance of the interval,  $(t-r)(t+r) = (t'-r'_T)(t'+r')$ , between the event  $t = t' = 0$  and any event on the plane front ([Pierseaux Y.(ellipse)]). With the tangency point T' of the front to the ellipse (image of T, **figure 3**) we have  $(t-r_T)(t+r_T) = (t'-r'_{T'})(t'+r'_{T'})$ :

$$\nu(t-r_T) = \Phi = \nu'(t'-r'_{T'}) = \Phi' \quad \text{or} \quad \omega t - kr_T = \Phi = \omega' t' - k' r'_{T'} = \Phi' \quad (23)$$

Poincaré's phase invariant  $\Phi = \Phi'$ , which is constructed with fundamental quadratic invariant (3), is necessarily a solution of the second order wave equation of electromagnetism ([Pierseaux Y. (2005-2)] & [Pierseaux Y & Rousseaux G]). The invariance of the phase for the wavefront is no longer coupled with the (prerelativistic) fixing of the time on the wavefront in each system. We immediately deduce from (23) a relativistic Doppler-Fizeau formula:

$$\left(\frac{\nu'}{\nu}\right)_{\text{Poincaré}} = \gamma(1 - \beta \cos \theta') \quad (24)$$

Let us call this formula, deduced from Poincaré's ellipse, "Poincaré's Doppler-Fizeau formula". **This is not the same as Einstein's one** (22) because the proper angle  $\theta$  is replaced with improper angle  $\theta'$  (see conclusion 29 for **experimental tests**). For the longitudinal propagation, the formulas are the same:

$$\left(\frac{\nu'}{\nu}\right)_{\text{Einstein-Poincaré}} = \sqrt{\frac{1-\beta}{1+\beta}} = \gamma(1-\beta) \quad (25)$$

Poincaré's formula can be immediately deduced from (14), with  $\nu = \frac{1}{t}$  et  $\nu' = \frac{1}{t'}$ , and therefore for polar LT (8), supposes the existence of a **wave 4-vector** *directly proportional* to the wavefront 4-vector defined in the paragraph 1 ( $c \neq 1$ ) (9)

$$(\lambda \cos \theta, \lambda \sin \theta, 0, cT)_{\text{poincaré}} \quad \left(\frac{1}{\lambda} \cos \theta, \frac{1}{\lambda} \sin \theta, 0, \frac{\nu}{c}\right)_{\text{einstein}} \quad (26)$$

Poincaré's 4-vector involves that the frequency (24) is transformed by LT *like the inverse of a the time*  $\nu = \frac{1}{t}$  et  $\nu' = \frac{1}{t'}$  whilst Einstein's four vector<sup>5</sup> involves that frequency (22) is transformed by LT *like the time* (see de Broglie's works). Poincaré's proportionality (9-26) involves that this Doppler-Fizeau formula for remote objects ( $S_\infty$ ) is not independant of the structure of space-time (Poincaré's **expansion of space** for remote objects [Pierseaux Y.(ellipse)]). We showed [Pierseaux Y & Rousseaux G.] that Poincaré's theory of wavefront is compatible with electromagnetic theory of waves (in particular with the transversality of the primed electric field to the primed direction of propagation).

### 3 Conclusion: Doppler formulas and experimental tests

In order to experimentally test both Doppler formulas, which are both compatible with the relativistic dilation of time [Pierseaux Y (2004)], we need length waves (or frequencies) emitted by moving atoms with respect to the observer (the relativistic transformation of the energy is not in question here). Mandelberg describes the SODS ("Second Order Doppler-Shift"):

<sup>5</sup>In Einstein-Minkowski's kinematics, the light has a double 4-vectorial representation : the Minkowski's light like four-vector and Einstein's wave 4-vector. This latter is inseparable du photon d'Einstein, défini en jauge transverse [Pierseaux Y. (2005-2)].

A moving beam of radiating hydrogen atoms with velocities ranging up to  $2.8 \cdot 10^8 \text{ cm/s}$  had been viewed from the incoming and the outgoing directions simultaneously. Averaging wavelength measurements of a particular spectral line for the two observations give a measurement of the quadratic shift... The wavelength of the light emitted from an oncoming atom of velocity  $v$  viewed at a small angle  $\theta'$  (near zero) to the beam direction by an observer stationary in the laboratory reference frame is given by (1) where  $\theta'$  is measured in the laboratory frame and  $\lambda$  is the wavelength observed in a reference frame at rest with respect to the radiating atom. The  $\beta \cos \theta$  term in the numerator is the first-order Doppler shift. The subscript B indicates a *Doppler shift to the blue*. The denominator contains the quadratic shift; the approximation expression is valid in this experiment because of the low velocity and the experimental uncertainty. Similarly the wavelength of the light emitted by the backward direction, i.e. as seen by a laboratory fixed observer looking at a receding atom is given by (2) [Ives H.E. and Stilwell, J.]

$$\lambda'_B = \lambda \frac{1 - \frac{v}{c} \cos \theta'}{\sqrt{1 - \beta^2}} = \lambda(1 - \beta \cos \theta' + \frac{1}{2}\beta^2) \quad (1) \quad \lambda'_R = \lambda \frac{1 + \frac{v}{c} \cos \theta'}{\sqrt{1 - \beta^2}} = \lambda(1 + \beta \cos \theta' + \frac{1}{2}\beta^2) \quad (2)$$

the subscript R indicated a *red Doppler shift*. The average of these two wavelengths

$$\lambda_Q = \frac{\lambda'_B + \lambda'_R}{2} = \lambda(1 + \frac{1}{2}\beta^2) \quad (28)$$

equals to the wavelength which would be observed at right angles to the beam corresponding  $\theta = \frac{\pi}{2}$  in previous equations. There are many reasons why a perpendicular observation of the beam is not feasible...

This kind of experiment measures in fact longitudinally (for  $\theta' \sim \theta \sim 0$ ) the factor  $1/2$  with respect to the second order term  $\beta^2$  (Mandelberg measures  $0,498 \pm 0.025$ ). Mandelberg defines a arithmetic average with the angle  $\theta'$ . With an arithmetic average based on the angle  $\theta$ , we obtain exactly the same result with Poincaré's formula  $(\frac{\lambda'}{\lambda})_{\text{Poincaré}} = \gamma(1 - \frac{v}{c} \cos \theta) \rightarrow \lambda_Q = \frac{\lambda'_B + \lambda'_R}{2} = \lambda(1 + \frac{1}{2}\beta^2)$ . Longitudinally Einstein's formula and Poincaré's formula are the same (25).

Given that SODS experiments directly measure (with the primed angle of the observer in the laboratory  $\theta'$ ) the length waves, we can write respectively "blueshift" and "redshift" Einstein's and Poincaré's formulas with  $\theta' \leq 90^\circ$  (22-24) in the following way:

$$(\frac{\lambda'_B}{\lambda})_{\text{Einstein}} = \gamma(1 - \frac{v}{c} \cos \theta') \quad (\frac{\lambda'_B}{\lambda})_{\text{Poincaré}} = \frac{1}{\gamma(1 + \frac{v}{c} \cos \theta')} \quad (29)$$

$$(\frac{\lambda'_R}{\lambda})_{\text{Einstein}} = \gamma(1 + \frac{v}{c} \cos \theta') \quad (\frac{\lambda'_R}{\lambda})_{\text{Poincaré}} = \frac{1}{\gamma(1 - \frac{v}{c} \cos \theta')} \quad (30)$$

Let us note that, without the relativistic factor  $\gamma$ , we find again both forms of classical formulas respectively of blueshift and redshift (the forms are inverted if we use unprimed proper angles  $\theta$ ). According to Mandelberg [Mandelberg H. I. & Witten L.], the measurement of transversal effect " ( $\theta' = \frac{\pi}{2}$ ) is not feasible<sup>6</sup>". Most of physicists have quoted the direct observation of the transversal effect to be extremely difficult or even impossible. This is the reason why most of experiments are on the Ives-Stilwell longitudinal configuration. At the first sight it seems impossible [Falk G, Ruppel W] to experimentally distinguish between Poincaré's and Einstein's formula. However, in 1979, Hasselkamp [Hasselkamp D. & all] introduces an experiment that directly measures the quadratic relativistic effect in the vicinity of the angle  $\theta' = 90^\circ$ . He rejects the argument according to which the Doppler-broadening of the line (which is a consequence of the finite opening angle of the optical system) could be much larger than the second order Doppler-shift). Hasselkamp specifies "the mechanical could only be performed to an accuracy in the real observation angle  $90^\circ < \theta' < 91^\circ$ . And also "The real angle of observation is  $90,5^\circ$ ". This is important for the discussion because Hasselkamp measures a redshift and if we use the blueshift formula (29 with  $90,5^\circ$ ) the quadratic effect is compatible with both formulas  $\cos \theta' = -\frac{v}{c}$ :

<sup>6</sup>Let us remark that the experimental point of view defined the transversality with observer angle whilst the theoretical point of view defined the transversality with the proper angle of the source. The question of  $\gamma$  the definition of transversality becomes very important with Poincaré's formula.

$(\frac{\lambda'}{\lambda})_{Poincaré} = \frac{1}{\gamma(1-\frac{v}{c}\cos\theta)}$  =  $\gamma$  if the magnitude of the velocities  $\frac{v}{c} \simeq 0.01$  but not compatible with Poincaré's formula for  $\frac{v}{c} \simeq 0.03$ . The function of velocity seems to confirm Einstein's formula. But the question is not completely clarified because according the figure 1 [Hasselkamp D. & all] , the domain of measure  $90^\circ < \theta' < 91^\circ$  is clearly a domain of *incoming* direction of atoms. So we must use the blueshift formulas (29,  $\theta' \leq 90^\circ$  ) between the angle  $90^\circ$  and  $89,5^\circ$  for which Einstein's formula gives respectively a redshift and a blueshift[Pierseaux Y.(2005)]. In order to experimentally make the difference between Einstein's formula and Poincaré's formula we have to measure within a symmetrical domain around the right angle  $89,5^\circ < \theta' < 91^\circ$ .

## References

- [Einstein A.(1905)] "Zur Elektrodynamik bewegter Körper", Ann. d. Ph., 17, p892-921, 1905, (we use English translation in "The Principle of relativity", introduction and comment by Sommerfeld, Dover, New York, p37-65, 1952).
- [Falk G, Ruppel W ] "Mechanik, RELativität, Gravitation, Berlin-Heidelberg-New York, Springer, 1973.
- [Moreau W.] "Wave front relativity", Am. J. Phys, **62** (5), May 1994.
- [Mandelberg H. I. & Witten L.] "Experimental verification of relativistic Doppler effect", Opt. Soc. Am. 52, number 5, 529, may 1962.
- [Hasselkamp D. & all] "Direct observation of the transversal Doppler-shift", Z. Physik, A 289, 151-155, 1979.
- [Ives H.E. and Stilwell, J. ] " An Experimental Study of the Rate of a Moving Atomic Clock", Opt. Soc. Am. 28, 215 (1938) 31, 369 (1941)
- [Pierseaux Y.(2005)] "Einstein's relativistic Doppler formula", soumis aux Annales de la fondation de Broglie.
- [Pierseaux Y (2004)] "Einstein's spherical light waves versus Poincaré's ellipsoidal light waves", London, pirt, September 2004, Annales de la Fondation de Broglie, vol 30, n°3 – 4, 2005 (<http://www.ensmp.fr/aflb/AFLB-303/table303.htm>).
- [Pierseaux Y. (2005-2)] "La cinématique relativiste avec photons d'Einstein et la cinématique relativiste avec ondes électromagnétiques de Poincaré", les Nouvelles d'Archimède, USTL, Octobre, Novembre, Décembre 2005.
- [Pierseaux Y & Rousseaux G.] "Les structures fines de la relativité restreinte et de l'électromagnétisme", article soumis en décembre 2005.
- [Pierseaux Y.(ellipse)] "La cinématique relativiste sous-jacente à l'ellipse de Poincaré", soumis aux Comptes-Rendus Physique en décembre 2005.
- [Poincaré H. 1905] "Sur la dynamique de l'électron", CR de l'Ac. des Sci. de Palerme, 27 juillet 1905. In Rendiconti d. Circ. mat. de Palermo, 21, 1906.
- [Poincaré H.(1908)] "La dynamique de l'électron". R. Gén. des Sciences Pures et Appl., 19, 386-402, 1908.
- [Poincaré H.(1912)] "La dynamique de l'électron", supplément aux annales des postes, télégraphes et téléphones (mars 1913), A. Dumas, Editeur, rue de la Chaussée d'Antin, 6, Paris.