

ION-AEROSOL INTERACTIONS IN THE LOWER ATMOSPHERE OF VENUS

K.L. Aplin

Space Science and Technology Department, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX UK

Introduction: the atmospheric electrical environment on Venus

All planetary atmospheres are electrified to some extent by cosmic ray ionization, and Venus is no exception. There is increasing awareness that ion-aerosol interactions could modulate terrestrial radiative processes, and this possibility will be investigated for the Venusian atmosphere. The likelihood of a Venusian global atmospheric electric circuit (defined below) will also be discussed.

Venus' atmosphere is dense enough for ions and electrons formed by cosmic rays to rapidly produce ion clusters¹. Common predicted positive species are $\text{H}_3\text{O}^+(\text{H}_2\text{O})_n$ ($n = 3$ or 4), $\text{H}_3\text{O}^+(\text{SO}_2)$ and $\text{H}_3\text{O}^+(\text{H}_2\text{O})(\text{SO}_2)$. Sulphate species dominate predicted negative ion species, and free electrons are also expected above 60km. Because of the ubiquity of the cloud cover (at ~50-70km), attachment to cloud particles is an important global ion loss mechanism, significantly reducing charged particle concentrations¹.

Could there be a global electric circuit on Venus?

A global atmospheric electric circuit needs, at the minimum, electrostatic discharge or charged particle precipitation balanced by current carried by mobile charged particles between a conductive ionosphere and surface². A global circuit causes transport of charged particles between different parts of the atmosphere. At the moment, the existence of a Venusian global electric circuit seems unlikely due to the lack of proof for electrostatic discharges³. If lightning is unambiguously observed, then there is a good basis for the existence of a global circuit because of the presence of ions and electrons and a conducting ionosphere.

Electrification and particle formation

A potentially important mechanism for cloud formation at the bottom of the cloud layer is vapour condensing onto aerosol particles⁴. The atmosphere is supersaturated with respect to sulphuric acid from 40km upwards, where H_2SO_4 vapour can condense onto hydrated sulphuric acid particles. As Venusian atmospheric ions are predicted to be sulphuric acid hydrates, appropriate conditions may exist for aerosols to form by ion-induced nucleation. The supersaturation required for ions to grow into ultrafine droplets by direct condensation can be determined using the Thompson equation⁵. This equation describes the equilibrium saturation ratio needed for ion-induced nucleation to become energetically favourable. The equilibrium condition is defined at a saturation ratio S , where r is radius, ρ fluid density, M the mass of the molecule, q charge, γ_T the surface tension, k_B Boltzmann's constant, T temperature, r_0 the initial radius (all in SI units), and ϵ_r relative permittivity:

$$\ln S = \frac{M}{k_B T \rho} \left[\frac{2\gamma_T}{r} - \frac{q^2}{32\pi^2 \epsilon_0 r^4} \left(1 - \frac{1}{\epsilon_r} \right) \right]. \quad (1)$$

(1) can be used to assess if condensation of gaseous H_2SO_4 onto ions is likely to occur in the lower cloud-forming regions at ~40km. To calculate S , H_2SO_4 concentrations and supersaturations determined from microwave absorption measurements are used⁶. Temperatures were obtained from a model atmosphere¹. Surface tension, density and dielectric constant for ~100% sulphuric acid at 400K were estimated and extrapolated from existing data².

(1) was then used to compute the saturation ratio needed for nucleation onto ions with 1, 2 and 5 electronic charges at a temperature T , for the latitudes at which the saturation ratios were measured.

At 88°S, supersaturation (SS) is expected at ~47km and 43km, but at the lower altitude the higher




Figure 1 Supersaturations required for H_2SO_4 nucleation onto ions with 1, 2 and 5 electronic charges for conditions at 88°S, 43 km.

temperatures reduce the SS needed for activation. Figure 1 shows the solution of (1) at 395K. SS is never high enough to activate particles with one electronic charge, but doubly charged particles of radius 1nm can nucleate at SS=7%, and particles with 5 charges of radius 2nm are activated at SS=1-2%, relatively easily attained SS levels⁶. The number of doubly charged particles can be estimated from the aerosol charge distribution arising from ion-aerosol attachment processes. The steady state charge distribution of a monodisperse aerosol population, represented as the ratio of the number of particles with charge j , N_j , to the number of neutral particles, N_0 , can be given by

$$\frac{N_j}{N_0} = x^j \frac{8\pi\epsilon_0 akT}{je^2} \sinh\left[\frac{je^2}{8\pi\epsilon_0 akT}\right] \exp\left[\frac{-j^2 e^2}{8\pi\epsilon_0 akT}\right] \quad (2)$$

where j is the electronic charge, a the mean aerosol radius, and x is the “ion asymmetry factor”⁷. The charge distribution can be calculated for T=395K and typical ion properties at 40km. The mobility of positive ions was assumed to be 7% higher than negative ions, with no free electrons below 60km¹. Positive and negative ions were assumed to exist in equal concentrations. The aerosol diameter at 40km was assumed to be 0.25µm¹. Substituting for a , T and x in (2) indicates that 84% of aerosols in this region carry some charge, with the most common single charge state being +1. The mean charge is +0.2, but, across the estimated charge distribution, Figure 2, $j \geq 2$ is relatively common with 54% of Venusian aerosols carry enough charge for ion-induced nucleation at supersaturations of 7%, and 7% of the aerosol population carries enough charge for nucleation at 1-2% supersaturation. These estimates suggest that it may be possible for gaseous sulphuric acid to condense onto ions and form aerosol particles.




Figure 2 Steady state charge distribution on Venus aerosols with mean diameter of 0.25 µm at 395 K.

Conclusions

- Venus almost certainly has active atmospheric electrical physics, though further measurements are needed to determine whether a global circuit exists. Because of cluster-ion formation in a dense gas, a global circuit could affect the location of charged particles and thus influence cloud or aerosol

particle formation and lifetime. From a comparative planetology viewpoint, a Venusian global circuit with its permanent cloud would be interesting to compare to the effect of variable cloud cover on Earth’s global circuit.

- Formation of aerosol particles by heterogeneous nucleation onto ions may be possible. If this does occur then cosmic rays modulated by the solar cycle could potentially affect climate.
- Any charged particle and aerosol measurements should be simultaneous so that ion-aerosol interactions can also be investigated. Larger charged particles should be measured, as well as the ions that contribute to air conductivity. This could be achieved with a relaxation probe, like the Huygens PWA probe, but using a higher voltage to capture larger particles.

References:

- ¹ Borucki W.J., et al *Icarus*, **51**, 302-321 (1982)
- ² Aplin K.L., *Surveys in Geophysics*, in press (2006)
- ³ Gurnett D.A. et al, *Nature* **409**, 313-315 (2001)
- ⁴ James E.P. et al, *Icarus*, **129**, 147-171, art. no. IS975763 (1997)
- ⁵ Mason B.J., *The physics of clouds*, Pergamon, Oxford (1971)
- ⁶ Kolodner M.A. and Steffes P.G., *Icarus* **132**, 151-169 (1998).
- ⁷ Clement C.F. and Harrison R.G., *J. Aerosol Sci.* **23**, 5, 481-504 (1992)