

# Influence of gravitational field on quantum-nondemolition measurement of atomic momentum in the dispersive Jaynes-Cummings model

M.Mohammadi<sup>1,2</sup> \*, M.H.Naderi<sup>3</sup> † and M.Soltanolkotabi<sup>3</sup> ‡

<sup>1</sup> Physics Department, Science and Research Campus Azad University Tehran, Iran.

<sup>2</sup> Physics Department, Shahreza and Islamic Azad University, Shahreza, Isfahan, Iran.

<sup>3</sup> Quantum Optics Group, University of Isfahan, Isfahan, Iran.

July 30, 2018

## Abstract

We present a theoretical scheme based on  $su(2)$  algebra to investigate the influence of homogeneous gravitational field on the quantum non-demolition measurement of atomic momentum in dispersive Jaynes-Cummings model. In the dispersive Jaynes-Cummings model, when detuning is large and the atomic motion is in a propagating light wave, we consider a two-level atom with quantized cavity-field in the presence of a homogeneous gravitational field. We derive an effective Hamiltonian describing the dispersive atom-field interaction in the presence of gravitational field. We can see gravitational influence both on the momentum filter and momentum distribution. Moreover, gravitational field decreases both tooth spacing of momentum and the width of teeth of momentum.

PACS numbers: 42.50.Ct, 42.50.Vk, 03.65.Ta, 03.75.Be

**Keyword:** Dispersive Jaynes-Cummings model, atomic motion, gravitational field, QND measurement

---

\*majid471702@yahoo.com

†mhnaderi2001@yahoo.com

‡soltan@sci.ui.ac.ir

# 1 Introduction

Among the models describing the interaction between light and matter, the Jaynes-Cummings model (JCM) [1] seems to be ideal. The JCM describes the interaction between a two-level atom and a single quantized mode of the electromagnetic field in a lossless cavity within the rotating wave approximation (RWA). This is the simplest model of the radiation-matter interaction. It is simple enough to be exactly solved on the one hand and complicated enough to exhibit many fascinating quantum features on the other hand. These pure quantum effects include quantum collapses and revivals of atomic inversion [2], squeezing of the radiation field [3], atomic dipole squeezing [4], vacuum Rabi oscillation [5] and dynamical entangling and disentangling of the atom field system in the course of time [6-8]. Further interest the JCM comes from the fact that its theoretical predictions have been extensively used in the context of quantum information [9], atoms and ions trapping [10,11] and quantum nondemolition (QND) measurements [12].

In a general QND measurement, an observable signal of a quantum system is measured by detecting a change in an observable of the probe system coupled to the quantum system during the measurement time, without perturbing the subsequent evolution of the observable signal. We can therefore make a sequence of precise measurements of an observable signal such that the result of each measurement is completely predictable from the result of the preceding measurement. Such an observable is called QND observable. Original QND ideas involved a dispersive coupling of the signal field to a material probe [13]. The QND method is quite generally based on dispersive and nonlinear effects. Methods to avoid the back action of the measurement on the detected observable have been proposed and implemented in the optical domain [14,15]. These experiments are the realization of the QND schemes introduced in [13]. They rely on nonlinear coupling of the signal field to be measured with a probe field whose phase is altered by a quantity depending on the number of photons in the signal beam. In a paper by Sleator and Wilkens [16] it was shown a complementary scheme in which a quadrature-component of a propagating laser wave acts as the probe for the QND measurement of the atomic momentum. It is based on the Doppler effect on the component of atomic momentum along the propagation direction of the light field.

On the other hand, experimentally, atomic beams with very low velocities are generated in laser cooling and atomic interferometry [17]. It is obvious that for atoms moving with a velocity of a few centimeters or meters per second for a time period of several milliseconds or more, the influence

of the Earth's acceleration becomes important and cannot be neglected [18]. For this reason it is of interest to study the temporal evolution of an atom moving in the gravitational field of the Earth and the cavity-field. Since any optical experiment in the laboratory is in a non-inertial frame it is important to estimate the influence of the Earth's acceleration on the outcome of these experiments. The effective Hamiltonian and time evolution for two-level atom that is simultaneously exposed to the field of a running laser wave and a homogeneous gravitational field is studied by Marzlin [19].

In this paper we investigate a complementary scheme based on  $\text{su}(2)$  algebra [20] to investigate the influence of gravitation on the QND measurement of atomic momentum in the dispersive JCM. In section 2 we apply scheme with  $\text{su}(2)$  algebra to describe the interaction of a two-level atom with quantized field and a homogeneous gravitational field in dispersive JCM. Recently, optical Schrödinger-cat states have been realized in dispersive JCM [21]. Also, these states have been verified experimentally, by Auffeves and coworkers, for a two-level atom interacting with a single mode of the electromagnetic field in dispersive JCM [22]. In the dispersive JCM, the atom is in ground state and detuning is large, so one can neglect spontaneous emission. In section 3 we investigate dynamical evolution of the system and show that how the gravitational field may be affected the dynamical properties of the dispersive JCM. In section 4 we investigate the influence of gravitational field on the QND measurement. Finally, we summarize our conclusions in section 5.

## 2 Dispersive Jaynes-Cummings Model in the presence of Gravitational Field

In the dispersive JCM, we assume that the atom is in its ground state initially and we consider the case of large detuning so that the excited state of the atom is almost never populated, so we neglect spontaneous emission. The importance of dispersive JCM is because of using in generation schrödinger cat states with superposition of coherent states. These states have been generated in different contexts. A great variety of methods have been proposed for generate such states, for example, in a Mach-Zehnder interferometer [23] and one of the first schemes due to Yurke and Stoler [24] who showed that a coherent state(CS) propagating in a Kerr medium could lead to a schrödinger cat state. We take into account the atomic motion along the position vector  $\hat{x}$ , so the evolution of the atom-field system in the

presence of gravitational field is governed by the Hamiltonian

$$\hat{H} = \frac{\hat{p}^2}{2M} - M\vec{g}\cdot\hat{\vec{x}} + \hbar\omega_c(\hat{a}^\dagger\hat{a} + \frac{1}{2}) + \frac{1}{2}\hbar\omega_{eg}\hat{\sigma}_z + \hbar\lambda[\exp(-i\vec{q}\cdot\hat{\vec{x}})\hat{a}^\dagger\hat{\sigma}_- + \exp(i\vec{q}\cdot\hat{\vec{x}})\hat{\sigma}_+\hat{a}], \quad (1)$$

where  $\hat{a}$  and  $\hat{a}^\dagger$  denote, respectively, the annihilation and creation operators of a single-mode traveling wave with frequency  $\omega_c$ ,  $\vec{q}$  is the wave vector of the running wave and  $\hat{\sigma}_\pm$  denote the raising and lowering operators of the two-level atom with electronic levels  $|e\rangle, |g\rangle$  and Bohr transition frequency  $\omega_{eg}$ . The atom-field coupling is given by  $\lambda$  and  $\hat{\vec{p}}, \hat{\vec{x}}$  denote, respectively, the momentum and position operators of the atomic center of mass motion and  $g$  is the Earth's gravitational acceleration. We find that an alternative representation of  $\text{su}(2)$  algebra arises naturally from the system. We construct a representation of  $\text{su}(2)$  algebra based on the generalized algebra and the pauli matrices. From Hamiltonian (1), it is apparent that there exist an operator  $\hat{K}$  with constant of motion

$$\hat{K} = \hat{a}^\dagger\hat{a} + |e\rangle\langle e|. \quad (2)$$

In addition, the operator  $\hat{a}\hat{\sigma}_+ = \hat{a}|e\rangle\langle g|$  commutes with  $\hat{K}$ . Now we introduce the following operators

$$\hat{S}_0 = \frac{1}{2}(|e\rangle\langle e| - |g\rangle\langle g|), \hat{S}_+ = \hat{a}|e\rangle\langle g|\frac{1}{\sqrt{\hat{K}}}, \hat{S}_- = \frac{1}{\sqrt{\hat{K}}}|g\rangle\langle e|\hat{a}^\dagger. \quad (3)$$

We can show that the operators  $\hat{S}_0, \hat{S}_\pm$  satisfy the following commutation relations

$$[\hat{S}_0, \hat{S}_\pm] = \pm\hat{S}_\pm, [\hat{S}_-, \hat{S}_+] = -2\hat{S}_0, \quad (4)$$

where  $S_0, S_\pm$  are the generators of  $\text{su}(2)$  algebra [25]. In terms of  $\text{su}(2)$  generators, the Hamiltonian(1) can be rewritten as

$$\hat{H} = \frac{\hat{p}^2}{2M} - M\vec{g}\cdot\hat{\vec{x}} + \hbar\omega_c\hat{K} + \frac{1}{2}\hbar\Delta\hat{S}_0 + \hbar\lambda\sqrt{\hat{K}}(\exp(-i\vec{q}\cdot\hat{\vec{x}})\hat{S}_- + \exp(i\vec{q}\cdot\hat{\vec{x}})\hat{S}_+), \quad (5)$$

where

$$\Delta = \omega_{eg} - \omega_c, \quad (6)$$

is the usual detuning parameter.

Now we start to find the exact solution for the dynamical evolution of the

total system governed by the Hamiltonian(5). The corresponding time evolution operator can be expressed as

$$\hat{u}(t) = \exp\left(\frac{iM\vec{g}\cdot\hat{x}t}{\hbar}\right)\hat{v}^\dagger\hat{u}_e(t)\hat{v}, \quad (7)$$

where

$$\hat{v} = \exp(-i\vec{q}\cdot\hat{x}\hat{S}_0), \quad (8)$$

$$\hat{u}_e = \exp\left(\frac{-i\hat{H}_e t}{\hbar}\right). \quad (9)$$

It can be shown that the operator  $\hat{u}_e(t)$  satisfy an effective Schrödinger equation governed by an effective Hamiltonian  $\hat{H}_e$ , that is

$$i\hbar\frac{\partial\hat{u}_e}{\partial t} = \hat{H}_e\hat{u}_e, \quad (10)$$

where

$$\hat{H}_e = \frac{\hat{p}^2}{2M} - \hbar\hat{\Delta}(\hat{p}, \vec{g})\hat{S}_0 + \frac{1}{2}Mg^2t^2 + \vec{g}\cdot\hat{p}t + \hbar\lambda(\sqrt{\hat{K}}\hat{S}_- + \sqrt{\hat{K}}\hat{S}_+) + \hat{H}_0, \quad (11)$$

$$\hat{H}_0 = \hbar\omega_c\hat{K} - \frac{\hbar}{2}\hat{\Delta}\hat{S}_0 - \frac{q^2\hbar^2}{2M}\hat{S}_0 + \frac{q^2\hbar^2}{8M}, \quad (12)$$

and the operator

$$\hat{\Delta}(\hat{p}, \vec{g}) = \omega_c - (\omega_{eg} + \frac{\vec{q}\cdot\hat{p}}{M} + \vec{q}\cdot\vec{g}t + \frac{q^2\hbar}{2M}), \quad (13)$$

has been introduced as the Doppler shift detuning at time t. Therefore, due to the Doppler shift of  $\frac{\vec{q}\cdot\hat{p}}{M}$ , recoil frequency  $\omega_{rec} = \frac{\hbar q^2}{2M}$  and gravitational field, the detuning between the cavity and the atomic transition frequency has been modified. The relevant time scale introduced by the gravitational influence is

$$\tau_a = \frac{1}{\sqrt{\vec{q}\cdot\vec{g}}}. \quad (14)$$

For an optical laser with ( $|\vec{q}| \simeq 10^7 m^{-1}$ ) and Earth's acceleration ( $|\vec{g}| = 9.8\frac{m}{s^2}$ ),  $\tau_a$  is about  $10^{-4}s$ . We remark that  $\hat{\Delta}(\hat{p}, \vec{g})$  does only depend on the product  $\vec{q}\cdot\vec{g}$ , so that the influence of the gravitational acceleration on the internal evolution vanishes if the acceleration is perpendicular to the laser field. Now we apply the interaction picture, i.e.,

$$\hat{u}_e = \exp\left(\frac{-it\hat{H}_0}{\hbar}\right)\hat{u}, \quad (15)$$

such that

$$i\hbar \frac{\partial \hat{u}}{\partial t} = \hat{H} \hat{u}, \quad (16)$$

where

$$\hat{H} = \frac{\hat{p}^2}{2M} - \hbar \hat{\Delta}(\hat{\vec{p}}, \hat{\vec{g}}) \hat{S}_0 + \frac{1}{2} M g^2 t^2 + \hat{\vec{p}} \cdot \hat{\vec{g}} t + \hbar (\kappa \sqrt{\hat{K}} \hat{S}_- + \kappa^* \sqrt{\hat{K}} \hat{S}_+), \quad (17)$$

and  $\kappa(t)$  is an effective coupling coefficient

$$\kappa = \lambda \exp\left(\frac{it}{2} (\hat{\Delta}(\hat{\vec{p}}, \hat{\vec{g}}) + \frac{\hbar g^2}{M})\right). \quad (18)$$

In the limit of very small values of  $|\langle \hat{\Delta}(\hat{\vec{p}}, \hat{\vec{g}}) \rangle|$  and  $\frac{\hbar g^2}{M}$ , the coefficient  $\kappa(t)$  is independent of time. As it stands, the effective Hamiltonian (17) has the form of the Hamiltonian of the JCM, the only modification being the dependence of the detuning on the conjugate momentum and the gravitational field. Now we use the JCM in the dispersive limit. In this limit, we assume that the atom is in its ground state initially and we consider the case of large detuning,  $|\delta\rangle\rangle \kappa \sqrt{\langle \hat{a}^\dagger \hat{a} \rangle}$ , with  $\delta \equiv \omega_c - \omega_{eg} - \omega_{rec}$ . In this case, the excited state of the atom is almost never populated, so we neglect spontaneous emission. Now we obtain effective Hamiltonian in dispersive regime. In the interaction picture the transformed Hamiltonian (17) takes the following form

$$\hat{H}_{int} = \exp\left(\frac{-i\hat{H}_0 t}{\hbar}\right) \hat{H}_I \exp\left(\frac{i\hat{H}_0 t}{\hbar}\right), \quad (19)$$

where

$$\hat{H}_0 = -\hbar \hat{\Delta}(\hat{\vec{p}}, \hat{\vec{g}}) \hat{S}_0, \quad (20)$$

$$\hat{H}_I = \hbar (\kappa \sqrt{\hat{K}} \hat{S}_- + \kappa^* \sqrt{\hat{K}} \hat{S}_+) + \hat{H}(\hat{\vec{p}}, \hat{\vec{g}}), \quad (21)$$

$$\hat{H}(\hat{\vec{p}}, \hat{\vec{g}}) = \frac{\hat{p}^2}{2M} + \hat{\vec{p}} \cdot \hat{\vec{g}} t + \frac{1}{2} M g^2 t^2. \quad (22)$$

Therefore we obtain

$$\hat{H}_{int} = \hbar (\kappa \sqrt{\hat{K}} \hat{S}_- \exp(-it \hat{\Delta}(\hat{\vec{p}}, \hat{\vec{g}})) + \kappa^* \sqrt{\hat{K}} \hat{S}_+ \exp(it \hat{\Delta}(\hat{\vec{p}}, \hat{\vec{g}}))) + \hat{H}(\hat{\vec{p}}, \hat{\vec{g}}). \quad (23)$$

We can obtain the effective Hamiltonian in dispersive limit [26]

$$\hat{H}_{eff} = \hat{H}(\hat{\vec{p}}, \hat{\vec{g}}) + \hbar \Omega(\hat{\vec{p}}, \hat{\vec{g}}) \hat{a}^\dagger \hat{a}, \quad (24)$$

where

$$\hat{\Omega}(\hat{\vec{p}}, \vec{g}) = \frac{|\kappa|^2}{\hat{\Delta}(\hat{\vec{p}}, \vec{g})}, \quad (25)$$

is the momentum-dependent frequency of the harmonic oscillator and identified as the Doppler modified ac stark shift of the atom-field interaction.

### 3 Dynamical Evolution

In section 2, we obtained the effective Hamiltonian atom-field system in the presence of the gravitational field in dispersive regime. In this section, we investigate dynamical evolution of system. We will show how the gravitational field may be affected the dispersive JCM. We will also investigate the dispersive JCM, in the short and long time limits. The Schrödinger equation reads as

$$i\hbar \frac{\partial |\psi\rangle}{\partial t} = \hat{H}_{eff} |\psi\rangle, \quad (26)$$

where

$$|\psi(t)\rangle = |\psi_g(t)\rangle \otimes |g\rangle. \quad (27)$$

In the dispersive regime, we define  $|\psi_g(t)\rangle$  as the state of the center of mass and the cavity field. We assume at  $t = 0$ , the atom-field system is described by the product state where the cavity field is initially prepared in a coherent state  $|\alpha\rangle$ . We apply evolution operator

$$\hat{u}(t) = \exp\left(\frac{-i}{\hbar} \int_0^t \hat{H}_{eff}(t') dt'\right), \quad (28)$$

on the initial state

$$|\psi_g(t=0)\rangle = \left(\int d^3p \phi_g(\vec{p}) |\vec{p}\rangle\right) \otimes |\alpha\rangle, \quad (29)$$

where  $\phi_g(\vec{p})$  is the probability amplitude for the center-of-mass motion of the ground-state atom in the momentum representation,  $|\vec{p}\rangle = |\hat{\vec{p}}|\vec{p}\rangle$ . When the atom leaves the cavity after an interaction time  $\tau$ , the state vector has evolved into the entangled state

$$|\psi_g(t=\tau)\rangle = \hat{u}(t=\tau) |\psi_g(t=0)\rangle, \quad (30)$$

$$\begin{aligned} |\psi_g(t=\tau)\rangle = & \int d^3p \exp\left(\frac{-i\tau p^2}{2M\hbar}\right) \exp\left(\frac{-i\vec{p}\cdot\vec{g}\tau^2}{2\hbar}\right) \exp\left(\frac{-iMg^2\tau^3}{6\hbar}\right) \phi_g(\vec{p}) |\vec{p}\rangle \\ & \otimes |\alpha \exp(-i\Omega(\vec{p}, \vec{g})\tau)\rangle. \end{aligned} \quad (31)$$

We now consider gravitational influence on the dynamical evolution of the system for two limiting case. The first, in the limit of small gravitational influence,  $t \ll \tau_a$ , means very small  $\vec{q} \cdot \vec{g}$ , i.e., the momentum transfer from the laser beam to the atom is only slightly altered by the gravitational acceleration because the latter is very small or nearly perpendicular to the laser beam. In this limit, the state vector is

$$|\psi_g(t = \tau)\rangle = \int d^3p \exp\left(\frac{-i\tau p^2}{2M\hbar}\right) \phi_g(\vec{p}) |\vec{p}\rangle \otimes |\alpha \exp(-i\Omega(\vec{p})\tau)\rangle, \quad (32)$$

where

$$\Omega(\vec{p}) = \frac{|\hat{\kappa}|^2}{\Delta(\vec{p})}, \Delta(\vec{p}) = \omega_c - \left(\omega_{eg} + \frac{\vec{q} \cdot \vec{p}}{M} + \frac{q^2 \hbar}{2M}\right). \quad (33)$$

The Doppler shift detuning is independent of gravitational field. The second, in the limit of long times,  $t \gg \tau_a$ , the atoms are accelerated by the Earth's gravity so that their velocity increases and the Doppler shift detuning in (13) is depends on the gravitational field.

## 4 The QND Measurement

The QND measurement has been the subject of numerous investigations in the past two decades [27-30]. For such a measurement on of a system, the system must be coupled to another system (called probe), and monitored an appropriately selected probe observable during the measurement. The system-probe interaction has to be chosen in such a way that the corresponding interaction Hamilton commutes with the system observable. The interaction of radiation with a single atom involves the electronic degrees of freedom of atom and center-of-mass degrees of freedom. The interaction of a two-level atom with a standing laser wave can result in a QND measurement of the atomic position when a quadrature component of the (sufficiently detuned) laser field is used as the probe [31,32]. In the previous section we show that how the gravitational field could be affected the dispersive Jaynes-Cummings model. In this section we introduce QND measurement and investigate the influence of gravitational field on the QND measurement of atomic motion. The Hermitian quadrature phase operator  $\hat{Y} = \frac{(\hat{a} + \hat{a}^\dagger)}{2}$  is used as a probe observable for a QND measurement of conjugate momentum  $\hat{p}$  when (17) satisfies condition  $\hat{H} = \hat{H}(\hat{p})$ ,  $[\hat{H}, \hat{p}] = 0$ ,  $[\hat{H}, \hat{Y}] \neq 0$ . The probability for obtaining value  $Y$  for the quadrature phase  $\hat{Y}$  may be expressed as

$$P(Y)dY = dY \int dp |\phi_g(\vec{p})|^2 |\langle Y | \alpha \exp(-i\Omega(\vec{p}, \vec{g})\tau) \rangle|^2. \quad (34)$$



The momentum distribution after a readout  $Y$  is given by the conditional probability  $P(\vec{p}|Y)$ , that the atom has a momentum vector  $\vec{p}$

$$P(\vec{p}|Y) = |\phi_g(\vec{p})|^2 |\langle Y | \alpha \exp(-i\Omega(\vec{p}, \vec{g})\tau) \rangle|^2 P(Y)^{-1}, \quad (35)$$

where

$$\begin{aligned} \langle Y | \alpha \exp(-i\Omega(\vec{p}, \vec{g})\tau) \rangle = & \left(\frac{2}{\pi}\right)^{\frac{1}{4}} \exp(-[|\alpha| \cos(\Omega(\vec{p}, \vec{g})\tau - \varphi_\alpha) - Y]^2) - \\ & 2i|\alpha|Y \sin(\Omega(\vec{p}, \vec{g})\tau - \varphi_\alpha). \end{aligned} \quad (36)$$

We define the momentum filter  $G(\vec{p}) = |\langle Y | \alpha \exp(-i\Omega(\vec{p}, \vec{g})\tau) \rangle|^2$ . The readout  $Y$  implies that the atom has a momentum that obeys  $\cos[\Omega(\vec{p}, \vec{g})\tau - \varphi_\alpha] \simeq \frac{Y}{|\alpha|}$ , with  $\alpha = |\alpha| \exp(i\varphi_\alpha)$ . Figures 1a and 2a show the form of the momentum filter  $|\langle Y | \alpha \exp(-i\Omega(\vec{p}, \vec{g})\tau) \rangle|^2$  and the momentum distribution  $P(\vec{p}|Y = 0)$  respectively assuming  $Y = 0$ . In these figures we assume  $(\frac{\kappa}{\delta})^2 \tau \omega_{rec} = 0.2$ ,  $\tau \omega_{rec} = 7.2$ ,  $\kappa \tau = 140$ ,  $\alpha = 2$ ,  $q = 10^7 m^{-1}$ ,  $M = 10^{-26} kg$ ,  $g = 9.8 \frac{m}{s^2}$ ,  $\tau = 14.4 \times 10^{-6} s$ , and  $\varphi_\alpha = \Omega(0)\tau + \frac{\pi}{2}$  [11,20]. Here we consider a beam of two-level atoms traversing an arm of an optical ring cavity, so that  $\vec{p} \cdot \vec{q} = pq \cos\theta$ ,  $\vec{q} \cdot \vec{g} = qg \sin\theta$ , with  $\theta = \frac{\pi}{4}$ . In figures 1a-1d and 2a-2d we plot the momentum filter and momentum distribution in terms of  $\frac{p}{\hbar q}$ . In these figures we show that gravitational field influence on the momentum filter and momentum distribution when the time increases. In figures 2a-2d one can see Oscillations. These oscillations result from quantum interference of translation motion [21]. To estimate the spacing for small  $\vec{p}$ , we expand  $\Omega(\vec{p}) \simeq \Omega(0) + (\frac{\kappa}{\delta - \vec{q} \cdot \vec{g} \tau})^2 \frac{\vec{q} \cdot \vec{p}}{M}$  and obtain

$$\Delta \vec{p} = |\vec{p}_{n+1} - \vec{p}_n| = \hbar q \frac{\pi}{2} \left( \frac{\delta - \vec{q} \cdot \vec{g} \tau}{\kappa} \right)^2 \frac{1}{\omega_{rec} \tau}. \quad (37)$$

The slow variation of  $\Delta \vec{p}$  in figures 1a-1d is due to the nonlinearity of  $\Omega(\vec{p}, \vec{g})$ , which leads to a decreasing tooth spacing for increasing momenta. In a simple Gaussian approximation, the width of the teeth near  $p = 0$  is given by

$$\sigma = \hbar q \frac{1}{4|\alpha|} \left( \frac{\delta - \vec{q} \cdot \vec{g} \tau}{\kappa} \right)^2 \frac{1}{\omega_{rec} \tau}. \quad (38)$$

From (37) and (38) one can see that the gravitational field decreasing both  $\Delta \vec{p}$  and  $\sigma$ .

## 5 Summary and conclusions

In this paper we have investigated the influence of gravitational field on the dynamical behavior of the JCM and on the QND measurement of atomic

momentum in dispersive JCM. For this purpose, based on  $su(2)$  algebra as the dynamical symmetry group of the model, we have derived an effective Hamiltonian describing the dispersive atom-field interaction in the presence of gravitational field. By finding an explicit form for the corresponding time evolution operator, we have explored the influence of gravitation on the atom-field coupling and detuning parameter. We have shown that due to the gravitational field the atomic transition frequency experiences a Doppler shift and atom-field coupling becomes time-dependent. Then we have investigated the influence of gravitational field on the QND measurement of atomic momentum in dispersive JCM. Moreover, we have shown that the gravitational field decreases both tooth spacing of momentum and the width of teeth of momentum.

### Acknowledgements

One of the authors (M.M) wishes to thank The Office of Graduate Studies of the Science and Research Campus Islamic Azad University of Tehran for their support.

### References

- [1] E.T.Jaynes and F.Cummings, Proc.IEEE **51**, 89 (1963).
- [2] H.I.Yoo, J.J.Sanchez-Mondragon and J.H.Eberly, J.Phys.A:Math.Gen **14**, 1383 (1981).
- [3] J.R.Kuklinski and J.Madajczyk, Phys.Rev.A **37**, 3175 (1988).
- [4] S.M.Barnett, Opt.Comm. **61** 432 (1982); P.Zhou and J.S.Peng, Phys.Rev.A **44** 3331 (1991)
- [5] G.S.Agarwal, J.Opt.Soc.Am.B **2** 480 (1985).
- [6] S.J.D.Phoenix and P.L.Knight, Ann.Phys.(N.Y.) **186** 381 (1988).
- [7] S.J.D.Phoenix and P.L.Knight, Phys.Rev.A **44** 6023 (1991).
- [8] A.Ekert and P.L.Knight, Am.J.Phys. **63** 415 (1995).
- [9] A.S.Soronsen and K.Molmer, Phys.Rev.Lett. **91**, 097905 (2003).
- [10] J.I.Cirac, R.Blatt, A.S.Parkins, and P.Zoller, Phys.Rev.Lett. **70**, 762 (1993).

- [11] D.M.Meekhof, C.Monroe, B.E.King, W.M.Itano, and D.J.Wineland, Phys.Rev.Lett. **76**, 1796 (1996).
- [12] W.J.Munro, Kao Nemoto, R.G.Beau Soleil, and T.P.Spiller, Phys.Rev.A **71**, 033819 (2005).
- [13] W.G.Unruh, Phys.Rev.D **18**, 1764 (1978).
- [14] G.J.Milburn and D.F.Walls, Phys.Rev.A **28**, 2065 (1980).
- [15] V.Braginsky and F.I.Khalili, Zh.Eksp.Teor.Fiz. **78**, 1712 (1980).
- [16] T.Sleator and M.Wilkens, Phys.Rev.A **48**, 3286 (1993).
- [17] C.Adamas, M.Sigel, and J.Mlynek, Phys.Rep. **240**, 143 (1994).
- [18] A.Kastberg, W.D.Philips, S.L.Rolston, R.J.C.Spreeuw, and P.S.Jessen, Phys.Rev.Lett. **74**, 1542 (1995).
- [19] S.Yu, H.Rauch, and Y.Zhang, Phys.Rev.A **54**, 2585 (1995).
- [20] R.L.de Matos Filho and W.Vogel, Phys.Rev.Lett. **76**, 608 (1996).
- [21] J.R.Kuklinski and J.Madajczyk, Phys.Rev.A **48**, 3291 (1993).
- [22] A.Auffeves, P.Maioli, T.Meunier, S.Gleyzes, G.Nogues, M.Brune, J.M.Raimond, and S.Haroche, Phys.Rev.Lett. **91**, 230405 (2003).
- [23] C.C.Gerry, Phys.Rev.A **59**, 4095 (1999).
- [24] B.Yurke and D.Stoler, Phys.Rev.Lett. **57**, 13 (1989).
- [25] K.P.Marzlin and J.Audertsch, Phys.Rev.A **53**, 1004 (1996).
- [26] W.P.Schleich, *Quantum optics in phase space*, (Springer,VCH,2001).
- [27] G.J.Milburn and D.F.Walls, Phys.Rev.A **28**, 2065 (1983).
- [28] N.Imoto, S.Watkins, and Y.Sasaki, Opt.commun. **61**, 159 (1987).
- [29] M.Brune, S.Haroche, V.Lefevre, J.M.Raimond, and N.Zagury, Phys.Rev.Lett. **65**, 976 (1990).
- [30] V.B.Braginsky and S.P.Vyatohanin, Phys.Lett.A **132**, 206 (1988).
- [31] P.Storey, M.Collett and D.Walls, Phys.Rev.Lett. **68**, 472 (1992).

[32] M.A.M.Marte and Zoller, Appl.Phys.B **54**, 477 (1992).

**FIGURE CAPTIONS:**

**FIG. 1** The momentum filter  $G(\vec{p}) = |\langle Y | \alpha \exp(-i\Omega(\vec{p}, \vec{g})\tau) \rangle|^2$  as a function of  $\frac{p}{\hbar q}$  that results a readout  $Y = 0$ . Here  $(\frac{\kappa}{\delta})^2 \tau \omega_{rec} = 0.2$ ,  $\tau \omega_{rec} = 7.2$ ,  $\kappa \tau = 140$ ,  $\alpha = 2$ ,  $q = 10^7 m^{-1}$ ,  $M = 10^{-26} kg$ ,  $g = 9.8 \frac{m}{s^2}$ .

a)  $\tau = 14.4 \times 10^{-6} sec$

b)  $\tau = 14.4 \times 10^{-5} sec$

c)  $\tau = 14.4 \times 10^{-4} sec$

d)  $\tau = 14.4 \times 10^{-3} sec$

**FIG. 2** Momentum distribution after a readout  $Y = 0$  has been detected. All parameters are as in fig. 1.

a)  $\tau = 14.4 \times 10^{-6} sec$

b)  $\tau = 14.4 \times 10^{-5} sec$

c)  $\tau = 14.4 \times 10^{-4} sec$

d)  $\tau = 14.4 \times 10^{-3} sec$