

We use the Fourier method to
strategies. Our results indicate
nities of banks, which can be c

*Corresponding author

To analyze the presence of comm we estimate the variance-covarian

The Italian electronic broker mar whole existing domestic overnight the Euro area in being a screen b interbank trades are largely bilate and foreign banks can exchange fu in 1999, 196 in 2000, 183 in 2001 a quotes to facilitate credit line che finalized if the ordering bank acc

In figure 2 we plot the distribution of highly active banks. We find that it follows a power-law distribution of the form $a \exp(-t/\tau)^\beta$ [1]. The parameter β is approximately 1.

We denote the signed trading volume of a single bank, where every transaction is counted with a plus sign if the bank lends some money and with a minus sign if the bank borrows a given liquidity, as $v_i(t)$. The total signed trading volume $V(t)$ is decreasing to zero thereafter.

implementation of a classical Pea

Bank code	a
1	3558
70	1031

Table 1: Parameter estimation for
for two highly active banks. The
form $a \exp(-t/\tau)^\beta$. $\langle dT \rangle$ is t
two banks.

the error and the largest the bias correlations among trading strategies with $M = 500$ are more centered around zero. There is not a great difference between the two cases. In the following results are obtained

3 Principal Components

We analyze the correlation matrix using Random Matrix Theory (RMT), in line with

a borrower). Thus, they transact among trading strategies is less st

4 Communities str

The aim of this section is to ide trading strategies. To help visual represent large Italian banks with Italian banks with green and sma

limits, while on a monthly level, in reserves at a lower frequency and other balance sheet positions over be able to play strategically at longer time scales. Nonetheless we matrices over the two time scales correlation network in figure 9 (to

To check that the method does not do not emerge purely from the f

¹For visualizations we used Graphviz

unless one knows, a priori, how many approaches based on the spectral analysis does not need any a priori knowledge (which is the most common case) of the optimization problem, in which we are interested by an harmonic potential energy.

Many spectral methods [31,32,33] use the (weighted) connectivity matrix A , and the functions of A adopted are the eigenvectors $N = K^{-1}A$, where K is the diagonal matrix of the number of nodes in the network.

Figure 9: Positive correlations graph
(top) and reshuffled transactions

Figure

figure 10.




Figure 13: (Left) Second eigenvalue reveals the presence of two common components for the reshuffled matrix.

for the correlation matrix obtained is not visible in this case.

As the second eigenvalue approach Figure 12 shows that near the EOM in contrast with our previous findings decreases when EOM approaches. correlation of strategies inside the relation between the two communities observed that very few banks cha

Finally we note that the first com

- [3] R. Reno, International Journal of Finance and Economics 87-102.
- [4] O. Precup, G. Iori, Physica A
- [5] M.O. Nielsen and P.H. Frederiksen, Journal of Economic Surveys 16(2002), 1-28
- [6] E. Barucci and R.Reno', Journal of Applied Money 12 (2002), 183-200
- [7] E. Barucci and R.Reno', Economic Letters 77(2002), 1-10
- [8] M. Mancino and R. Reno', Applied Financial Economics 13(2003), 1-10

- [27] J. Duch and A.Arenas, *Physica*
- [28] L. Danon, A.Diaz-Guilera, J. J. *Phys. Rev. E* 71 (2005) P09008 (2005).
- [29] M. Girvan, and M.E.J. Newman, *Phys. Rev. E* 65 (2002) of USA 99 (12) (2002), 7821.
- [30] A. Capocci., V.D.P. Servedio, *Phys. Rev. E* 71 (2005), 669-676.
- [31] K.M. Hall, *Management Science*

