Mechanical model of the Lorentz force and Coulomb interaction

Valery P. Dmitriyev Lomonosov University P.O.Box 160, Moscow 117574, Russia (Dated: 9 December 2007)

The centripetal and Coriolis accelerations experienced by a cart traveling over a rotating turntable are usually calculated proceeding from the known kinematics of the problem. Respective forces can be regarded as due to the entrainment of the cart in the moving solid environs. We extend the approach to the general case of a particle entrained in the flow of the surrounding medium. The expression for the driving force on the particle obtained from the kinematics of the entrainment prescribed appears to be isomorphic to the Lorentz and Coulomb force on a positive electric charge. The inverse direction of the electromagnetic force on a negative charge implies that a growing applied flow induces the upstream motion of the particle. A possible microscopic mechanism for it may be the Magnus force dynamics of a kink in a vortex tangle. The loop on a straight vortex filament can be taken as a model of the electron, the loop with a cavitation models the positron. The Lorentz force is concerned with the Coriolis acceleration. The Coulomb interaction is due to the centripetal or centrifugal force that arises in the turbophoresis of the kink in the perturbation field generated in the medium by the center of pressure.

1. INTRODUCTION

When considering a man walking in an accelerated train or a car traveling over the Earth's surface we introduce inertial forces. These fictitious forces can be calculated from kinematics of the system. Then, returning to the inertial reference frame, we may obtain real forces that the train and the Earth exert upon the bodies. In this context the moving plane and the rotating surface are actually flows of the solid medium that entrain the man and the car respectively. It is interesting to extend the approach to the general case of a particle carried by the stream of a medium. In this event we will not at first look into a physical origin of the entrainment but simply postulate that the particle initially at rest acquires the velocity of the entraining medium, i.e. it is advected as a tracer. We will assume also that the particle retains its own motion.

The special interest is the flow of an ideal fluid, or the Euler velocity field. A particle entrained in the Euler field will be shown to obey the equation that is isomorphic to the motion equation of the positive charge in the electric and magnetic fields.

The loop on a vortex filament possesses the property that it unfolds and begins to move upstream under the action of the accelerated flow. Choosing the kink on a vortex filament as the model of the electrically charged particle we may reproduce both signs of the electric charge. In this event the positive charge corresponds to the cavitated vortex.

2. CALCULATING REAL FORCES VIA THE KINEMATICS PRESCRIBED

The problem can be stated as follows. Given a velocity field $\mathbf{u}(\mathbf{r}, t)$. There is a particle at $\boldsymbol{\xi}$ that is affected by this field. We will assume that at any instant of time the full velocity \mathbf{v} of the particle can be represented as the sum of the particle's own velocity (relative velocity) \boldsymbol{v} and the surroundings' velocity:

$$\mathbf{v} = \mathbf{v} + \mathbf{u}(\boldsymbol{\xi}, t) \tag{1}$$

where the time dependence $\check{\boldsymbol{v}}(t)$ of the relative velocity in the reference frame $\check{\mathbf{r}}$ accompanying the velocity field at $\boldsymbol{\xi}$ is known. The relative motion of the particle will be assumed to be uniform and rectilinear, i.e.

$$\check{\boldsymbol{v}} = \frac{d\boldsymbol{\xi}}{dt},\tag{2}$$

$$\frac{d\tilde{\boldsymbol{v}}}{dt} = 0. \tag{3}$$

We want to find the particle's acceleration due to (1)-(3), or equivalently the force **f** on the particle (driving force) needed in order to ensure (1)-(3).

Formally \mathbf{f} can be found using (1) in the equation

$$m\frac{d\mathbf{v}}{dt} = \mathbf{f} \tag{4}$$

where m is the mass of the particle. The difficulty in the direct differentiating of (1) is that we do not know the dependence v(t) on time in the inertial reference frame underlying the medium. Still we know $\check{v}(t)$ as given by (3).

When v = 0, equation (1) can be taken as modeling the advection of a tracer particle.

Below the particular case of the kinematics will be firstly discussed that is known as the centripetal and Coriolis accelerations. Then the Lagrange technique will be used for a general treatment.

3. THE CORIOLIS FORCE

We will consider a particular yet the important case. A particle moving in two dimensions with relative velocity \boldsymbol{v} is entrained in a rotating field. We are interested in the evolution of the full velocity \mathbf{v} of the particle. Looking at the problem more generally the evolution of any vector entrained in a rotating field can be primarily discussed [1].

Let some vector **G** be rotated about an axis through an angle φ . As can be seen from geometrical considerations the angle between the vectors in initial and final positions will be φ as well. The increment of **G** due to rotation through a small angle $d\varphi$ is given by

$$(d\mathbf{G})_{\mathsf{rot}} = d\boldsymbol{\varphi} \times \mathbf{G} = dt\boldsymbol{\omega} \times \mathbf{G} \tag{5}$$

where $\boldsymbol{\omega}$ is the angle velocity. Substituting **r** for **G** in (5) we obtain the very rotating field

$$\mathbf{u} = \boldsymbol{\omega} \times \mathbf{r} \tag{6}$$

with the center of rotation at the origin of coordinates. Taking into account the rate of change of the vector in the rotating reference frame $\check{\mathbf{r}}$, (5) can be generalized to

$$\frac{d\mathbf{G}}{dt} = \frac{d\mathbf{\check{G}}}{dt} + \boldsymbol{\omega} \times \mathbf{G}.$$
(7)

Putting (6) in (1) we obtain the velocity of a particle entrained in the rotating field:

$$\mathbf{v} = \boldsymbol{v} + \boldsymbol{\omega} \times \boldsymbol{\xi}.\tag{8}$$

Differentiating (8) with respect to time we get

$$\frac{d\mathbf{v}}{dt} = \frac{d\boldsymbol{\upsilon}}{dt} + \boldsymbol{\omega} \times \frac{d\boldsymbol{\xi}}{dt} = \frac{d\boldsymbol{\upsilon}}{dt} + \boldsymbol{\omega} \times \mathbf{v}.$$
(9)

Using equation (7) for \boldsymbol{v} in the right-hand side of (9) and substituting (8) for \mathbf{v} gives

$$\frac{d\mathbf{v}}{dt} = \frac{d\tilde{\boldsymbol{v}}}{dt} + 2\boldsymbol{\omega} \times \boldsymbol{v} + \boldsymbol{\omega} \times (\boldsymbol{\omega} \times \boldsymbol{\xi}).$$
(10)

Using (3) and (4) in (10) and performing the vector multiplication we obtain the driving force of the entrainment

$$\mathbf{f} = 2m\boldsymbol{\omega} \times \boldsymbol{v} - m\omega^2 \boldsymbol{\xi}.$$
(11)

The normal inertial force $2m\boldsymbol{\upsilon}\times\boldsymbol{\omega}$ found from (11) is referred to as the Coriolis force.

4. DYNAMICS OF A PARTICLE ENTRAINED IN A VELOCITY FIELD

We will consider the general case of the dynamics of the entrained particle and then verify it by the rotating field. Converting the right-hand part of (1) into the reference frame accompanying the velocity field at $\boldsymbol{\xi}$ we get for the kinetic energy of the particle

$$K = \frac{1}{2}m[\check{\boldsymbol{v}}(t) + \check{\mathbf{u}}(\check{\boldsymbol{\xi}})]^2.$$
(12)

By the construction $\mathbf{\check{u}}(\boldsymbol{\check{\xi}}) \equiv 0$. However we retain it in (12) since may be $\mathbf{\check{u}}(\mathbf{\check{r}}) \neq 0$ for $\mathbf{\check{r}} \neq \boldsymbol{\check{\xi}}$. Lagrange equations for the evolution of the particle's relative velocity (2) are

$$\frac{d}{dt}\frac{\partial K}{\partial \check{\boldsymbol{v}}} = \frac{\partial K}{\partial \check{\boldsymbol{\xi}}} + \check{\mathbf{f}}.$$
(13)

Substituting (12) in (13) we have with the account of (2)

$$\frac{d}{dt}\frac{\partial K}{\partial \check{\boldsymbol{v}}} = m\frac{d}{dt}\left(\check{\boldsymbol{v}} + \check{\mathbf{u}}\right) = m\left[\frac{d\check{\boldsymbol{v}}}{dt} + \left(\check{\boldsymbol{v}}\cdot\check{\boldsymbol{\nabla}}\right)\check{\mathbf{u}}\right],\tag{14}$$

$$\frac{\partial K}{\partial \boldsymbol{\check{\xi}}} = \frac{1}{2} m \boldsymbol{\check{\nabla}} \left(\boldsymbol{\check{\upsilon}} + \boldsymbol{\check{u}} \right)^2 = m \left(\boldsymbol{\check{\upsilon}} \cdot \boldsymbol{\check{\nabla}} \right) \boldsymbol{\check{u}} + m \boldsymbol{\check{\upsilon}} \times \left(\boldsymbol{\check{\nabla}} \times \boldsymbol{\check{u}} \right) + \frac{1}{2} m \boldsymbol{\check{\nabla}} \boldsymbol{\check{u}}^2$$
(15)

where the well-known vector relation $\nabla (\mathbf{a} \cdot \mathbf{b}) = (\mathbf{a} \cdot \nabla) \mathbf{b} + (\mathbf{b} \cdot \nabla) \mathbf{a} + \mathbf{a} \times (\nabla \times \mathbf{b}) + \mathbf{b} \times (\nabla \times \mathbf{a})$ was used. Using (14) and (15) in (13) we get

$$m\frac{d\check{\boldsymbol{\upsilon}}}{dt} = m\check{\boldsymbol{\upsilon}} \times (\check{\boldsymbol{\nabla}} \times \check{\mathbf{u}}) + \frac{1}{2}m\check{\boldsymbol{\nabla}}\check{\mathbf{u}}^2 + \check{\mathbf{f}}.$$
(16)

Using (3) in (16) gives for the driving force of the entrainment

$$\check{\mathbf{f}} = -\frac{1}{2}m\check{\boldsymbol{\nabla}}\check{\mathbf{u}}^2 - m\check{\boldsymbol{\upsilon}} \times (\check{\boldsymbol{\nabla}} \times \check{\mathbf{u}}).$$
⁽¹⁷⁾

Next we may convert (17) into the inertial reference frame taking advantage of that gradient and curl are coordinateindependent and $\check{\boldsymbol{v}} = \boldsymbol{v}$:

$$\mathbf{f} = -\frac{1}{2}m\nabla\mathbf{u}^2 - m\mathbf{v} \times \text{curl}\mathbf{u}.$$
 (18)

In order to interpret terms in the right-hand part of (18) we will apply (18) to the particular case, the rotating field. We have for (6) in two dimensions

$$\operatorname{curl} \mathbf{u} = 2\boldsymbol{\omega},$$
 (19)

$$\frac{1}{2}\boldsymbol{\nabla}\mathbf{u}^2 = \omega^2 \boldsymbol{\xi}.$$
(20)

Substituting (19)-(20) into (18) we come to the relation (11) previously obtained from kinematical considerations.

5. PARTICLE DRIVEN BY THE EULER FIELD

Equation (18), (4) can be viewed as describing dynamics of a particle entrained in the stream of a fluid. By (18) particles will drift to the region where the velocity is low. This theoretical prediction is in accord with observations that the litter floating in the river is carried away to the bank. The flow of an ideal fluid is governed by the Euler equation. The latter can be written down in the Gromeka-Lamb form

$$\frac{\partial \mathbf{u}}{\partial t} + \frac{1}{2} \nabla \mathbf{u}^2 - \mathbf{u} \times \operatorname{curl} \mathbf{u} + \frac{1}{\varrho} \nabla p = 0$$
(21)

where ρ is the density of the fluid. Multiplying (21) by m and using it in (18) gives with the account of (1)

$$\mathbf{f} = m\frac{\partial \mathbf{u}}{\partial t} + \frac{m}{\varrho}\boldsymbol{\nabla}p - m\mathbf{v} \times \text{curl}\mathbf{u}.$$
(22)

6. THE ELECTROMAGNETIC ANALOGY

Expression (22) has the same structure as the force on the positron moving in the electromagnetic field. Supposing that the field \mathbf{A} entrains the positron we will postulate the aether flow with the velocity \mathbf{u} , and define

$$\mathbf{A} = \kappa c \langle \mathbf{u} \rangle \tag{23}$$

where

$$\kappa = -\frac{m_{\rm e}}{e} \tag{24}$$

 m_{e} is the mass, e the charge of the positron, and $\langle .. \rangle$ an averaging. Further we may define [2] the mechanical analogy for the scalar potential

$$\varrho\varphi = \kappa \langle p - p_0 \rangle \tag{25}$$

where p_0 is the background pressure of the aether. Taking $m = m_e$ and substituting (23)-(25) into (22) properly averaged gives the electromagnetic force on the positron

$$\mathbf{f}_{+} = -\frac{e}{c}\frac{\partial \mathbf{A}}{\partial t} - e\boldsymbol{\nabla}\varphi + \frac{e}{c}\mathbf{v} \times \text{curl}\mathbf{A}.$$
(26)

In order to model the dynamics of the electron we must find an entity that may move upstream. It is a vortex formation in the fluid that possesses such the property.

7. VORTEX RINGS IN THE APPLIED FLUID FLOW

In the external fluid flow vortices are undergone to the lateral thrust referred to as the Magnus force [3]. Thus, the force on a linear vortex is given by $\rho \mathbf{u} \times \boldsymbol{\kappa}$ where κ is the fluid circulation about the vortex. Due to the Magnus force we have the specific behavior of an ideal vortex ring in the flow of the fluid. Normally the vortex ring is traveling in the fluid with a constant velocity \boldsymbol{v} of self-induction. Under the action of the Magnus force it turns and becomes oriented along the flow (see Fig.1). In this event the ordinary vortex ring (non-dilatational vortex ring) may move upstream [4] (see Fig.1 right), and the cavitated vortex ring [5] moves downstream.



FIG. 1: The Magnus force $\mathbf{f}_{\mathsf{M}} \sim \mathbf{u} \times \boldsymbol{\kappa}$ on a vortex ring in the ambient velocity field \mathbf{u} , where $\boldsymbol{\kappa}$ is the vorticity; \boldsymbol{v} velocity of the non-dilatational ring in a quiescent fluid (the cavitated ring travels in the opposite direction). A possible initial position (*left*), and the final configuration (*right*). Circles show the circulation of the fluid in the cross section of the ring.

In terms of wave motion the upstream-oriented vortex ring is concerned with a pressure wave, and the downstreamoriented vortex ring corresponds to a rarefaction wave.

8. THE LOOP ON A VORTEX FILAMENT

A quantum particle can be modeled [6] by the loop on a straight vortex filament [7]. The loop can be seen as a vortex ring attached to the filament after the appropriate reconnection accomplished. Being flat the loop is translationally at rest, and rotates steadily about the filament (see Fig.2 left). When in a non-planar configuration the kink travels along the filament with a velocity \boldsymbol{v} . By the analogy with the vortex ring, a non-dilatational or rather compressional kink moves in the direction of the right-hand screw (see Fig.2 middle and right). If associated with a cavity the same configuration of the kink travels in the opposite direction. Similarly to the vortex ring, the kink on the vortex filament is carried away by the flow so that the fluid velocity \mathbf{u} adds to its own velocity \boldsymbol{v} as is written in (1).

The loop unfolds into the non-planar configuration under the action of an external force. In particular, the external force may be caused by the accelerated fluid flow. As we see from the direction of the Magnus force, in the growing flow the compressional loop begins to move upstream (see Fig.2 middle and right). The cavitated loop moves downstream.

We may assign to the kink a translational momentum and the respective effective mass $m_e > 0$ that has the meaning of the kink's self-energy, i.e. the fluid kinetic energy concerned with the loop on the vortex filament [8].



FIG. 2: The loop on a vortex filament with the direction of vorticity shown by the arrow. The flat configuration, side view (*left*). The growing velocity of the flow is **u** and of the kink v: side view (*middle*), top view(*right*).

9. THE NEGATIVE MASS

Under the action of an external force the ordinary kink (non-dilatational kink) moves in the direction opposite to the direction of the force. So, we may assign to the non-dilatational kink, or rather a compressional kink the negative mass $-m_{e}$. Alternatively we may change the sign of the force in the dynamic law (4) to the opposite. The cavitated kink has the positive effective mass. We will try the compressional vortex loop as the model of the electron, and cavitated vortex loop as the model of the positron. Insofar as the pressure wave may be symmetric to the dilatational wave masses of the solitons considered should be identical. Changing the sign of (22) gives the force on the compressional kink

$$\mathbf{f} = -m_{\mathbf{e}} \frac{\partial \mathbf{u}}{\partial t} - \frac{m_{\mathbf{e}}}{\varrho} \nabla p + m_{\mathbf{e}} \mathbf{v} \times \text{curl} \mathbf{u}.$$
(27)

Substituting (23)-(25) into (27) we obtain the force on the electron

$$\mathbf{f}_{-} = \frac{e}{c} \frac{\partial \mathbf{A}}{\partial t} + e \nabla \varphi - \frac{e}{c} \mathbf{v} \times \text{curl} \mathbf{A}.$$
(28)

Generalizing (26) and (28) to the case of the charge q of any sign and value we come to the well-known expression for the electromagnetic force on a charged particle

$$\mathbf{f} = -\frac{q}{c}\frac{\partial \mathbf{A}}{\partial t} - q\boldsymbol{\nabla}\varphi + \frac{q}{c}\mathbf{v} \times \text{curl}\mathbf{A}.$$
(29)

10. THE VORTEX SPONGE

In three dimensions the kink model of the positron or electron should be seen as a reconnecting vortex ring in the structure of the vortex sponge (cf. the reconnection vortex-tangle model [9]). The vortex sponge was postulated [10] as an ideal fluid pierced in all directions by straight vortex filaments. We may see the vortex tangle as the vortex sponge on a large scale. In this event the tangle may be constituted of a single filament. The latter may be the first cause of that all electrons are identical.

11. THE LORENTZ FORCE

Let us discuss the hydrodynamics of the electromagnetic interaction as it is given by equations (18), (21) and (22). The crucial point is the origin of the Lorentz force. We have $\nabla p = 0$ and $\partial \mathbf{u}/\partial t = 0$ in the solenoid, and the field $\mathbf{u}(\mathbf{r})$ the same as on the turntable. The question is how it may be the Coriolis driving force $-m\mathbf{v} \times \text{curl}\mathbf{u}$ from (18) without the centripetal force $-m\nabla \mathbf{u}^2/2$? The answer is that the latter is somehow compensated by $m\mathbf{u} \times \text{curl}\mathbf{u}$ from (21). In the result we obtain the term $-m\mathbf{v} \times \text{curl}\mathbf{u}$ in (22) for the Lorentz force.

So, the Lorentz force is due to the Coriolis driving force that bends the velocity of the particle in the rotational flow of the fluid. Because of the peculiar behavior of the vortex loop the compressional kink will be rotated in the opposite direction.

12. THE COULOMB INTERACTION

The electric interaction is concerned with the centripetal force term $-m\nabla \mathbf{u}^2/2$ in (18). The question is what may be the flow that provides the term 1/r of the Coulomb field for \mathbf{u}^2 ? The answer is that it is the system of the

vortex sponge, or a perturbation of the turbulence, in a macroscopic description. Following Osborn Reynolds the fluid velocity can be expanded into the sum of the average value $\langle \mathbf{u} \rangle$ and fluctuation \mathbf{u}' :

$$\mathbf{u} = \langle \mathbf{u} \rangle + \mathbf{u}' \tag{30}$$

where $\langle .. \rangle$ is the symbol of the short-time averaging, so that

$$\langle \mathbf{u}' \rangle = 0. \tag{31}$$

The essential point in (30) is that the turbulent fluctuation is much greater than the average fluid flow:

$$\langle \mathbf{u}^{\prime 2} \rangle \gg \langle \mathbf{u} \rangle^2$$

$$\tag{32}$$

where $\langle \mathbf{u}'^2 \rangle = \langle u'_i u'_i \rangle$ and $\langle \mathbf{u} \rangle^2 = \langle u_i \rangle \langle u_i \rangle$, and the summation over the recurrent index is implied throughout. Substituting (30) into $\langle \mathbf{u}^2 \rangle$, using (31) in it and by (32) dropping the quadratic term $\langle \mathbf{u} \rangle^2$ we obtain for the averaged first term from (18)

$$\mathbf{f}_{\mathsf{C}} = -\frac{1}{2} m_{\mathsf{e}} \boldsymbol{\nabla} \langle \mathbf{u}^{\prime 2} \rangle. \tag{33}$$

Formula (33) models the force exerted upon the charge in the electrostatic field. The exact expression of the electric force term from (22) via Reynolds stresses $\langle u'_i u'_k \rangle$ looks as follows [2]

$$\frac{\partial \langle \mathbf{u} \rangle}{\partial t} + \frac{1}{\varrho} \nabla \langle p \rangle = \partial_k \langle \mathbf{u}' u_k' \rangle.$$
(34)

By (33) in the non-uniform turbulence the particle will drift towards the regions of decreasing turbulence intensity. Because of the peculiar behavior of the vortex loop the compressional kink will move in the opposite direction.

So, the Coulomb interaction should be attributed [2] to the turbophoresis [11] of the vortex kink. The field 1/r of the perturbation of the turbulence energy density $\rho \langle u'_i u'_i \rangle / 2$ is generated by a center of pressure. The positive $p > p_0$ pressure center models the electron. The negative $p < p_0$ pressure center corresponds to the positron or proton [2].

13. THE MASS PARTICLE

We used the kinematical approach in order to calculate real forces exerted on the vortices in the laminar and turbulent fluid flows. Next the forces thus found can be applied to more complex formations.

The neutron was earlier modeled [2] by the vapor bubble in the turbulent aether. The mass of the neutron is measured by the mass of the vapor in the bubble. The bubble associated with a vortex loop can be taken as the model of the proton. We will assume that the bubble by itself is not advected by the aether flow. Hence all the drive on the proton will be concerned with the vortex attached. Then the equation of motion of the proton can be obtained uniting (4) and (18), and replacing m by m_e in the right-hand part:

$$m\frac{d\mathbf{v}}{dt} = -\frac{1}{2}m_{\mathbf{e}}\boldsymbol{\nabla}\mathbf{u}^{2} - m_{\mathbf{e}}\boldsymbol{\upsilon} \times \operatorname{curl}\mathbf{u}$$
(35)

where m is the mass of the proton. Multiplying (21) by m_e , using it in (35), averaging and substituting (23)-(25) in the result we obtain the proper electromagnetic equation. This rounds off outlines of the mechanical model of the magnetic and electric interaction.

- [9] K.W.Shwarz, 'Three-dimensional vortex dynamics in superfluid H_4 : homogeneous superfluid turbulence', *Phys.Rev.* B 38, 2398-2417 (1988).
- [10] E.M.Kelly, 'Vacuum electromagnetics derived exclusively from the properties of an ideal fluid', Nuov.Cim. 32 B, No 1, 117-137 (1976).
- [11] M.W.Reeks, 'The transport of discrete particles in inhomogeneous turbulence', J. Aerosol Science **310**, 729-739 (1983).

^[1] H.Goldstein, *Classical mechanics*, Cambridge Mass, Addison-Wesley Press, 1st edition.

 ^[2] V.P.Dmitriyev, 'Mechanical realization of the medium imitating electromagnetic fields and particles', *Meccanica* 42, No 3, 283-291 (2007).

^[3] E.R.Huggins, 'Exact Magnus-force formula for three-dimensional fluid-core vortices', Phys. Rev. A 1, No 2, 327-337 (1970).

^[4] A.H.M.Eisenga, R.Verzicco, and G.J.F.van Heijst, 'Dynamics of a vortex ring moving perpendicularly to the axis of a rotating fluid', J.Fluid Mech. 354, 215-239 (1998).

^[5] K.Shariff, A.Leonard, 'Vortex rings', Annu. Rev. Fluid Mech. 24, 235-279 (1992).

^[6] V.P.Dmitriyev, 'Mechanical analogy for the wave-particle: helix on a vortex filament', J. Appl. Math. 2, No 5, 241-263 (2002).

^[7] V.P.Dmitriyev, 'Helical waves on a vortex filament', Am. J. Phys. **73**, No 6, 563-565 (2005).

^[8] V.P.Dmitriyev, 'Particles and charges in the vortex sponge', Z.Naturforsch. 48a, No 8/9, 935-942 (1993).