

# Statistical Modelling of the Relationship Between Main Development Region Sea Surface Temperature and *Landfalling* Atlantic Basin Hurricane Numbers

Roman Binter (RMS and LSE)  
 Stephen Jewson (RMS) \*  
 Shree Khare (RMS)

May 29, 2018

## Abstract

We are building a hurricane number prediction scheme that relies, in part, on statistical modelling of the empirical relationship between Atlantic sea surface temperatures and landfalling hurricane numbers. We test out a number of simple statistical models for that relationship, using data from 1900 to 2005 and data from 1950 to 2005, and for both all hurricane numbers and intense hurricane numbers. The results are very different from the corresponding analysis for basin hurricane numbers.

## 1 Introduction

We are interested in developing practical methods for the prediction of the distribution of the number of hurricanes that might make landfall in the US over the next 5 years. One possible way to make such predictions is via a 2-step method that involves predicting main development region (MDR) sea surface temperature (SST), and then predicting landfalling hurricane numbers as a function of the MDR SST. The first step of predicting MDR SST has been considered in Meagher and Jewson (2006) and Laepple et al. (2006). This paper investigates the second step, and considers statistical relations that one might use to model the relationship between MDR SST and the number of landfalls. This paper closely follows an earlier paper (Binter et al., 2006) in which we modelled the relationship between MDR SST and the number of hurricanes in the basin. The data and the models we use, and the format of the results we present, are all taken from that paper. Readers should refer to that paper for further details including a short discussion, with references, giving an overview of the physical relationship between SST and hurricanes.

The rest of this article proceeds as follows: in section 2 we show the results from our tests on the landfalling hurricane data, and in section 3 we discuss what we have found.

## 2 Results

We now present results from our comparisons of the ability of various statistical models to represent the relationship between MDR SST and landfalling hurricane numbers. First we consider models for the total number of landfalling hurricanes, for the periods 1900-2005 and 1950-2005, and then we consider models for intense landfalling hurricane numbers for the same two periods.

### 2.1 All hurricanes, 1900-2005

The first results we present are based on all landfalling hurricanes, and data from 1900 to 2005. The scatter plot shown in figure 1 shows the number of hurricanes versus the SST during this period. The picture is dramatically different from what we saw when we considered the relationship between SST and the number of hurricanes in the basin in Binter et al. (2006). In that case there was a clear positive relationship. In this case, there is, at least *prima facie*, no relationship at all. Table 1 shows that the linear correlation is 0.16 and the rank correlation is 0.12.

---

\* Correspondence email: [stephen.jewson@rms.com](mailto:stephen.jewson@rms.com)

Table 2 shows the score comparisons for the six models for this data set, and table 4 shows the p-values for the pairwise comparison of these models. The best model, in terms of out-of-sample RMSE performance, is the exponential negative binomial model, but the score is only slightly better than the trivial flat poisson model which has no relationship between landfalling numbers and SST. The point-wise comparisons show that the non-trivial models are not statistically distinguishable from the trivial model. As far as the log-likelihood scores in table 2 are concerned, we find that the flat poisson model defeats the linear and damped linear normal models in a statistically significant way. The remaining models do not beat the flat-line model in a statistically significant way.

In table 3 we see that the slope parameters of all the non-trivial models are not significantly different from zero. In other words, based on these parameter estimates and standard errors, we certainly couldn't reject a null-hypothesis that there is no relation at all between MDR SST and the number of landfalling hurricanes.

The damping parameter in the damped linear trend model is much less than 1 (0.73), in response to the weak (or perhaps non-existent) signal that we are trying to identify.

The linear relationships between SST and hurricane numbers, for what they are worth, give a sensitivity of between 0.64 and 0.69 hurricanes per degree.

In summary: we don't see any indication of a relationship between MDR and SST and landfalling hurricane numbers.

## 2.2 All hurricanes, 1950-2005

Given the lack of a significant relationship between SST and hurricane numbers on the data from 1900 to 2005, it is interesting to see if we can find one using the more recent data. On the one hand, the more recent data is more accurate, and so it might be more likely we can detect a relationship. On the other hand, using less data will make it even harder to estimate the parameters of the models.

Table 6 shows that all the non-trivial models defeat the flat poisson model. The results for linear normal, damped linear normal and linear poisson are statistically significant.

As far as the log-likelihood scores are concerned, the linear normal and damped linear normal models are defeated by the flat poisson model in a statistically significant way. The linear poisson model defeats the trivial model in a statistically significant way (for RMSE). However, based on these results, it's hard to conclude definitively that the linear poisson model is best. For instance, in the pairwise comparisons for RMSE and log-likelihood, the exponential poisson model also defeats the linear poisson model in a statistically significant way.

Once again the slope parameters in the non-trivial models are all indistinguishable from zero. The linear relationships now have slopes between 0.99 and 1.2, but with standard errors of up to 0.63.

Overall, there is at least now *some* statistical evidence of a relationship between SST and landfalling hurricane numbers, although the parameters of the relationship are very poorly estimated.

## 2.3 Intense hurricanes

We have only just found a relationship between MDR SSTs and the total number of landfalling hurricanes. Could there be a relationship between MDR SSTs and the number of *intense* landfalling hurricanes, for which there is even less data? The scatter plots (figures 7 and 8) suggest that there is no such relationship. Our statistical results (tables 10 to 17) confirm that none of the non-trivial models defeat the flat trivial model in a statistically significant way for both data sets 1900-2005 and 1950-2005. In fact, for the 1950-2005 data set, the flat-line model defeats the linear and damped linear normal model in a statistically significant way in the probabilistic comparison.

## 3 Summary

We have investigated whether there is a statistical relationship between MDR SST and the number of hurricanes making landfall in the US. In previous work we've seen a strong relationship between MDR SST and the number of Atlantic basin hurricanes, and so our *a priori* assumption is that there must be some relationship for landfall numbers as well. Our analysis, however, finds only a weak relationship for total hurricane numbers and no relationship at all for intense hurricane numbers.

Why could this be? We see two possible explanations:

- There *is* a physical relationship between SST and landfalling hurricane numbers, but this relationship is mostly obscured by the signal-to-noise ratio, which is very poor because there are so few landfalling hurricanes.

- Even though there is a strong physical relationship between SST and basin hurricane numbers, there is only a weak physical relationship between SST and landfalling hurricane numbers. The effects of SST conspire to change the proportion of hurricanes that make landfall in such a way that the effects that are seen in the basin numbers almost disappear when we consider the landfalls. For instance, higher SSTs may mean higher numbers of hurricanes in the basin (on average), but they may also mean a lower proportion making landfall, and these two effects may combine in such a way that the actual number making landfall remains the same.

And of course reality may be a combination of these two effects.

We are investigating both of these possibilities. W.r.t. the first, we are considering simple statistical systems to see whether this is really the behaviour that we would expect under idealized assumptions. W.r.t. the second, we are performing statistical tests to investigate the hypothesis that the proportion of hurricanes making landfall really does vary in such a way as to mask the effects of SST on numbers of hurricanes.

## References

- R Binter, S Jewson, and S Khare. Statistical modelling of the relationship between Main Development Region Sea Surface Temperature and Atlantic Basin hurricane numbers. *arXiv:physics/0701170*, 2006. RMS Internal Report E04a.
- T Laepple, S Jewson, J Meagher, A O'Shay, and J Penzer. Five-year ahead prediction of Sea Surface Temperature in the Tropical Atlantic: a comparison of simple statistical methods. *arXiv:physics/0701162*, 2006.
- J Meagher and S Jewson. Year ahead prediction of hurricane season SST in the tropical Atlantic. *arXiv:physics/0606185*, 2006.

Table 1: Linear and Rank Correlations

	Linear Correlation	Rank Correlation
1900 - 2005 Landfall vs SST	0.16	0.12
1950 - 2005 Landfall vs SST	0.25	0.16
1900 - 1949 Landfall vs SST	0.17	0.25
1900 - 2005 Intense Landfall vs SST	0.28	0.24
1950 - 2005 Intense Landfall vs SST	0.25	0.09

Table 2: RMSE comparison 1900 - 2005 Landfall vs SST

	model name	RMSE (in)	RMSE (out)	100-100*RMSE/RMSEconst	LL (in)	LL (out)
model 1	Flat Poisson	1.399	1.412	0	-1.655	-1.666
model 2	Linear Normal	1.381	1.411	0.165	-1.841	-1.864
model 3	Damped Linear Normal	1.382	1.415	-0.359	-1.842	-1.866
model 4	Linear Poisson	1.381	1.41	0.372	-1.642	-1.664
model 5	Exponential Poisson	1.379	1.411	0.264	-1.641	-1.664
model 6	Exponential Neg. Bin.	1.379	1.387	3.577	-1.638	-1.649

Table 3: Model parameters incl. out of sample RMSE 1900 - 2005 Landfall vs SST

	$\hat{\alpha}$	s.e.	$\hat{\beta}$	s.e.	$k$	cov	corr	RMSE (out of sample)
model 1	0.541	0.074						1.412
model 2	1.717	0.135	0.686	0.415		0	0	1.411
model 3	1.717		0.502		0.732			1.415
model 4	1.717	0.127	0.645	0.39		0.006	0.123	1.41
model 5	0.532	0.075	0.395	0.225		-0.002	-0.128	1.411
model 6	0.532	0.078	0.391	0.236		-0.002	-0.115	1.387

Table 4: Winning count for particular model 1900 - 2005 Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	44 (0.897)	44 (0.897)	44 (0.897)	44 (0.897)	43 (0.928)
model 2	56 (0.143)	0 (1)	53 (0.314)	52 (0.385)	57 (0.103)	45 (0.857)
model 3	56 (0.143)	47 (0.752)	0 (1)	47 (0.752)	46 (0.809)	49 (0.615)
model 4	56 (0.143)	48 (0.686)	53 (0.314)	0 (1)	50 (0.539)	45 (0.857)
model 5	56 (0.143)	43 (0.928)	54 (0.248)	50 (0.539)	0 (1)	49 (0.615)
model 6	57 (0.103)	55 (0.191)	51 (0.461)	55 (0.191)	51 (0.461)	0 (1)

Table 5: Winning count (LL) for particular model 1900 - 2005 Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	76 (0)	75 (0)	44 (0.897)	44 (0.897)	43 (0.928)
model 2	24 (1)	0 (1)	53 (0.314)	25 (1)	25 (1)	22 (1)
model 3	25 (1)	47 (0.752)	0 (1)	22 (1)	23 (1)	22 (1)
model 4	56 (0.143)	75 (0)	78 (0)	0 (1)	50 (0.539)	47 (0.752)
model 5	56 (0.143)	75 (0)	77 (0)	50 (0.539)	0 (1)	49 (0.615)
model 6	57 (0.103)	78 (0)	78 (0)	53 (0.314)	51 (0.461)	0 (1)




Figure 1: 1900 - 2005 Landfall vs. SST




Figure 2: 1900 - 2005 Landfall vs. SSTs




Figure 3: Fitted Lines for all Models 1900 - 2005 Landfall vs SST

Table 6: RMSE comparison 1950 - 2005 Landfall vs SST constant

	model name	RMSE (in)	RMSE (out)	100-100*RMSE/RMSEconst	LL (in)	LL (out)
model 1	Flat Poisson	1.424	1.45	0	-1.637	-1.66
model 2	Linear Normal	1.379	1.441	1.24	-1.852	-1.91
model 3	Damped Linear Normal	1.381	1.449	0.057	-1.854	-1.914
model 4	Linear Poisson	1.38	1.435	2.023	-1.604	-1.65
model 5	Exponential Poisson	1.366	1.431	2.528	-1.598	-1.649
model 6	Exponential Neg. Bin.	1.367	1.361	11.935	-1.592	-1.615

Table 7: Model parameters incl. out of sample RMSE 1950 - 2005 Landfall vs SST constant

	$\hat{\alpha}$	s.e.	$\hat{\beta}$	s.e.	$k$	cov	corr	RMSE (out of sample)
model 1	0.463	0.106						1.45
model 2	1.589	0.188	1.18	0.622		0	0	1.441
model 3	1.589		0.923		0.783			1.449
model 4	1.589	0.168	0.991	0.562		0.018	0.187	1.435
model 5	0.439	0.108	0.72	0.342		-0.008	-0.213	1.431
model 6	0.44	0.116	0.696	0.37		-0.008	-0.179	1.361

Table 8: Winning count for particular model 1950 - 2005 Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	38 (0.978)	38 (0.978)	38 (0.978)	39 (0.959)	41 (0.93)
model 2	62 (0.041)	0 (1)	57 (0.175)	57 (0.175)	46 (0.748)	45 (0.825)
model 3	62 (0.041)	43 (0.886)	0 (1)	39 (0.959)	43 (0.886)	50 (0.553)
model 4	62 (0.041)	43 (0.886)	61 (0.07)	0 (1)	38 (0.978)	50 (0.553)
model 5	61 (0.07)	54 (0.344)	57 (0.175)	62 (0.041)	0 (1)	48 (0.656)
model 6	59 (0.114)	55 (0.252)	50 (0.553)	50 (0.553)	52 (0.447)	0 (1)

Table 9: Winning count (LL) for particular model 1950 - 2005 Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	79 (0)	79 (0)	38 (0.978)	39 (0.959)	41 (0.93)
model 2	21 (1)	0 (1)	57 (0.175)	23 (1)	23 (1)	21 (1)
model 3	21 (1)	43 (0.886)	0 (1)	23 (1)	23 (1)	23 (1)
model 4	62 (0.041)	77 (0)	77 (0)	0 (1)	38 (0.978)	48 (0.656)
model 5	61 (0.07)	77 (0)	77 (0)	62 (0.041)	0 (1)	50 (0.553)
model 6	59 (0.114)	79 (0)	77 (0)	52 (0.447)	50 (0.553)	0 (1)






Figure 4: 1950 - 2005 Landfall vs. SST




Figure 5: 1950 - 2005 Landfall vs. SST




Figure 6: Fitted Lines for all Models 1950 - 2005 Landfall vs SST

Table 10: RMSE comparison 1900 - 2005 Intense Landfall vs SST

	model name	RMSE (in)	RMSE (out)	100-100*RMSE/RMSEconst	LL (in)	LL (out)
model 1	Flat Poisson	0.836	0.844	0	-1.074	-1.084
model 2	Linear Normal	0.801	0.823	4.932	-1.24	-1.351
model 3	Damped Linear Normal	0.802	0.825	4.538	-1.24	-1.356
model 4	Linear Poisson	0.803	0.816	6.624	-1.019	-1.03
model 5	Exponential Poisson	0.799	0.827	3.997	-1.031	-1.054
model 6	Exponential Neg. Bin.	0.799	0.807	8.605	-1.031	-1.04

Table 11: Model parameters incl. out of sample RMSE 1950 - 2005 Intense Landfall vs SST

	$\hat{\alpha}$	s.e.	$\hat{\beta}$	s.e.	$k$	cov	corr	RMSE (out of sample)
model 1	-0.429	0.12						0.844
model 2	0.651	0.079	0.73	0.241		0	0	0.823
model 3	0.651		0.659		0.902			0.825
model 4	0.651	0.078	0.912	0.11		0.009	1	0.816
model 5	-0.494	0.128	1.094	0.361		-0.016	-0.338	0.827
model 6	-0.494	0.128	1.094	0.361		-0.015	-0.338	0.807

Table 12: Winning count for particular model 1900 - 2005 Intense Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	43 (0.928)	43 (0.928)	43 (0.928)	44 (0.897)	44 (0.897)
model 2	57 (0.103)	0 (1)	55 (0.191)	45 (0.857)	67 (0)	57 (0.103)
model 3	57 (0.103)	45 (0.857)	0 (1)	45 (0.857)	59 (0.032)	58 (0.072)
model 4	57 (0.103)	55 (0.191)	55 (0.191)	0 (1)	63 (0.004)	59 (0.032)
model 5	56 (0.143)	33 (1)	41 (0.98)	37 (0.998)	0 (1)	53 (0.314)
model 6	56 (0.143)	43 (0.928)	42 (0.951)	41 (0.98)	47 (0.752)	0 (1)

Table 13: Winning count (LL) for particular model 1900 - 2005 Intense Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	50 (0.539)	54 (0.248)	43 (0.928)	44 (0.897)	44 (0.897)
model 2	50 (0.539)	0 (1)	55 (0.191)	37 (0.998)	37 (0.998)	37 (0.998)
model 3	46 (0.809)	45 (0.857)	0 (1)	37 (0.998)	37 (0.998)	37 (0.998)
model 4	57 (0.103)	63 (0.004)	63 (0.004)	0 (1)	63 (0.004)	59 (0.032)
model 5	56 (0.143)	63 (0.004)	63 (0.004)	37 (0.998)	0 (1)	54 (0.248)
model 6	56 (0.143)	63 (0.004)	63 (0.004)	41 (0.98)	46 (0.809)	0 (1)




Figure 7: 1900 - 2005 Intense Landfall vs SST




Figure 8: 1900 - 2005 Intense Landfall vs SST




Figure 9: Fitted Lines for all Models 1900 - 2005 Intense Landfall vs SST

Table 14: RMSE comparison 1950 - 2005 Intense Landfall vs SST

	model name	RMSE (in)	RMSE (out)	100-100*RMSE/RMSEconst	LL (in)	LL (out)
model 1	Flat Poisson	0.875	0.891	0	-1.085	-1.107
model 2	Linear Normal	0.848	0.903	-2.692	-1.267	-1.514
model 3	Damped Linear Normal	0.85	0.906	-3.454	-1.268	-1.528
model 4	Linear Poisson	0.848	0.896	-1.211	-1.055	-1.112
model 5	Exponential Poisson	0.837	0.903	-2.873	-1.05	-1.117
model 6	Exponential Neg. Bin.	0.837	0.825	14.307	-1.049	-1.067

Table 15: Model parameters incl. out of sample RMSE 1950 - 2005 Intense Landfall vs SST

	$\hat{\alpha}$	s.e.	$\hat{\beta}$	s.e.	$k$	cov	corr	RMSE (out of sample)
model 1	-0.442	0.167						0.891
model 2	0.643	0.115	0.71	0.383		0	0	0.903
model 3	0.643		0.551		0.775			0.906
model 4	0.643	0.107	0.635	0.35		0.011	0.302	0.896
model 5	-0.495	0.175	1.06	0.532		-0.029	-0.306	0.903
model 6	-0.493	0.181	1.03	0.557		-0.027	-0.278	0.825

Table 16: Winning count for particular model 1950 - 2005 Intense Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	50 (0.553)	50 (0.553)	50 (0.553)	54 (0.344)	54 (0.344)
model 2	50 (0.553)	0 (1)	50 (0.553)	50 (0.553)	54 (0.344)	48 (0.656)
model 3	50 (0.553)	50 (0.553)	0 (1)	50 (0.553)	46 (0.748)	52 (0.447)
model 4	50 (0.553)	50 (0.553)	50 (0.553)	0 (1)	59 (0.114)	46 (0.748)
model 5	46 (0.748)	46 (0.748)	54 (0.344)	41 (0.93)	0 (1)	50 (0.553)
model 6	46 (0.748)	52 (0.447)	48 (0.656)	54 (0.344)	50 (0.553)	0 (1)

Table 17: Winning count (LL) for particular model 1950 - 2005 Intense Landfall vs SST

	model 1	model 2	model 3	model 4	model 5	model 6
model 1	0 (1)	64 (0.022)	62 (0.041)	50 (0.553)	54 (0.344)	54 (0.344)
model 2	36 (0.989)	0 (1)	50 (0.553)	36 (0.989)	36 (0.989)	39 (0.959)
model 3	38 (0.978)	50 (0.553)	0 (1)	36 (0.989)	36 (0.989)	41 (0.93)
model 4	50 (0.553)	64 (0.022)	64 (0.022)	0 (1)	59 (0.114)	46 (0.748)
model 5	46 (0.748)	64 (0.022)	64 (0.022)	41 (0.93)	0 (1)	50 (0.553)
model 6	46 (0.748)	61 (0.07)	59 (0.114)	54 (0.344)	50 (0.553)	0 (1)




Figure 10: 1950 - 2005 Intense Landfall vs. SST




Figure 11: 1950 - 2005 Intense Landfall vs. SST




Figure 12: Fitted Lines for all Models 1950 - 2005 Intense Landfall vs SST



