A note on the light velocity anisotropy

Bruno Preziosi

Dipartimento di Scienze Fisiche - Università Federico II di Napoli

ABSTRACT

In the framework of linear transformations between inertial systems, there are no physical reasons for assuming any anisotropy in the one-way velocity of light.

Pacs - 03.30 Special relativity Key words: inertial systems, Lorentz transformations, speed of light

1 Introduction

The anisotropy in the microwave background⁽¹⁾ has suggested the existence of a preferred frame Σ which sees an isotropic background and of a correspondent anisotropy in the one-way velocity of light, when measured in our system *S*, which moves with respect to Σ at the velocity of about 377 Km/s. This possibility has been exploited from the theoretical point of view^(2,3); many important and precise experiments have then been carried out with the purpose of detecting this anisotropy. No variation was observed at the level of 2×10^{-13} ⁽⁴⁾, 3×10^{-9} ⁽⁵⁾ and 2×10^{-15} ⁽⁶⁾.

We discuss here the generalized Lorentz transformations which have stimulated these experiments.

2 Local inertial systems

The starting point of our discussion is the fact that the background radiation intensity appears to be anisotropic to an observer O, at the origin of a reference frame S in our "region" of the universe R. It is quite reasonable to suppose, as we do, that the universe mass distribution too is anisotropic from the point of view of O, while it is seen isotropic by the observer Ω at the origin of a preferred reference frame Σ . We suppose moreover that all the reference frames we are speaking about are settled in accordance with the following procedure:

a) the origins are attached to bodies, which are "far enough" from other celestial bodies and are well oriented with respect to identical very far objects;

b) identical conventional space and time standards are used in all the reference systems.

The important point is that the absolute system Σ and the relative frame S of our region of the universe appreciate differences in the radiation background, but cannot "locally" appreciate any force due to the different mass distribution. The "region" R behaves like the world inside the Einstein elevator; the Einstein equivalence principle ensures that, if Ω and O perform identical experiments in their respective "regions", which are not influenced by the presence of local masses (Earth, Sun, ...), they obtain identical results. An immediate consequence is that the inertia principle is valid for all the local inertial systems.

This concept is very clearly stated by Hans Reichenbach⁽⁷⁾:

It can be shown within the framework of Newton's theory that one can obtain local inertial systems by transforming away the gravitational field, although these systems are in a different state of motion provided that the equivalence of inertial and gravitational mass is presupposed. The gravitational field, which as such is still present, is compensated in these local systems by their acceleration relative to absolute space and the resulting inertial forces. According to Einstein, however, only these local systems are the actual inertial systems. In them the field, which generally consists of a gravitational and an inertial component, is transformed in such manner that the gravitational component disappears and only the inertial component remains. The astronomical inertial systems of Newton can at best be approximations which gradually change in the neighborhood of stars. Only because distances in space are large compared to the masses of the stars, and because the stars have very low speeds, are astronomical inertial systems possible as approximations.

The words "far enough" used in a) express this approximation.

3 Linear transformations

If the transformations between these inertial systems are taken to be linear, then the condition that any inertial motion is such for any local inertial system is implicity satisfied. Many authors have added, to this hypothesis of linearity, other hypotheses with the purpose of assuring consistency between the measures of different observers; many of them^(8–23) have not postulated, a priori, the invariance of the light speed. They conclude that these transformations must be of Lorentz-type, characterized by a velocity **c** which, in principle, may take different absolute value in the different astronomical directions, but which is, for any fixed astronomical direction, equal for all the local inertial systems.

The consequences of the existence of a preferred reference frame Σ have been extensively discussed by Mansouri and Sexl⁽²⁾; in particular they analyze the form the linear transformation from Σ to an other frame S must take in order that the classical special relativity experiments could be explained, without introducing *a priori* the invariance of the speed of light. Moreover they compare the Einstein and the transport syncronization procedure and derive the dependence of the one-way velocity of light on its motion direction.

One of the most general transformations between the inertial systems Σ and S, which relatively move along the *x*-direction which connects the origins Ω and O, has been given by Robertson⁽⁴⁾ in the form:

$$t' = a_0 \left(t + \frac{va_1}{a_0 c^2} x \right), \ x' = a_1 \left(x + \frac{va_0}{a_1} t \right), \ y' = a_2 y, \ z' = a_2 z,$$
(1)

where a_0 , a_1 and a_2 may depend on v. This transformation, which is expressed in terms of the parameter v and which reduces to the identity when v = 0, is derived under the hypotheses that: i) space is euclidean for both Σ and S which use identical rods and clocks; ii) in Σ all clocks are syncronized and light moves with a speed c which is independent of direction and position; iii) the one-way speed of light in S in the azymuth plane is direction independent.

Notice that the relative velocity of S with respect to Σ is not given by v, but is equal to $\tilde{v} \equiv va_0/a_1$; analogously, the one-way velocity of light is given by $\tilde{c} \equiv ca_0/a_1$. In terms of these true velocities, equations (1) take the form

$$t' = a_0 \left(t + \frac{\tilde{v}}{\tilde{c}^2} x \right), \ x' = a_1 \left(x + \tilde{v} t \right), \ y' = a_2 y, \ z' = a_2 z.$$
(2)

The inverse transformation in the x-direction is given by

$$x = \frac{1}{a_1(\tilde{v})(1 - \frac{\tilde{v}^2}{\tilde{c}^2})} \left(x' - \tilde{v} \frac{a_1(\tilde{v})}{a_0(\tilde{v})} t' \right).$$
(3)

Moreover, the velocity \tilde{c} is seen by S to take the value

$$\tilde{c}' = \frac{a_1(v)}{a_0(v)}\tilde{c}.$$
(4)

In other words, the one-way velocity of light scales as the relative velocity of the two frames. As a consequence, if S and S' are any two reference frames which are moving (along the x-axis) with respect to Σ , they appreciate different relative velocities and different same-way velocities of light which are in the same ratio. It follows that, if the unit lenght of S is such a way that O and Ω appreciate the same relative speed, then $a_0 = a_1$ and viceversa.

It is interesting to notice that, if the Robertson hypotheses are released and one looks for the general one-dimensional linear transformations along the x-axis between the preferred frame Σ and any S which satisfies the condition that

-not only Σ and S appreciate the same relative speed but also any two frames S and S', uniformly moving at finite velocities with respect to Σ , appreciate the same finite speed,

-then one fundamental velocity (which obviously must be identified with the light speed) exists, the syncronization between two points of a frame is in agreement with the Einstein rule and the transformations take the form of the first two of eqs. (2) with $a_0 = a_1$ (see references (6)-(20) and, in particular, (21)).

The transformation (2) is simply the product of the scale transformation which scales the four dimensions respectively by the factors

$$a_0(\tilde{v})\sqrt{1-\frac{\tilde{v}^2}{\tilde{c}^2}}, \ a_1(\tilde{v})\sqrt{1-\frac{\tilde{v}^2}{\tilde{c}^2}}, \ a_2(\tilde{v}), \ a_2(\tilde{v})$$
 (5)

by a standard Lorentz transformation.

The scale transformations, first considered by Galilei in *Discorsi sovra* due nuove Scienze, were carefully analyzed by J. Fourier, H. von Helmoltz and in particular by H. Poincaré, who reached the conclusion that a pure spatial deformation of the universe has no physical consequences; moreover, if the space and time coordinates of the universe are deformed in such manner that all the space-time coincidences are conserved, then the universe remains unchanged.⁽²⁴⁾

If we consider the particular case of eqs. (2) in which $\tilde{v} = 0$, we have

$$ds^{2} = dx'^{2} + dy'^{2} + dz'^{2} = a_{1}^{2}dx^{2} + a_{2}^{2}dy^{2} + a_{2}^{2}dz^{2}.$$
 (6)

This case is typical of a tetragonal crystal; the position of the second nearest neighborhood is described in S by the numbers (1, 1, 0) and in S' by the components $(a_1, a_2, 0)$. The distances remain the same. It is worthwhile to notice that, in this case, if we scale the units of S' in such a way that they become equal, the times required by the light to travel the reticular distances in the different directions remain unchanged, but the light velocities are corresponding scaled; the anisotropy of the tetragonal crystal remains untouched.

4 Conclusions

The general transformation (2) reduces to the Lorentz one if one requires that Ω and O evaluate that the clock of the other observer is going slower in the same ratio (or, equivalently, if each of them appreciate an identical contraction for the reciprocal rods).

The effect of the scale transformations (5) must be carefully analyzed. For what concerns the lenghts in different directions, if there is no way for independent measures of lenghts and light velocities, in other words, if unit lenghts are measured by fixing the light velocity (or viceversa), there is no way of controlling the change in lenght of a rod with the orientation. The only thing to do is to use the Poincaré simplicity criterion and consider equal the lenghts of the rods and the one-way speeds of light in the different directions.

In conclusion, isotropy in the one-way speed of light is a matter of definition and the experiments conducted by J. Hall and coworkers at very sofisticated levels must be considered significant improvements of some classical experiments in the frame of the special relativity.

There are however experiments which cannot be explained in the frame of generalized linear Lorentz transformation; this is the case when the time required by the light to cover a poligonal is not in the ratio with the time required to cover the apotheme, which is provided by the Euclidean geometry. Notice that the explanation is not found in supposing that the light speed along the perimeter is different from the one along the apotheme (on the contrary, from the point of view of a local observer the velocity of a photon along its trajectory is always $c^{(25)}$), but in requiring that the space structure is no more Euclidean.

Analogously, the experiments by Shapiro *et al.*⁽²⁶⁾ and Reasenberg *et al.*⁽²⁷⁾ on the radar echo delay show that an e.m. wave travelling close to

the Sun appears not only deviated, but also slightly retarded to an observer in a far flat region. This phenomenon too cannot be explained by supposing a delay for the light which comes from the Sun to the Earth, but must be settled in the general relativity frame.

5 Acknowledgments

Thanks are due to professors John L. Hall and Giuseppe Marmo for useful discussions.

6 References

 G.F.Smoot, M.V. Gorenstein and R.A. Muller, *Phys. Rev. Lett.* **39**, 898 (1977)

R.M.Mansouri and R.U.Sexl, J. Gen. Rel. Grav. 8, 497 (1977); 8,515 (1977); 8,809 (1977)

3. H.P.Robertson, *Rev. Mod. Phys.* **21**, 378 (1949); H.P.Robertson and T.W.Noonan, *Relativity and Cosmology*, Saunders, Philadelphia, (1968) p. 69

4. A. Brillet and J.L. Hall, *Phys. Rev. Lett.*, **42**, 549 (1979)

5. E. Riis, L. Aaen Andersen, N. Bjerre, O.Poulsa, S.A.Lee, J. L. Hall, *Phys. Rev. Lett.* **60**, 81, (1988)

6. Dieter Hils and J.L. Hall, Phys. Rev. Lett. 64, 1697 (1990)

7. Hans Reichenbach, *Philosophie der Raum-Zeit-Lehre* (Walter de Gruyter, Berlin and Leipzig, 1928) translated by M. Reichenbach and J. Freund as *The Philosophy of space and time* (Dover New York, N. Y. 1958)

8. W. von Ignatowsky, Arch. Math. Phys. (Leipzig) 17, 1 (1910)

9. P. Frank and H. Rothe Ann. Phys. Leipzig 34, 825 (1911)

10. F. Severi, *Rend. Regia Acc. Lincei* 1924 and *Cinquant'anni di relativita'*, Sansoni ed., Firenze, 1955

11. C. Cattaneo, Rend. Acc. Lincei, 24, 256, (1958)

12. R. Weinstock, Am. J. Phys. 33, 640, (1965)

13. V. Mitvalsky, Am. J. Phys. 34, 825 (1966)

14. L. J. Eisenberg, Am. J. Phys. 35, 649 (1967)

15. V. Berzi and V. Gorini *J. Math. Phys.* 10, 1518 (1969) 16. A.R.Lee and T.M.Kalotas, *Am. J. Phys.* **43**, 5, 1975

17. Jean-Marc Lévy-Leblond, Am. J. Phys. 44, 271 (1976) and Riv. N.C.

7, 187 (1977); Jean-Marc Lévy-Leblond and Jean Pierre Provost, Am. J.

Phys. 47, 12, (1979)

- 18. A.M. Srivastava, Am. J. Phys. 49, 504 (1981)
- 19. N. D. Mermin, Am. J. Phys. 52, 119 (1984)
- 20. H. M. Schwartz, Am. J. Phys. 53, 1007 (1985)
- 21. S. Singh, Am. J. Phys. 54, 183 (1986)
- 22. A. W. Ross, Am. J. Phys. 55, 174 (1987)
- 23. B.Preziosi, N. Cim. 109 B, 1331, (1994)
- 24. Moritz Schlick, *Raum und Zeit in der gegenwärtigen Physik*, (Springer Verlag, Berlin, 1922)
- 25. I.R. Kenyon, General relativity, Oxford University Press, 1990, p. 44
- 26. I.I. Shapiro *et al.*, *Phys. Rev. Lett.* **26**, 1132 (1971) 27. R.D. Reasenberg *et al. Astr. J.* **234**, 1219 (1979)