Landau damping: is it real?

V. N. Soshnikov *

Plasma Physics Dept., All-Russian Institute of Scientific and Technical Information of the Russian Academy of Sciences (VINITI, Usievitcha 20, 125219 Moscow, Russia)

To calculate linear oscillations and waves in dynamics of gas and plasma one uses as a rule the old classical method of dispersion equation. The latter connects frequencies ω and wave numbers k (both are in general case complex values): $\epsilon(\omega, k) = 0$. In the plasma case ϵ is called generalized dielectric susceptibility. Dispersion equation is derived by substitution of asymptotical solution exp $(-i\omega t + i\vec{k}\vec{r})$ into the coupled system of linearized dynamical and field partial differential equations. However, this method appears to be inapplicable, f.e., in the case of waves in Maxwellian collisionless plasma when dispersion equation has no solutions. By means of some refined sophistication L. Landau in 1946¹ has suggested in this case actually to replace the dispersion equation with another one, having a specific solution ("Landau damping") and being now widely used in plasma physics. Recently we have suggested² a quite new universal method of two-dimensional Laplace transformation (in coordinate x and time t for plane wave case), that allows to obtain asymptotical solutions of original Vlasov plasma equations as inseparable sets of coupled oscillatory modes (but not a single wave like $\exp(-i\omega t + ikx)$). The mode parameters are defined in this case by doublepoles (ω_n, k_n) of Laplace image $E(\omega_n, k_n)$ of electrical field E(x, t) or, correspondingly, by double-zeros of inverse expression $1/E(\omega_n, k_n) = 0$. This method, in contrast with classical one, allows to obtain the whole set of oscillatory modes for every concrete problem. It leads to some new ideology in the theory of plasma oscillations, the latter's being a set of coupled oscillatory modes (characterized by pairs (ω_n, k_n) and amplitudes) and depending not only on the intrinsic plasma parameters, (as is the case in classic theory), but also on mutually dependent self-consistent initial and boundary conditions and on method of plasma oscillations excitation.

PACS numbers: 52.25 Dg; 52.35 Fp.

Key words: Landau damping; plasma oscillations; electron and ion waves; plasma dielectric susceptibility; plasma dispersion equation.

More than 50 years ago there appeared the work of L. Landau¹ with an original asymptotical solution of Vlasov equations for collisionless Maxwellian plasma in the form of a single travelling damping electromagnetic wave (longitudinal or transversal one). In this way there appeared some novel conceptions in plasma physics: Landau damping and Landau rule of bypassing around poles in calculating some indefinite Cauchy-type integrals along the real axis, which arise in standard procedures of deriving dispersion equation after substituting in Vlasov equations the wave exp $(-i\omega t + ikx)$. Landau searched for a solution f(t) of initial problem for perturbation of electron distribution function and electrical field in a form $\sim \exp(ikx) \cdot f(t)$, where k is real wave number, using the method of Laplace transformation of f(t). By means of analytical continuation of Laplace image of f(t), that is f_p , in complex transform parameter $p \equiv -i\omega$ he has obtained an asymptotical solution corresponding to a pole p_n of the analytically continued function in the form of a slowly damping travelling wave with complex frequency $\omega \equiv \omega_0 - i\delta$. This work of Landau was attributed to the rank of outstanding discoveries in the physics of plasma³.

The most surprising turned out the fact that the substitution of this solution back into the Vlasov equations results in the dispersion equation which has no solutions at $\omega_0 \neq 0$ for any δ (see in more details^{4,5}). Yet, it was decided that the right one is not the dispersion equation obtained in this way, but some *other* dispersion equation with evident natural additions in the Vlasov equations. These additions arise at calculation of indefinitely divergent Cauchy integral in velocity v_x with the pole $\omega = kv_x$ (x is direction of wave movement, v_x is x-component of particle velocity and integration along real v_x -axis is carried out with bypassing this pole along

^{*}Correspondence should be addressed to: Soshnikov V. N., Krasnodarskaya str., 51-2-168, Moscow 109559, Russia.

the half-circle contour in the complex plane v_x).

These results came into the all plasma physics text-books (see, f.e.⁶⁻⁹). There have appeared experimental works^{10,11,16–18}, which seemed to have proved Landau damping. Although sometimes there appeared some doubts about rightness of these theoretical results^{12–15}, all of them were searching for either something mathematically incorrect in Landau's derivation or some diversifications, and also philosophical justifications like the causality principle or expansion in asymptotically divergent series in small ratio δ/ω_0 , and so on. To date Landau's results are considered as irrefragable, and they compose an essential part of the theoretical plasma physics^{3,6-9}. Attempts to find mathematical incorrectnesses in Landau's derivation, in particularly, in^{4,5}, have failed (see²). Apparently, his derivation is really mathematically irreproachable.

These results are looked now in a quite another light after the successful attempt to use in solving Vlasov equations two-dimensional (in coordinate x and time t) Laplace transforma $tion^{2,4,5}$. In this case the asymptotical solution which must satisfy the original Vlasov equations after its substitution there, is found as the sum of oscillatory modes $\sum_{n} a_n \exp(-i\omega_n t + ik_n x)$ which does not reduce to the simple product $f(t)\phi(x)$. And besides, coefficients a_n , frequencies and wave numbers being found not from the dispersion equation $\epsilon(\omega, k) = 0^{7,8}$ (ϵ is generalized plasma susceptibility), but as some values determining residues and double-poles of the Laplace transform $E(\omega, k)$ or, correspondingly, double zeros (in ω and k) of the inverse expression $1/E(\omega, k)$. In this case the solution, including ω_n , k_n , is determined not only by inherent parameters of plasma (as it is in the Landau theory), but in the same extent by initial as well as boundary conditions and by the ways of excitation of plasma oscillations/waves. The initial and boundary conditions are not arbitrary, but are connected by conditions of the field selfconsistency (supposing finiteness) in Vlasov equations, so a dividing to the so called properly initial or properly boundary problems in Landau theory is incorrect. In this way, even at the initial condition of perturbation of the distribution function $f_1(x, t = 0, \vec{v}) = \alpha(\vec{v}) \exp(ikx)$ initial single-mode solution is factorized in time, and other oscillatory modes appear inevitably in the asymptotical solution. The coupled modes which are constituents of the solution include Langmuir waves, standing waves, damping and non-damping waves and oscillations, but do not include whatever specific Landau damping². Vlasov equations must be satisfied indeed only by the totality of the modes of $f_1(x, t, \vec{v})$ and $\vec{E}(x, t)$, but not by whatever mode separately. Besides that, from the condition of self-consistency (finiteness) of solution it follows that the modes of f_1 are proportional to the amplitude of electrical field with the natural result that there are no oscillations in the absence of restoring forces of electrical field (when the latter is damping). Properties of the system of linear equations which connect asymptotical amplitudes of E(x,t) and $f_1(x,t,\vec{v})$ determine the form of solution: one, or more independent sets of modes including a possible case of unavoidable exponentially divergent modes (probably, it is the peculiar case of an external non-self-consistent source).

Strictly speaking, in order to keep self-consistency in advance, one could after setting boundary and initial conditions for the distribution function f_1 also obtain such conditions for electric field E in accord with Poisson equation for $\partial E/\partial x$.

Whether these results being contrary to all the educational textbooks on plasma physics are right?

The answer is, that Landau derivation appears only as some mathematically right part of the classical proof by reductio ad absurdum: from some arbitrary supposition, what would be, if the solution of Vlasov equations should have the form $f(t) \exp(ikx)$, follows at asymptotical limit $f(t) \to \exp(-\omega t)$ with Landau's ω , k (and also at any other values), that these solutions do not satisfy the original Vlasov equations, consequently, the initial supposition about the assumed form of the solution was wrong. It is a strikingly simple and classical solution of the "paradox of Landau", including some mysterious appearance of non-damping oscillations of $f_1(x, t, \vec{v})$ in the absence of any restoring force (electrical field $E \to 0$ according to Landau damping!)⁶.

However the contradiction in Landau's logic is even more profound. He proceeds from the Laplace image f_p^+ (correspondingly, electrical field E_p^+) in the upper half-plane ω , where poles in ω should lead to exponentially divergent solutions, what appears illogical. But in the upper half-plane poles are absent, so the dispersion equation has no solutions. But Landau's analytical continuation in ω into the lower half-plane leads to poles in ω , which contrary to the initial supposition correspond to exponentially damping solutions (what already is paradoxical). Besides, these solutions satisfy some another dispersion equation (the original dispersion equation which follows from substitution of a travelling wave into Vlasov equations, as it was already said before, is not satisfied in this case). On the contrary, it should be more logical to proceed from the Laplace images in the lower half-plane f_p^- , E_p^- (different from f_p^+ , E_p^+) with damping solutions. However, in this case the poles exist only for analytical continuation of f_p^- , E_p^- into the upper half-plane and correspond, contrary to the initial supposition, to divergent solutions.

So, our basic principle is to find the correct solution of original Vlasov equations, but not to find some modified equations for fitting them to be satisfied by Landau damping solution.

In this way, the dispersion equation in the form $\epsilon(\omega, k) = 0$, as well as generally the method of one-mode derivation of dispersion equation have rather limited applicability. The suggested here method of two-dimensional Laplace transformation gives effectively the total set of coupled oscillatory modes in every concrete case, and it can lead to many new results not only in plasma physics (with revision of the results obtained on the base of Landau "bypassing around poles"), but also in gasdynamics and other fields. One can also conclude that with the new method a completely new ideology of oscillatory solutions arises which is quite different with respect to Landau's sophisticated conceptions, "predicting" physical phenomenon ("Landau damping"), which probably does not exist in nature.

Experimental "Landau damping" (see^{3,10,11,16–18}) may be mimicked by a damping caused by finiteness of plasma tube diameter (in the real plasma slab modelling) or/and non-selfconsistent (external) field term of the plasma waves excitation (in the field term of kinetic equation), or more really by a non-zero source term in the right-hand side of the more general kinetic equation(s) (cf.³), according to some concrete experimental conditions (Langmuir probes type, excitation transparent grids or compact flat or wire electrodes etc.). So, this experimental damping must satisfy some other physical equations, not the original Vlasov equations. In this case the actual problem is the analysis and physical groundings of other different modifications of these equations in their conformity with the different concrete experimental conditions, including a distinct demarcation between the electric external (non-self-consistent) and intrinsic Poissonian (self-consistent) fields. Such plasma wave equations can be solved then with the proposed method of two-dimensional Laplace transformation.

ACKNOWLEDGMENTS

I am very grateful to Dr. A. P. Bakulev for his active and very constructive edition and help in amendment of this paper as well as his realization of electronic sending procedures.

¹ Landau L. D., J. Phys. (USSR), **10** (1946) 25

² Soshnikov V. N. Preprint physics/9706041 (see http://xxx.lanl.gov/e-print/physics/9706041)

³ Kadomtsev B. B., Uspekhi Fiz. Nauk, **95** (1968) 111 (in Russian)

⁴ Soshnikov V. N., Manuscript, Dep. VINITI (1990) No. 2982-B90;

See also: http://xxx.lanl.gov/e-print/plasm-ph/9511001

⁵ Soshnikov V. N., J. Moscow Phys. Soc., **6** (1996) 279

⁶ Clemmow P. C., Dougherty J. P., "Electrodynamics of Particles and Plasmas", 2nd ed., Addison-Wesley, NY (1990); (Rus. transl. Moscow, Mir, 1996)

- ⁷ Lifshits E. M., Pitaevsky L. P., "Physical kinetics", Nauka, Moscow (1979) (in Russian)
- ⁸ Alexandrov A. F., Bogdankevich L. S., Rukhadze A. A., "Foundations of Electrodynamics of Plasma", 2nd ed., Vysshaya Shkola, Moscow (1988) (in Russian)
- ⁹ Kadomtsev B. B., "Collective Phenomena in Plasma", 2nd ed. Nauka, Moscow (1988) (in Russian)
- ¹⁰ Malmberg I. H., Wharton C. B., Phys. Rev. Lett., **17** (1966) 175
- ¹¹ Malmberg I. H., Wharton C. B., Drummond W. E., Plasma Phys., **1** (1966) 485
- ¹² Weitzner H., Phys. Fluids, **6** (1963) 1123
- ¹³ Sagan D., Am. J. Phys., **62** (1994) 450
- 14 Alexeff I., Osamu I., IEEE Trans. Plasma Sci., ${\bf 6}$ (1978) 212
- ¹⁵ Alexeff I., Rader M., Int. J. Electronics, **68** (1990) 385
- ¹⁶ Malmberg I. H., Wharton C. B., Phys. Rev. Lett., **19** (1967) 775
- ¹⁷ Derfler H., Simonen T. C., 8-th Intern. Conf. Phenom. in Ionized Gases, Contrib. Pap., Vienna, p. 335 (1967)
- ¹⁸ Van-Hoven G., Phys. Rev. Lett., **17** (1966) 169