

DETERMINATION OF THE FINE STRUCTURE CONSTANT BY π

A. H. Anastassov
University of Sofia, Faculty of Physics *

October 31, 1997

Abstract

In the present report the author presents a simple and systematically defined formula for the fine structure constant based only on the number π . The difference between the suggested value and the currently known experimental one is about 60 times smaller than the uncertainty interval.

The fundamental physical constants have attracted the attention of physicists and mathematicians for long time. Especially challenging is the idea that they can be reduced to mathematical constants (Hilbert [1], Einstein [2]).

The greatest is the challenge offered by the dimensionless constant α of Sommerfeld [3], currently known as the fine structure constant or as electromagnetic constant. As it is well known, it is determined by the dimensionless relation $e^2/\hbar c$, where e is the elementary electric charge, \hbar is Plank's constant divided by 2π , and c - the absolute velocity (the speed of light). Its reciprocal value $1/\alpha$ is very closed to the integer 137 and this fact has grown the attention of many researches starting with Sommerfeld himself. Eddington's special interest towards this issue is well known, too [4]. Similarly Dirac assumed that some constants, including the fine structure constant α ,

*petersl@phys.uni-sofia.bg

where build up of simple quantities like 4π [5]. In this line of research were published interesting formulae such as:

$$\begin{aligned}(2^4 5! / \pi^5)^{1/4} (8\pi^4 / 9) &= 137,036\,082 \\ 108\pi(8/1843)^{1/6} &= 137,035\,915 \\ 4\pi^5/9 + 37/36 &= 137,036\,527 \\ (137^2 + \pi^2)^{1/2} &= 137,036\,016\end{aligned}$$

considered in more details in [6].

The currently recommended by The National Institute of Standards and Technology (USA) value of Sommerfeld's constant and its reciprocal are [7]:

$$\alpha_R = \mathbf{7,297\,353\,08(33)} \times 10^{-3}, \quad (1)$$

$$\alpha_R^{-1} = \mathbf{137,035\,989\,5(61)}. \quad (2)$$

In this note, following Dirac's idea I will show that α could be presented with an absolute (for the moment) accuracy in **a simple and systematical way only through the numbers 1, 2, 3, 4 and π** . In order to do that I will introduce the quantities:

$$\begin{aligned}\alpha_0 &:= 1/\sqrt{\frac{1}{2}\left(\frac{1}{2}+1\right)}(4\pi)^2, \\ \rho &:= \alpha_0/2\pi(\pi^2-2\pi), \\ L &:= 2\pi\rho, \\ S &:= \pi\rho^2, \\ V &:= \frac{4}{3}\pi\rho^3.\end{aligned}$$

Proposition: the number

$$\alpha_\rho := \alpha_0/(1 + L - 2S + 3V/2)$$

has the value

$$\alpha_\rho = \mathbf{7,297\,353\,085\,4...} \times 10^{-3} \quad (3)$$

and its reciprocal value is

$$\alpha_\rho^{-1} = \mathbf{137,035\,989\,392...}. \quad (4)$$

The comparison shows that the number (3) not only is in the limits of the error of the experimental value (1) but hits its central value. The same is valid, of course, for the reciprocal values (4) and (2).

It comes to prove that Sommerfeld's constant can in fact be presented through simple mathematical quantities, based on the number π .

References

- [1] Hilbert D., *Die Grundlagen der Physik*, Nachrichten K. Gessellschaft Wiss Göttingen, Math.-phys. Klasse, 1915, Heft3, p.395.
- [2] Einstein A., *Autobiographical Notes*. In: "Albert Einstein - Philosopher-Scientist", (ed. P.A.Schilpp, Evanston, Illinois, 1949).
- [3] Sommerfeld A., *Atombau und Spectrallinien*, (Friedr. Vieweg & Sohn, Brounschweig, 1951).
- [4] Eddington A.S., *Fundamental Theory*. (Cambridge Univ. Press, London, 1948).
- [5] Dirac P.A.M., *Directions in Physics*, (Wiley, New York, 1978).
- [6] Eagles D.M., Intern. J. Theor. Phys., 1976, **15**, No4, p. 265.
- [7] Cohen E.R. and B.N. Taylor, J. of Res. of the Nat. Bureau of Standards, **92**, 85, (1987).