

CAN MECHANICALLY EXTENDED SOURCES OF RADIATION IN CLASSICAL ELECTRODYNAMICS SOLVE ALL PROBLEMS OF RADIATING POINT-LIKE SOURCES?

Alexander A. Vlasov

High Energy and Quantum Theory
Department of Physics
Moscow State University
Moscow, 119899
Russia

On simple example is shown that a model of mechanically extended source of radiation in classical electrodynamics indeed has some problems with radiation reaction force. So one cannot, in general case, state, that the present situation with classical dynamics of mechanically extended sources of radiation is "very satisfactory".

03.50.De

In classical electrodynamics there are known problems of radiation reaction force for point-like sources of radiation: preacceleration, runaway and other pathological (unphysical) solutions.

In the literature one can find the opinion that if the finite size of a radiating source is fully taken into account all the above problems can be solved.

In this paper we examine the well known model of "nonrelativistically rigid charged sphere" to show that nevertheless there remain problems with radiation reaction force.

It should be mentioned that "extended (mechanically) electron models" were proposed long time ago in the works of Abraham, Lorentz, Sommerfeld and others (for good reviews see, for ex., [1,2,3,4]).

Thus in Sommerfeld works [5] was derived the expression of self-force acting on "nonrelativistically rigid charged sphere", i.e sphere with radius a , its center moving along trajectory $\vec{R}(t)$, with total charge Q and charge density (in laboratory reference frame)

$$\rho(t, \vec{r}) = \frac{Q}{4\pi a^2} \delta(|\vec{r} - \vec{R}| - a)$$

In the case of shell rectilinear motion in [7] was given for the self-force

$$\vec{F}_{self} = \int dV \rho (\vec{E} + [\vec{v}, \vec{H}]/c)$$

the following expression:

$$\begin{aligned} F_{self} = & \frac{Q^2 c}{4a} \int_{-1}^{+1} dx \int_{t_1}^t dt' \frac{ax + L'}{[a^2 + 2axL' + (L')^2]^{3/2}} \\ & + \frac{Q^2}{4a} \int_{-1}^{+1} dx \frac{x - v/c}{a(1 - xv/c)} \\ & - \frac{Q^2}{4a} \int_{-1}^{+1} dx \frac{N_1 - v_1/c}{[a^2 + 2axL_1 + (L_1)^2]^{1/2} (1 - N_1 v_1/c)} \end{aligned} \quad (1)$$

here

$$L' = R(t) - R(t'), \quad L_1 = R(t) - R(t_1), \quad v = dR(t)/dt, \quad v_1 = v(t_1),$$

$$N_1 = \frac{ax + L_1}{[a^2 + 2axL_1 + (L_1)^2]^{1/2}}$$

and t_1 is the solution of the equation (equation of retardation):

$$t_1 = t - [a^2 + 2axL_1 + (L_1)^2]^{1/2}/c - a/c,$$

After integration by parts, expression (1) can be put in the form

$$\begin{aligned} F_{self} = & \frac{Q^2}{4a^2} \left(\frac{1 - \beta^2}{\beta^2} \ln \frac{1 + \beta}{1 - \beta} - \int_{L_-}^{L_+} \frac{dL_1}{L_1} \frac{1 - \beta^2(t_1(L_1))}{\beta^2(t_1(L_1))} \right) \\ & + \frac{Q^2 c}{4a} \int_{-1}^{+1} dx \int_{t_1}^t dt' \frac{\dot{v}'}{v'^2 [a^2 + 2axL' + L'^2]^{1/2}} \end{aligned} \quad (2)$$

here

$$L_{\pm} = L_1(x = \pm 1), \quad v' = v(t'), \quad \beta = v/c$$

With (2) equation of shell motion is

$$m \frac{d}{dt}(\gamma v) = F_{self} + F_{ext} \quad (3)$$

here m is the total mass of the shell, F_{ext} - some external force.

Let us mention some features of the eq.(3).

1). If $F_{ext} = 0$, then eq. (3) has the trivial solution

$$v = v_0 = const$$

because the expression (1,2) for the self-force identically vanishes.

2). But this solution is not unique. An example of so called radiationless motion of a rigid sphere was given in [6].

Now we consider small deviations from shell uniform rectilinear motion for linear analysis of stability of such motion.

For this goal it will be convenient to use for the self-force the Sommerfeld expression [2], which is equivalent to (1) (this can be shown spreading with the help of step functions limits of integration over t' in the double intergal in (1) to the whole axis $(-\infty, +\infty)$ and changing the order of integration over x and t'):

$$F_{self} = \frac{Q^2}{4a^2} \left[-c \int_{T^-}^{T^+} dT \frac{cT - 2a}{L^2} + \ln \frac{L^+}{L^-} + \left(\frac{1}{\beta^2} - 1 \right) \ln \frac{1 + \beta}{1 - \beta} - \frac{2}{\beta} \right] \quad (4)$$

here $cT^\pm = 2a \pm L^\pm$, $L^\pm = |R(t) - R(t - T^\pm)|$, $L = |R(t) - R(t - T)|$.

Let us take

$$R(t) = R_0 + vt + \delta(t), \quad R(t-T) = R_0 + v(t-T) + \delta(t-T), \quad L = vT + \delta(t) - \delta(t-T),$$

here $\delta(t)$ - small perturbation, $v \neq c, \neq 0$.

Then in linear approximation the self-force (4) reads

$$F_{self} = \frac{Q^2}{4a^2} \left[\frac{4\dot{\delta}}{\beta^2} \left(1 - \frac{1}{2\beta} \ln \frac{(1 + \beta)}{(1 - \beta)} \right) - \frac{2c}{v^3} \int_{T^-}^{T^+} dT \frac{cT - 2a}{T^3} \delta(t - T) \right] \quad (5)$$

here $T^\pm = \frac{2a}{c(1 \mp \beta)}$, and the equation of shell motion is

$$m\gamma^3 \ddot{\delta} = F_{self} \quad (6)$$

Solution of eq. (6) we take as $\delta \sim \exp(pt)$. Then the parameter p must obey the following equation

$$p^2 = \frac{Q^2}{a^2 m \gamma^3} \left[\frac{p}{c \beta^2} \left(1 - \frac{1}{2\beta} \ln \frac{(1+\beta)}{(1-\beta)} \right) - \frac{c}{2v^3} \int_{T^-}^{T^+} dT \frac{cT - 2a}{T^3} \exp(-pT) \right] \quad (7)$$

Now consider two boundary values of shell velocity.

For analysis of stability of the solution $v_0 = 0$ we examine the approximation of small velocities. This yields for the self-force known expression [2]:

$$F_{self} = \frac{Q^2}{3ca^2} [v(t - 2a/c) - v(t)] \quad (8)$$

and the eq. of motion

$$m\dot{v} = \frac{Q^2}{3ca^2} [v(t - 2a/c) - v(t)] \quad (9)$$

does not possess growing with time solutions.

For analysis of stability of the solution $v \rightarrow c$ it will be convenient to use the expression (2) for the self-force. Let's put in (2) $R(t) = R_0 + ct + \delta(t)$, then the main contribution to the eq. of motion gives the first term in (2) and

$$F_{self} \sim \dot{\delta} \ln |\dot{\delta}|,$$

as the L.H.S. of eq. of motion takes the form $\sim \ddot{\delta} (-\dot{\delta})^{-3/2}$, then full eq. of motion is

$$\ddot{\delta} = -k (-\dot{\delta})^{+5/2} \ln |\dot{\delta}|$$

here k is positive parameter.

This eq. has the solution

$$\delta \sim t^{1/3} \ln^{-2/3} t, \quad t \rightarrow \infty$$

that is the uniform motion of the sphere with $v \rightarrow c$ is relatively stable in that sense that

$$\dot{\delta}/c \rightarrow 0, \quad \delta/(R = ct) \rightarrow 0, \quad t \rightarrow \infty \quad (10).$$

Thus there can be solutions asymptotically tending to light cone.

More general analysis of eq.(7) will be done in the next preprint (under preparation).

3). There is one more problem, actual to all intergal equations. If the small perturbation $\delta(t)$ is zero for $t < 0$, then the Laplace transformation of (5,6) yields the equation for eigenvalues λ of this integral eq. equal to the eq. for p (7).

Thus at least for $\lambda a/c \ll 1$ this eq. is solvable and eigenvalue is not zero.

Then, as the theory of intergal equations tells, not for all forms of external forces the solution of the inhomogeneous equation (3) with $F_{ext} \neq 0$ exists!

The similar situation occurs for rotating shell - see [8].

Thus our conclusion is:

a model of mechanically extended source of radiation in classical electrodynamics with retardation indeed possesses some problems with radiation reaction force.

So one cannot, in general, state, that the present situation with classical dynamics of mechanically extended sources of radiation is "very satisfactory".

And to solve problems of radiation reaction force one must search for new ideas and approaches.

REFERENCES

1. T.Erber, Fortschr.Phys., 9, 342 (1961).
2. P.Pearle, in *Electromagnetism*, ed. D.Tepliz, Plenum, NY, 1982, p.211.
3. A.D.Yaghjian, *Relativistic Dynamics of a Charged Sphere*, Lecture Notes in Physics, 11, Springer, Berlin, 1992.
4. F.Rohrlich, Am.J.Physics, 65(11), 1051 (1997).
5. A.Sommerfeld, Gottingen Nachrichten, 29 (1904), 363 (1904), 201 (1905).
6. G.A.Schott, Phil.Mag., 15, 752 (1933).
7. Alexander A.Vlasov, physics/9711024.
8. Alexander A.Vlasov, physics/9801017.