All-electron Dirac-Coulomb and RECP calculations of excitation energies for mercury atom with combined CI/MBPT2 method.

N. S. Mosyagin^{*}, M. G. Kozlov, and A. V. Titov

Petersburg Nuclear Physics Institute, Gatchina, Petersburg district 188350, RUSSIA (November 14, 2018)

Abstract

Calculations of transition energies between low-lying states of mercury atom are performed in the frame of combined CI/MBPT2 method. Results of all-electron relativistic calculations (using the Dirac-Coulomb Hamiltonian) are compared with experimental data and results of other four-component calculations. The results of the RECP calculations are compared with the corresponding all-electron results in order to estimate accuracy of different RECPs. Contributions from correlations in different shells to the calculated excitation energies as well as effects of basis set truncation at different orbital angular momenta, nuclear models, errors in gaussian approximation of the GRECP components are reported. Analysis of the obtained results shows that at least 34 external electrons of mercury atom should be correlated and the one-electron basis set should contain up to h angular momentum functions in order to reach a reliable agreement with experimental data within 200 cm⁻¹. It is concluded that correlations of the 4f electrons can be efficiently taken into account for 20 electron GRECP at the generation stage.

SHORT NAME: CI/MBPT2 calculations of mercury.

KEYWORDS FOR INDEXING: Configuration Interaction, Many Body Perturbation Theory, correlation structure (electronic structure), excitation energies (transition energies), Relativistic Effective Core Potential (pseudopotential), four-component calculations, mercury, heavy atoms.

I. INTRODUCTION

During the last few years, a considerable number of papers devoted to electronic structure calculations of heavy atoms have been published (e.g., see [1-3]) that is not only due to experimental requirements but because they are good test systems to check or estimate accuracy of different approximations before using them in more expensive molecular calculations.

In paper [4], Generalized Relativistic Effective Core Potentials (GRECPs) were tabulated for atoms Hg through Rn and were tested in numerical Hartree-Fock (HF) calculations by comparison with the all-electron Dirac-Fock (DF) and other RECP ones. However, the question

of quality of the GRECPs for description of correlation effects was not clarified yet. Whereas paper [5] contains an answer on this question from theoretical point of view, the present paper is devoted to our correlation structure calculations for mercury atom. For this atom, the RECPs were generated by other groups [6,7] where the same number of electrons (20) was explicitly included into calculations as in the case of the GRECP [4]. Therefore, one

should expect about the same computational expenses in calculations with

all these RECPs and the comparison of their accuracy appears to be of practical interest.

II. METHODS AND BASIS SETS

The GRECP method was described in details in papers [8,4,5]. The main distinguishing features of this method are presence of non-local terms with the projectors on the outercore (OC) pseudospinors together with the standard semi-local ones in the effective potential operator and generation of the effective potential components for smoothed pseudospinors which may have nodes [9].

Theory of the CI/MBPT2 method is presented in papers [2]. In this method, the correlations in the valence (V) region of an atom which are the most important ones are treated by the Configuration Interaction (CI) method (which is able to provide excellent results for the small number of correlated electrons) whereas the relatively small contributions to the considered low excitation energies from the large number of the core-valence correlations are taken into account with the help of less expensive second order Many Body Perturbation Theory (MBPT2).

The program (in the jj-coupling scheme) for all-electron relativistic four-component CI/MBPT2 calculations (using the Dirac-Coulomb Hamiltonian) was modified to make possible two-component RECP calculations (with the nonrelativistic kinetic energy operator and relativistic j-dependent potentials). This program allows one to use two different basis sets of numerical spinors at the CI and MBPT2 calculation stages for description of the correlations taking into account a space separation of the core and valence regions. Basis functions for the present calculations were obtained from numerical SCF (DF or HF) calculations of the corresponding spinors for positive ion states. For example, the [4,4,3,2,1] basis set for the 2 electron valence CI (2e-CI) was derived from the calculations for the following nonrelativistically averaged configurations of Hg: $5d^{10}6s^2$, $[5d^9]7s^1$, $[5d^6]8s^1$, $[5d^3]9s^1$,

 $[5d^{10}6s^1]6p^1$, $[5d^9]7p^1$, $[5d^8]8p^1$, $[5d^7]9p^1$, $[5d^9]6d^1$, $[5d^6]7d^1$, $[5d^3]8d^1$, $[5d^8]5f^1$, $[5d^7]6f^1$, $[5d^4]5g^1$, where the shells in the square brackets were frozen in the calculations and the 1s-5p shells are dropped out for brevity. For convenience of comparison, a $[k_s, k_p, k_d, k_f, k_g, k_h]$ basis set for the case of N correlated electrons will refer here to a basis set consisting of $n_ss_{1/2}, \ldots, (n_s + k_s - 1)s_{1/2}; n_pp_{1/2}, \ldots, (n_p + k_p - 1)p_{1/2}; n_pp_{3/2}, \ldots, (n_p + k_p - 1)p_{3/2}; n_dd_{3/2}, \ldots, (n_d + k_d - 1)d_{3/2}; n_dd_{5/2}, \ldots, (n_d + k_d - 1)d_{5/2}; n_ff_{5/2}, \ldots, (n_f + k_f - 1)f_{5/2}; n_ff_{7/2}, \ldots, (n_f + k_f - 1)f_{7/2}; 5g_{7/2}, \ldots, (4 + k_g)g_{7/2}; 5g_{9/2}, \ldots, (4 + k_g)g_{9/2}; 6h_{9/2}, \ldots, (5 + k_h)h_{9/2}; 6h_{11/2}, \ldots, (5 + k_h)h_{11/2}$ spinors where $n_s = 5$ if $N \ge 20$, $n_p = 5$ if $N \ge 18$, $n_d = 5$ if $N \ge 12$, $n_f = 4$ if $N \ge 34$, otherwise $n_s = 6$, $n_p = 6$, $n_d = 6$, $n_f = 5$. Additional information on the used basis sets can be found on http://www.qchem.pnpi.spb.ru. The above listed basis set has provided the lowest total energy in our CI calculations as compared with other tested

similar-sized basis sets. More important advantage of this basis set is a possibility of one-to-one comparison between results of the all-electron

and RECP calculations not only for large (close to full) basis set sizes

but for the small ones as well. In fact, the calculated RECP errors depend only slightly on the size of such basis sets.

III. RESULTS AND DISCUSSION

Series of the correlation structure calculations were performed for the number of correlated electrons varied from 2 to 34 and one-electron basis sets truncated at the orbital quantum numbers from 2 to 5. The all-electron relativistic calculations were implemented for two nuclear models: a point nucleus and an uniformly charged ball with $1.334 \cdot 10^{-4}$ a.u. radius. The GRECP calculations were made with both the numerical GRECP components and their gaussian expansions from [4]. Moreover, 34 electron GRECP variant (where the outercore 4f, 5s, 5p, 5d and valence 6s, 6p shells of mercury atom are explicitly treated in calculations) was generated and tested in these calculations.

The results of the 2e-CI calculations (where all the possible excitations of 2 valence electrons were considered and the core $1s_{1/2}$ - $5d_{5/2}$ spinors were frozen from the SCF calculations of the ground state with the $6s^2$ configuration) are presented in tables I and II for [8, 8, 7, 6, 5, 4] and [4, 4, 3, 2, 1] basis sets. One can see from these tables that the use

of the [4, 4, 3, 2, 1] basis set allows one to describe adequately the V-V

correlations and that the RECP errors are rather stable in respect to the basis set size variation. Whereas the correlations only in the valence region are considered, the 20 electron GRECP is about 8 times more accurate than the 20 electron RECP of Ross *et al.* [6] and about 30 times more accurate than the 20 electron energy-adjusted PseudoPotential (PP) [7] (first of all, due to the neglect of the difference between the outercore and valence potentials in the RECPs [6,7]; see [8,5] for details).

From tables III and IV in [4] one can see that the errors of the energy-adjusted PP, Ross *et al.*'s RECP, and the GRECP for the excitations from the $6s_{1/2}^2$ state to the $6s_{1/2}^16p_{1/2}^1(J=0)$

and $6s_{1/2}^1 6p_{3/2}^1 (J = 2)$ states¹ are -667 and +347 cm⁻¹, +182 and +224 cm⁻¹, -9 and -9 cm⁻¹, correspondingly. They are in agreement with the errors from tables I and II here with an exception of the GRECP errors (because the main contribution to the GRECP errors is due to errors of reproducting the two-electron integrals rather than drawbacks of the effective potential operator).

In tables II–V, results of both the all-electron relativistic and 20 electron GRECP calculations are presented for different numbers of correlated electrons² using the equivalent basis sets. One can see that the 2e-CI (table II) gives only rough description of the correlation structure of mercury atom, the errors in energies of excitations from the ground state are $3000-6000 \text{ cm}^{-1}$ in respect to the experimental data or the most elaborated calculations in table V. Such a large

deviation is mainly due to neglect of the OC-V correlations involving the 5d shell. Their consideration (in the 12e-CI/MBPT2, table III) reduces these errors to a level of about 1000 cm⁻¹. The contributions from the correlations with the 5p and 5s shells (the 18e-CI/MBPT2 and 20e-CI/MBPT2, tables III and IV) to the excitation energies are 400– 600 cm⁻¹ and about -300 cm⁻¹, correspondingly. However, these contributions are mainly compensated and the total contribution from the correlations with both these shells is only 100–300 cm⁻¹. The correlations with the 4f shell (the 34e-CI/MBPT2, table V) give an essential contribution that is 600–700 cm⁻¹. Our final all-electron results for transitions between the first four states are within 200 cm⁻¹ with the experimental data [10] and the Relativistic Coupled Cluster (RCC) calculation results of Eliav *et al.* [3]. A relatively large deviation for the last state is rather due to the MBPT2 approximation than due to basis set incompleteness or correlations in more inner shells. Our estimates show that the contribution from the correlations with the 4d shell to the excitation energies is of order of 100 cm⁻¹.

When only the V-V correlations are considered (tables I and

II), the GRECP errors in reproducting the all-electron results are within 30 cm^{-1} . This is a good confirmation to our previous estimates [4] that the GRECP describes the electronic structure in the valence region with a high accuracy. However, the consideration of the correlations in the outercore region (tables III and IV) leads to an increase of these

errors (due to a rather large smoothing region for the nodeless outercore 5s, 5p, 5d pseudospinors³) up to 200 cm⁻¹. These results are in a good agreement with that from table III in paper [4] where a significant increase in the GRECP errors can be observed for the case of excitations from the 5d shell. One should expect that the contribution from the correlations

¹ Analysis for the case of the $6s_{1/2}^1 6p_{1/2}^1 (J = 1)$ and $6s_{1/2}^1 6p_{3/2}^1 (J = 1)$ SCF states is more complicated because they are strongly mixed in the CI calculations.

² Electrons are included in the calculations in the following order: $6p, 6s; 5d; 5p; 5s; 4f; \ldots$

³ We suggest that these GRECP errors can be seriously reduced due to more artificial generator configuration selection and smoothing procedure and with the help of corrections to the GRECP operator (see [5], subsection 4.5) because these errors arise from electronic structure reorganization in the outercore region.

the outercore region to low excitation energies and GRECP errors in their reproducting will be reduced when passing from Hg toward Rn.

The errors of different RECPs for the case of 20 correlated electrons are presented in table IV. The errors of the GRECP are only about 4 times less than that of Ross *et al.*'s RECP [6] and about 7

times less than that of the energy-adjusted PP [7] for the same number of electrons explicitly included in calculations. This is also in agreement with the results from table III in paper [4] where the ratio between GRECP errors and errors of the other RECPs was reduced from one order of magnitude (when excitations only for the valence electrons were under consideration) to 1.5-2 times (for the case of excitations from the outercore shells described by nodeless pseudospinors). In fact, additional changes in the RECP errors when the OC-V correlations are taken into account (tables II and IV) are within 200 cm⁻¹ for the used GRECP variants and within 400 cm⁻¹ for the other RECPs. The main reason is the spinor smoothing which has similar features for all these RECPs.

The data from the MRCI calculations with the energy-adjusted PP using the CIPSO method for the transitions between the first four states from table 6 in paper [7] are within 100 cm^{-1} with the experimental data but the 20 electron PP does not take into account contributions from the correlations with the 4f shell (which are up to

700 cm⁻¹) and the used basis set does not contain functions with h orbital momentum (their contributions are up to 300 cm⁻¹). Therefore, these data are results from cancelation of a few contributions: PP errors (e.g., the $6s_{1/2}^16p_{1/2}^1(J=0)-6s_{1/2}^16p_{3/2}^1(J=2)$ splitting is overestimated about 1000 cm⁻¹ by the energy-adjusted PP because of the features of the spin-orbit simulation in the *LS*-based variant of the energy-adjusted scheme), a neglect of the correlations with the 4f shell, a basis set incompleteness, etc. The estimation of error for these data within 5% [7] appears to be correct. The data in paper [11] were obtained with the 12 electron RECP of Ross *et al.* which differs from the used here 20 electron RECP.

One can see from tables I, II, and IV that the errors of the 20 and 34 electron GRECPs are of the same order of magnitude because the main distinctions between these GRECP variants are

inclusion of the 4f electrons in calculations with the 34 electron GRECP and smoothing the 5f spinors for the 20 electron GRECP. However, the 34 electron GRECP allows one to take into account explicitly the correlations with the 4f shell which are important for an agreement with the experimental data within 200 cm⁻¹. As one can see from table V, the 34 electron GRECP errors are within 200 cm⁻¹ in this case.

The basis set truncation effect at different orbital quantum numbers on energies of excitations from the ground state can be observed in tables V and VI for the 34e-CI/MBPT2 calculations. It is clear that (spd) correlation basis set⁴ does not allow one to take into account the correlations with the 5*d* shell properly. This leads to the excitation energies (table VI) rather close to that from the 2e-CI calculations. Addition of functions

5

⁴ The [12, 12, 11, 1] basis set contains only such functions with f orbital momentum which correspond to the $4f_{5/2}$ and $4f_{7/2}$ spinors.

with f orbital momentum is crucial for description of these correlations and leads to an increase in the excitation energies on 2000–4000 cm⁻¹. The excitation energies for the (spdf) basis set are within 900 cm⁻¹ from that for the (spdfgh) basis set (tables V and VI). In turn, (spdf) basis set is inadequate for description of the correlations with the 4f shell. The results for this basis set are rather close to that from the 20e-CI/MBPT2 calculations in table IV. Addition of functions with g orbital momentum is necessary for the correct description of these correlations and gives the contribution 300–600 cm⁻¹ to the excitation energies (table VI) whereas addition of h functions contributes about 300 cm⁻¹ (table V). Our test calculations showed that the contribution from functions with i orbital momentum is of order 50 cm⁻¹.

One can see from comparison between the 6 and 7 columns in tables I, II, and IV that the errors

due to the gaussian approximation of the GRECP components are approximately one order of magnitude less than the errors of the numerical GRECP. As one can see from tables I, II, IV, and V, the effects of different nuclear models may be neglected for accuracy within 200 cm⁻¹.

IV. CONCLUSIONS

The GRECP allows one to reproduce the electronic structure in the valence and outercore regions essentially better than the other tested RECPs for

the same number of explicitly treated electrons.

At least 34 external electrons (occupying the 4f, 5s, 5p, 5d, 6s, 6p shells) of mercury atom should be correlated and the one-electron basis set should contain up to h angular momentum functions in order to obtain a reliable agreement with experimental data for low excitation energies within 200 cm⁻¹ whereas the errors of the gaussian approximation of the GRECP components and the effects of different nuclear models are negligible for this accuracy. However, our test calculations show that the main contribution from the correlations with the 4f shell is due to the one-electron correction from the self-energy diagrams [2], therefore, this contribution can be taken into account for 20 electron GRECP with the help of the technique proposed in [5] (subsection 5.2).

ACKNOWLEDGMENTS

This work was implemented under the financial support of the Russian Foundation for Basic Research (N. M. and A. T.: grants N 96–03–33036 and 96–03–00069g; M. K.: grant N 98–02–17663).

REFERENCES

- * E-mail: Mosyagin@hep486.pnpi.spb.ru for correspondence; http://www.qchem.pnpi.spb.ru
- E. Eliav, U. Kaldor, Y. Ishikawa, M. Seth, and P. Pyykko, Phys. Rev. A 53, 3926 (1996); F. Rakowitz and C. M. Marian, Chem. Phys. Lett. 257, 105 (1996); U. Wahlgren, M. Sjovoll, H. Fagerli, O. Gropen, and B. Schimmelpfenning, Theor. Chim. Acc. 97, 324 (1997); R. J. Buenker, A. B. Alekseyev, H.-P. Liebermann, R. Lingott, and G. Hirsch, Phys. Rev. A, in press.
- [2] V. A. Dzuba, V. V. Flambaum, and M. G. Kozlov, Pis'ma v ZheTF 63, 844 (1996);
 V. A. Dzuba, V. V. Flambaum, and M. G. Kozlov, Phys. Rev. A 54, 3948 (1996);
 M. G. Kozlov and S. G. Porsev, ZheTF 111, 838 (1997) [in Russian].
- [3] E. Eliav, U. Kaldor, and Y. Ishikawa, Phys. Rev. A 52, 2765 (1995).
- [4] N. S. Mosyagin, A. V. Titov, and Z. Latajka, Int. J. Quant. Chem. 63, 1107 (1997).
- [5] A. V. Titov and N. S. Mosyagin, Preprint PNPI No. 2182 (Petersburg Nuclear Physics Institute, Gatchina, St.-Petersburg district, 1997), 81 p., submitted for publication.
- [6] R. B. Ross, J. M. Powers, T. Atashroo, W. C. Ermler, L. A. Lajohn, and P. A. Christiansen, J. Chem. Phys. 93, 6654 (1990).
- [7] U. Haussermann, M. Dolg, H. Stoll, H. Preuss, P. Schwerdtfeger, and R. M. Pitzer, Mol. Phys. 78, 1211 (1993).
- [8] I. I. Tupitsyn, N. S. Mosyagin, and A. V. Titov, J. Chem. Phys. **103**, 6548 (1995).
- [9] A. V. Titov, A. O. Mitrushenkov, and I. I. Tupitsyn, Chem. Phys. Lett. 185, 330 (1991).
- [10] C. E. Moore, Circ. Natl. Bur. Stand. (U.S.) 467 (1958).
- [11] A. B. Alekseyev, H.-P. Liebermann, R. J. Buenker, and G. Hirsch, J. Chem. Phys. 104, 4672 (1996).

TABLES

TABLE I. All-electron and RECP correlation energies^a, all-electron transition energies (TE) and absolute errors (AE) of different RECPs in their reproducting from the 2 electron valence CI calculations of low-lying states of Hg for the [8, 8, 7, 6, 5, 4] basis set (in cm⁻¹).

							20 el.	20 el.
Sym-	Leading	All	-el.	34 el.	20	el.	RECP	energy-
metry	conf.	finite	point	GRECP	GR	ECP	of Ross	adjusted
(J_{parity})		nucl.	nucl.	num.	num.	gaus.	et al. ^c	PP^{d}
				Corre	elation en	$ergies^{a}$		
0_g	$6s_{1/2}^2$	-5991	-5987	-6019	-6023	-6024	-6006	-5990
0_u	$6s_{1/2}^{1'}6p_{1/2}^{1}$	-922	-922	-927	-927	-927	-926	-933
1_u	$6s_{1/2}^{1^{'}}6p_{1/2}^{1^{'}}{}^{\mathrm{b}}$	-6919	-6916	-6936	-6937	-6937	-6923	-6735
2_u	$6s_{1/2}^16p_{3/2}^1$	-1108	-1108	-1111	-1111	-1112	-1111	-1116
1_u	$6s_{1/2}^16p_{3/2}^1$ b	171	172	164	163	163	171	17
		TE	TE	AE	AE	AE	AE	AE
0_g	$6s_{1/2}^2$	0	0	0	0	0	0	0
0_u	$6s_{1/2}^{1'}6p_{1/2}^{1}$	31832	31891	-1	17	16	203	-687
1_u	$6s_{1/2}^{1}6p_{1/2}^{1}^{\mathbf{b}}$	33704	33764	-3	17	16	215	-439
2_u	$6s_{1/2}^{1^{'}}6p_{3/2}^{1^{'}}$	38073	38136	-8	18	17	248	360
1_u	$6s_{1/2}^{1}6p_{3/2}^{1^{'}b}$	50108	50162	7	34	33	245	-68

^aThe correlation energies were calculated as a difference between the total energies from the above mentioned CI calculations and the numerical SCF calculations with the frozen core $1s_{1/2}$ - $5d_{5/2}$ spinors for the corresponding terms.

^bThe $6s_{1/2}^1 6p_{1/2}^1$ and $6s_{1/2}^1 6p_{3/2}^1$ configurations are strongly mixed for these terms in the CI calculations.

^cThe RECP from Ref. [6].

^dThe PP from Ref. [7] with the corrected V_{so} by the factors (2l + 1)/2 (M. Dolg, private communication).

							20 el.	20 el.
Sym-	Leading	All	-el.	34 el.	20	el.	RECP	energy-
metry	conf.	finite	point	GRECP	GRI	ECP	of Ross	adjusted
(J_{parity})		nucl.	nucl.	num.	num.	gaus.	et al. ^c	PP^{d}
				Corre	elation en	ergies ^a		
0_g	$6s_{1/2}^2$	-5927	-5923	-5954	-5964	-5965	-5941	-5924
0_u	$6s^{1'}_{1/2}6p^{1}_{1/2}$	-909	-910	-914	-915	-915	-913	-919
1_u	$6s_{1/2}^{1^{'}}6p_{1/2}^{1^{'}}{}^{\mathrm{b}}$	-6903	-6900	-6920	-6922	-6923	-6907	-6718
2_u	$6s_{1/2}^16p_{3/2}^1$	-1097	-1097	-1100	-1101	-1101	-1100	-1106
1_u	$6s_{1/2}^{1}6p_{3/2}^{1^{'}}{}^{\mathrm{b}}$	237	238	227	221	221	235	79
		TE	TE	AE	AE	AE	AE	AE
0_g	$6s_{1/2}^2$	0	0	0	0	0	0	0
0_u	$6s_{1/2}^{1'}6p_{1/2}^{1}$	31780	31839	-1	22	21	202	-687
1_u	$6s_{1/2}^16p_{1/2}^{1-\mathrm{b}}$	33655	33716	-4	21	21	214	-438
2_u	$6s_{1/2}^16p_{3/2}^1$	38020	38083	-9	22	21	247	358
1_u	$6s_{1/2}^{1}6p_{3/2}^{1}{}^{\mathrm{b}}$	50109	50164	3	31	31	241	-73

TABLE II. All-electron and RECP correlation energies^a, all-electron transition energies (TE) and absolute errors (AE) of different RECPs in their reproducting from the 2 electron valence CI calculations of low-lying states of Hg for the [4, 4, 3, 2, 1] basis set (in cm⁻¹).

^aThe correlation energies were calculated as a difference between the total energies from the above mentioned CI calculations and the numerical SCF calculations with the frozen core $1s_{1/2}$ - $5d_{5/2}$ spinors for the corresponding terms.

^bThe $6s_{1/2}^1 6p_{1/2}^1$ and $6s_{1/2}^1 6p_{3/2}^1$ configurations are strongly mixed for these terms in the CI calculations.

^cThe RECP from Ref. [6].

^dThe PP from Ref. [7] with the corrected V_{so} by the factors (2l + 1)/2 (M. Dolg, private communication).

TABLE III. All-electron transition energies (TE) and absolute errors (AE) of the 20 electron GRECP in their reproducting from the 12 and 18 electron CI/MBPT2 calculations of low-lying states of Hg for the [4, 4, 3, 2, 1] CI, [11, 11, 11, 9, 8, 7] and [11, 12, 11, 9, 8, 7] MBPT basis sets (in cm⁻¹).

Sym-		All-el.	20 el.	All-el.	20 el.
metry	Leading	point	GRECP	point	GRECP
(J_{parity})	conf.	nucl.	gaus.	nucl.	gaus.
Number of	correlated electrons:	12	12	18	18
		TE	AE	TE	AE
0_g	$6s_{1/2}^2$	0	0	0	0
0_u	$6s^{1'}_{1/2}6p^{1}_{1/2}$	36753	-194	37264	-211
1_u	$6s_{1/2}^1 6p_{1/2}^1$	38526	-178	39049	-192
2_u	$6s^{1^{'}}_{1/2}6p^{1^{'}}_{3/2}$	43149	-167	43751	-172
1_u	$6s_{1/2}^{1^{'}}6p_{3/2}^{1^{'}}$	52919	-30	53325	-24

TABLE IV. All-electron transition energies (TE) and absolute errors (AE) of different RECPs in their reproducting from the 20 electron CI/MBPT2 calculations of low-lying states of Hg for the [4, 4, 3, 2, 1] CI and [12, 12, 11, 9, 8, 7] MBPT basis sets (in cm⁻¹).

							20 el.	20 el.
Sym-	Leading	All	-el.	34 el.	20	el.	RECP	energy-
metry	conf.	finite	point	GRECP	GR	ECP	of Ross	adjusted
(J_{parity})		nucl.	nucl.	num.	num.	gaus.	et al. ^a	$\rm PP^b$
		TE	TE	AE	AE	AE	AE	AE
0_g	$6s_{1/2}^2$	0	0	0	0	0	0	0
0_u	$6s_{1/2}^{1'}6p_{1/2}^{1}$	36900	36963	-137	-135	-130	570	-468
1_u	$6s^1_{1/2}6p^1_{1/2}$	38690	38754	-128	-115	-111	581	-233
2_u	$6s^{1}_{1/2}6p^{1}_{3/2}$	43415	43481	-127	-94	-90	656	795
1_u	$6s^1_{1/2}6p^1_{3/2}$	53004	53062	-22	50	52	545	303

^aThe RECP from Ref. [6].

^bThe PP from Ref. [7] with the corrected V_{so} by the factors (2l + 1)/2 (M. Dolg, private communication).

TABLE V. All-electron transition energies (TE) and absolute errors (AE) of the 34 electron GRECP in their reproducting from the 34 electron CI/MBPT2 calculations of low-lying states of Hg for the [4, 4, 3, 2, 1] CI and [12, 12, 11, 10, 8, 7] MBPT basis sets in comparison with experimental data and results of RCC calculations (in cm⁻¹).

			$\mathrm{RCC}^{\mathrm{b}}$			
Sym-	Leading	Exper.	(all-el.,	All	-el.	34 el.
metry	conf.	$data^{a}$	finite	finite	point	GRECP
(J_{parity})			nucl.)	nucl.	nucl.	num.
		TE	TE	TE	TE	AE
0_g	$6s_{1/2}^2$	0	0	0	0	0
0_u	$6s_{1/2}^{1}6p_{1/2}^{1}$	37645	37453	37569	37634	153
1_u	$6s_{1/2}^16p_{1/2}^1$	39412	39302	39361	39426	169
2_u	$6s_{1/2}^{1}6p_{3/2}^{1}$	44043	44190	44157	44224	225
1_u	$6s_{1/2}^{1'}6p_{3/2}^{1'}$	54069	55453	53553	53612	224

^aThe data from Ref. [10].

^bThe results from Ref. [3] for the [27, 23, 21, 16, 10, 6] basis set (in the notations of the present paper) and 34 correlated electrons.

TABLE VI. All-electron transition energies (TE) from the 34 electron CI/MBPT2 calculations of low-lying states of Hg for different basis sets (in cm^{-1}).

Sym-		All-el.	All-el.	All-el.
metry	Leading	point	point	point
(J_{parity})	conf.	nucl.	nucl.	nucl.
CI basis:		$\left[4,4,3,2,1\right]$	[4, 4, 3, 2]	[4, 4, 3]
MBPT basis:		$\left[12,12,11,9,8\right]$	[12, 12, 11, 9]	[12, 12, 11, 1]
		TE	TE	TE
0_g	$6s_{1/2}^2$	0	0	0
0_u	$6s_{1/2}^{1^{'}}6p_{1/2}^{1}$	37370	37068	32680
1_u	$6s_{1/2}^{1'}6p_{1/2}^{1'}$	39151	38798	34530
2_u	$6s_{1/2}^{1^{'}}6p_{3/2}^{1^{'}}$	43906	43296	38847
1_u	$6s_{1/2}^1 6p_{3/2}^1$	53272	52726	50604