## THE QUANTUM MECHANICAL CURRENT OF THE PAULI EQUATION

Marek Nowakowski Grup de Física Teòrica, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

## Abstract

We argue that the process of constructing the quantum mechanical current of the Pauli equation by copying the line of arguments used in the spin-0 case, i.e. the Schrödinger equation, is ambiguous. We show that a non-relativistic reduction of the relativistic Dirac four-vector current is, however, capable of fully resolving the problem. This analysis reveals that the non-relativistic current of the Pauli equation should include an extra term of the form  $\nabla \times (\psi^{\dagger} \sigma \psi)$ . We present an initial exploration of the potential consequences of this new 'spinterm' by solving the Pauli equation for crossed magnetic and electric fields and calculating the corresponding current.

Most of the applications of Quantum Mechanics (QM) can be found in the realm of the three phases of matter: gases, liquids and solids where a non-relativistic quantum mechanical description is fully adequate. This shows the importance of nonrelativistic QM, even though, in principle, a strictly correct treatment should implement relativity as well. The basic constituents of matter are atoms whose building blocks in turn are nuclei and electrons. These last control the properties of matter such as chemical bonds and conductivity. Since electrons are spin-1/2 fermions, the non-relativistic wave equation describing them, namely the Pauli equation, has a somewhat distinguished position in our understanding of the matter surrounding us. But whereas in most books on QM a considerable effort is spent on the interpretation of the Schrödinger equation (i.e. the spin-0 wave equation) in terms of the probability density  $\rho$  and the current  $\mathbf{i}$ , a corresponding discussion of the spin-1/2 case (the Pauli equation) is rarely to be found [1]. This might have to do with the tacit assumption that the construction of the spin-1/2 current goes along the same line of arguments as in the spin-0 case. Hence one might conclude that, apart from a trivial replacement of 'complex-conjugate' by 'hermitian-conjugate', there is conceptually nothing new in the current of the Pauli equation. This, as will be shown below, is misleading. Indeed, in constructing the current for spin-1/2 case one can copy all the steps known from the Schrödinger case to obtain an expression for  $\mathbf{j}$ . However, due to the presence of the spin this construction is ambiguous, i.e. there exist terms which can be added to j and which do not emerge from the above mentioned construction. They can be added since they do not spoil the continuity equation, they are of first order in the derivative and second order of the wave function which means that they are of 'velocity-type' as are the rest of the conventional terms known from the spin-0 case. This ambiguity cannot be resolved by means of non-relativistic QM alone. Or, to put it differently, this ambiguity only appears from the point of view of non-relativistic QM. However, the current must be fixed uniquely as it is an observable.

It is clear that something new is required to solve the problem in a satisfactory way. As it is often the case in physics a new symmetry imposed on the system restricts the number of possible terms and can therefore resolve an otherwise persistent ambiguity. The symmetry we have in mind here, is the Lorentz symmetry. A relativistic wave equation for spin-1/2 particles is the Dirac equation which has a positive definite probability density and a continuity equation [2]. The non-relativistic reduction of the Dirac equation and of the corresponding current will then answer the problem unambiguously. From a pedagogical point of view it seems even desirable to postpone the discussion of the Pauli current until relativistic QM is introduced. Unfortunately, the main stream of interest diverges here and from relativistic QM one usually proceeds to relativistic Quantum Field Theory. Hence it seems that there is a problem which most of the books on non-relativistic QM do not mention. Considering the importance of the current in interpreting QM, it seems that it is worthwhile to fill this gap.

To make the problem concrete, let us start with the Pauli equation for an

electron in the presence of a electromagnetic field  $A_{\mu}$ 

$$i\frac{\partial\psi}{\partial t} = \left[\frac{1}{2m}(-i\boldsymbol{\nabla} - e\boldsymbol{A})^2 - \frac{e}{2m}\boldsymbol{\sigma}\cdot\boldsymbol{B} + eA_0\right]\psi = H_{Pauli}\psi \tag{1}$$

where  $\psi$  is a two component spinor. We have set  $\hbar = c = 1$ . Although mostly we will be concerned with electromagnetic interactions, the point we are making (namely the correct form of the current) is in fact independent of the detailed form of the interaction. We will comment on this later in the text. The probability density

$$\rho = \psi^{\dagger}\psi \tag{2}$$

only has a consistent, well-defined interpretation if it satisfies the continuity equation

$$\frac{\partial \rho}{\partial t} + \boldsymbol{\nabla} \cdot \boldsymbol{j} = 0 \tag{3}$$

Extending the standard prescription for construction of  $\boldsymbol{j}$  from the Schrödinger case (i.e. we use eq.(1) and its hermitian conjugate in computing  $\partial \rho / \partial t$ ) one finds a current which we denote here by  $\boldsymbol{j}'$ 

$$\boldsymbol{j}' = -\frac{i}{2m} \left( \psi^{\dagger} \boldsymbol{\nabla} \psi - (\boldsymbol{\nabla} \psi^{\dagger}) \psi \right) - \frac{e}{m} \boldsymbol{A} \psi^{\dagger} \psi$$
$$= \frac{1}{m} \Im m(\psi^{\dagger} \boldsymbol{\nabla} \psi) - \frac{e}{m} \boldsymbol{A} \rho$$
(4)

This expression is gauge invariant (thanks to the  $A\rho$  term) and could be, in principle, a good candidate for the complete quantum mechanical current of spin-1/2 fermions if we could make it plausible that (4) is in some sense unique. This is, however, not the case. We can trivially add to j' a gauge invariant term proportional to

$$\boldsymbol{\nabla} \times (\psi^{\dagger} \boldsymbol{\sigma} \psi) \tag{5}$$

without changing the continuity equation (3). Note that in the Schrödinger case it is not possible to construct a 'curl-term' which is first order in the derivative and second order in the wave functions.

Hence there is a priori no way to decide (not only for the electromagnetic interaction), from the point of view of non-relativistic QM, whether a term like in eq.(5) should be added to  $\mathbf{j}'$  or not (and if yes what is the proportionality factor). Since the current is a physical observable, this apparent ambiguity must have a unique resolution. Indeed, as will be evident below, there is no such ambiguity in the full physical theory as we can fix the current uniquely by using relativistic arguments. Note also that once this question is settled, the electric current has to be  $\mathbf{J} = e\mathbf{j}$ .

One could of course argue that only 'orbital-terms' like  $\psi^{\dagger} \nabla \psi$  should enter in **j** and hence also in **J**. This has a classical flavour and cannot be regarded as a compelling argument. The correct approach should use a non-relativistic reduction of both, the relativistic wave equation *and its current* and we will see below that the above naive argument does not hold.

Before performing the non-relativistic reduction for the Dirac equation let us emphasize here two points. Relativistic QM including external fields is well-defined below the particle anti-particle production threshold which implies that the external fields should not be too strong. This is to stress the correctness of the relativistic external field problem. While historically one of the first checks of any relativistic theory has been to test that it yielded the standard non-relativistic limit, by now the relativistic theory, and in particular here the Dirac equation, is well-established. Thus when there is a non-relativistic ambiguity such as we have seen above, we may safely use the relativistic Dirac theory to find the correct non-relativistic limit.

The Dirac equation in the Dirac representation of the  $\gamma_{\mu}$ -matrices reads [3]

$$i\frac{\partial}{\partial t}\left(\begin{array}{c}\psi\\\chi\end{array}\right) = \boldsymbol{\sigma}\cdot\boldsymbol{\pi}\left(\begin{array}{c}\chi\\\psi\end{array}\right) + eA_0\left(\begin{array}{c}\psi\\\chi\end{array}\right) - 2m\left(\begin{array}{c}0\\\chi\end{array}\right)$$
(6)

where  $\boldsymbol{\pi} = -i\boldsymbol{\nabla} - e\boldsymbol{A}$ , and  $\psi$  and  $\chi$  are both two component spinors. The non-relativistic reduction starts by assuming the kinetic energy and field strength to be small compared to the mass m. Then one of the equations in (6) is approximately

$$\chi \simeq \frac{\boldsymbol{\sigma} \cdot \boldsymbol{\pi}}{2m} \psi \tag{7}$$

Inserting this in (6) we obtain the Pauli equation (1) for the spinor  $\psi$ . A similar reduction of the probability density yields eq.(2) up to terms of order  $v^2$  which are of the form  $(1/4m^2)(\boldsymbol{\sigma}\cdot\boldsymbol{\pi}\psi)^{\dagger}(\boldsymbol{\sigma}\cdot\boldsymbol{\pi}\psi)$ . Of course we should also apply the same approximation to the spatial components of the Dirac current

$$\boldsymbol{j}_{Dirac} = \Psi^{\dagger} \boldsymbol{\alpha} \Psi = -\left(\psi^{\dagger} \boldsymbol{\sigma} \chi + \chi^{\dagger} \boldsymbol{\sigma} \psi\right)$$
(8)

where  $\boldsymbol{\alpha} = \gamma_0 \boldsymbol{\gamma}$ . Inserting in (8) the non-relativistic approximation (7) and using  $\sigma_i \sigma_j = i \epsilon_{ijk} \sigma_k + \delta_{ij}$  we find the non-relativistic version of the current

$$\boldsymbol{j} = \boldsymbol{j}' + \frac{1}{2m} \boldsymbol{\nabla} \times (\psi^{\dagger} \boldsymbol{\sigma} \psi) + \mathcal{O}(v^2/c^2)$$
(9)

where j' has been already defined in (4). Equation (9) is the correct non-relativistic spin-1/2 current of the Pauli equation. The question about the ambiguity of the Pauli current posed at the beginning has been completely answered.

We see that the correct electric current to be used, say, in addressing questions about conductivity indeed contains a 'spin-term' of the form (5) as well as the usual 'orbital-terms'  $\mathbf{j}'$ . Some comments are in order here. First note that whereas  $\mathbf{j}'$  depends explicitly on the interaction (potential) used in the Pauli equation (1), the 'spin-term'  $(1/2m)\nabla \times (\psi^{\dagger}\boldsymbol{\sigma}\psi)$  does not. This is clear from eq. (4) where the vector potential enters explicitly. Had we used an interaction other than the electromagnetic one,  $\mathbf{j}'$  could then still be constructed as in the Schrödinger case, but the 'spin-term' would then follow from a corresponding non-relativistic reduction of the Dirac equation (coupled to this interaction) and its current. In other words this term will always be present, regardless of the interaction, and indeed even in the interaction-free case (it is even an easier excercise to perform the non-relativistic reduction for free electrons). From the above it follows that while j' is closely related to the detailed form of the Pauli equation, the 'spin-term' is not. That is why from the point of view of non-relativistic quantum mechanics there seems to be an ambiguity. It seems that, in general, the non-relativistic result contains less information when viewed independently from its relativistic 'parent'. This is not surprising as imposing a symmetry, here the Lorentz symmetry, limits the number of choices and can therefore 'seal the fate' of a possible structure of a term. In the case of electromagnetic interactions we can relate the 'spin-term' to the  $(e/2m)\boldsymbol{\sigma}\cdot\boldsymbol{B}$  term in the Hamiltonian of the Pauli equation, provided we are allowed to use some arguments from field theory where a part of the interaction Hamiltonian is given by  $e \int d^3x \mathbf{i} \cdot \mathbf{A}$ . Inserting here the 'spin-term' of the current (9) we recover, after partial integration, the  $(e/2m)\boldsymbol{\sigma}\cdot\boldsymbol{B}$  term of the Hamiltonian (we assume here  $\boldsymbol{B}$  to be constant and the wave-packets localized). Note that this argument, in the case of electromagnetic interactions, would link the Pauli equation with the full current (9) and produce the correct result seemingly without the non-relativistic reduction. However, there are several drawbacks to it. For instance, we should not mix (quantum) field theory with 'point-particle' (quantum) physics. Indeed, in the latter the current is a functional of the wave function which in turn is the solution of the Pauli equation. A 'confirmation' of the Hamiltonian via the field theory method can thus be understood as a useful heuristic argument. As mentioned above, the presence of the 'spin-term' in (9) is also independent of the form of interaction we use. But the argument given above is limited to the electromagnetic interaction. It can, however, always be used heuristically to check the correctness of (9).

To show the relevance of the 'new' term in eq. (9) explicitly, let us solve the Pauli equation (1) for an electron in an uniform magnetic field  $\boldsymbol{B}$  (pointing in the positive z-direction) and an uniform electric field  $\boldsymbol{E}$  (in the negative x-direction, say). The electromagnetic field configuration is then like that of the Hall effect [4]. We are considering here the case of one electron in the presence of electromagnetic fields, but in vacuum otherwise. This is then unlike the Hall effect where the electrons are moving in solids (to avoid confusion, we are not suggesting here a new way to solve the Quantum Hall effect). In the so-called Landau gauge  $A_0 = \mathcal{E}x$ ,  $A_2 =$  $\mathcal{B}x$ ,  $A_1 = A_3 = 0$ ,  $\mathcal{E} = |\boldsymbol{E}|$ ,  $\mathcal{B} = |\boldsymbol{B}|$  the Hamiltonian  $H_{Pauli}$  from (1) commutes with  $-i\partial/\partial y$ ,  $-i\partial/\partial z$  and with  $\sigma_3$ . We make therefore the ansatz for an unnormalized wave function

$$\psi_{\xi n} = e^{-iE_{\xi n}} e^{ip_y y} e^{ip_z z} \Phi_{\xi n}$$
  
$$\Phi_{\xi=1,n} = \begin{pmatrix} \varphi_n \\ 0 \end{pmatrix}, \quad \Phi_{\xi=-1,n} = \begin{pmatrix} 0 \\ \varphi_n \end{pmatrix}$$
 (10)

where the quantum number  $\xi$  denotes different polarizations. We leave the wave function unlocalized in the y and z direction. The eigenvalue problem can be then brought into a form familiar from the one-dimensional harmonic oscillator

$$\left[E_{\xi n} - \frac{p_y^2 + p_z^2}{2m} + \frac{1}{2}\omega\xi + \frac{1}{2}m\omega^2\beta^2\right]\varphi_n$$
$$= (n + \frac{1}{2})\omega\varphi_n = \left[-\frac{1}{2m}\frac{\partial^2}{\partial x'^2} + \frac{1}{2}m\omega^2\eta^2\right]\varphi_n$$
(11)

where

$$\omega = \frac{e\mathcal{B}}{m}, \quad \beta = -\frac{e\mathcal{E} - \omega p_y}{m\omega^2}$$
$$x' = x - \beta, \quad \eta = \sqrt{m\omega}x' \tag{12}$$

The energy eigenvalues can be easily shown to be

$$E_{\xi n} = (n + \frac{1}{2})\omega - \frac{1}{2}\omega\xi + \frac{p_y^2 + p_z^2}{2m} - \frac{1}{2}m\omega^2\beta^2$$
(13)

while the eigenfunctions are given in terms of Hermite polynomials in the variable  $\eta$ 

$$\varphi_n = N_n e^{-\frac{1}{2}\eta^2} H_n(\eta)$$

$$N_n = \sqrt{\sqrt{\frac{m\omega}{\pi} \frac{1}{2^n n!}}}$$
(14)

We can now compute the components of the electric current J = ej with j given in (9). For definiteness we do it for  $\xi = 1$ .

$$J_x = \frac{e}{m} \frac{\partial}{\partial y} \rho_n = 0$$
  

$$J_z = e \frac{p_z}{m} \rho_n$$
  

$$J_y = e \rho_n \frac{\mathcal{E}}{\mathcal{B}} - \frac{e}{m} 2n e^{-\eta^2} H_n(\eta) H_{n-1}(\eta)$$
(15)

where  $\rho_n = \psi_{\xi=1,n}^{\dagger} \psi_{\xi=1,n}$ . For n = 0, i.e. in the ground state, the second term in  $J_y$  vanishes. The first term, viz.  $e\rho_n \mathcal{E}/\mathcal{B}$ , corresponds in this form to the classical result (see e.g. [4]). In the ground state therefore, the classical and quantum mechanical results coincide. For higher excited states there is, however, a new contribution to  $J_y$  (proportional to 2ne/m) which is of purely quantum mechanical origin and which can be traced back to the 'spin-term' in eq.(5). As already stated, we have not tried to solve here the Quantum Hall effect [5]. Our main motivation was to point out the relevance of the 'spin-term' to the (electric) current. It would clearly be interesting to study the effects of the extra term in the current in various physical applications.

We feel it is reasonable to speculate that this term might play a role in problems concerning conductivity in solids. Finally, we recall that the proof of the need for such a term in the non-relativistic current was decided essentially by relativistic QM. This is one of the few places where relativistic QM can resolve a problem of the nonrelativistic theory. We think that text books on non-relativistic QM should include at least a note on the different (as compared to the Schrödinger case) nature of the quantum mechanical Pauli current in order not to give the impression that in constructing the Pauli current it suffices to copy the steps from the Schrödinger case. The quantum mechanical current is not only important for the correct interpretation of QM, but also in calculating conductivity etc. in solid matter. In teaching this subject, one can point out the power of symmetry arguments which limits our choice of possible terms. Since the appearance of the ambiguity (from the non-relativistic point of view) in the Pauli current is physically not acceptable (the current is an observable), this hints towards the need of a more general theory which includes relativity. In spite of the fact that in practical calculations concerning properties of solids relativity does not play a big part, it plays a role, as shown above, on a more fundamental level. This is then also a lesson on the unity of physics.

**Acknowledgments**. I would like to thank R. Godbole and M. Lavelle for many valuable discussions and suggestions. This work has been supported by the Spanish Ministerio de Educacion y Ciencia.

## References

- [1] A disclaimer is in order at this point. The author of the present paper does not claim that there do not exist any books with a discussion of the spin-1/2 current. It is then possible that a discussion can be found where the results are similar or opposite to these presented here. If so, the exercise of this note here is to point out the problem and its solution.
- [2] The author noticed the problem of the Pauli current while working on a different but related problem of constructing a current for neutral spin-0  $K^0$  and  $\bar{K^0}$  mesons. There one of the problems is to have a relativistic well-defined current.See, B. Ancochea, A. Bramon, R. Munoz-Tapia, M. Nowakowski, "Spacedependent probabilities for  $K^0 - \bar{K^0}$  oscillations", Phys. Lett. **B389**, 149-156 (1996)

- [3] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics*, (McGraw-Hill, 1964), pp. 1-10
- [4] For an overview see in A. P. Balachandran, E. Ercolesi, G. Morandi and A. M. Srivastava, *Hubbard Model and Anyon Superconductivity*, (World Scientific, 1990), pp. 121-135
- [5] For a general reference see *The Quantum Hall Effect*, eds. R. E. Prange and S. M. Girvin, (Springer-Verlag, 1990) 2nd edition; also G. Morandi, *Quantum Hall Effect*, Bibliopolis 1988