

An Extended Liouville Equation for Variable Particle Number Systems

Michael H. Peters^{a)}

Department of Chemical Engineering, Florida State University

and Florida A&M University, Tallahassee, FL 32310

(Submitted September 1998)

It is well-known that the Liouville equation of statistical mechanics is restricted to systems where the total number of particles (N) is fixed. In this paper, we show how the Liouville equation can be extended to systems where the number of particles can vary, such as in open systems or in systems where particles can be annihilated or created. A general conservation equation for an arbitrary dynamical variable is derived from the extended Liouville equation following Irving and Kirkwood's² technique. From the general conservation equation, the particle number conservation equation is obtained that includes general terms for the annihilation or creation of particles. It is also shown that the grand canonical ensemble distribution function is a particular stationary solution of the extended Liouville equation, as required. In general, the extended Liouville equation can be used to study nonequilibrium systems where the total number of particles can vary.

^{a)} e-mail: peters@eng.fsu.edu

I. Introduction

Modern molecular theories of equilibrium thermodynamics and nonequilibrium thermodynamics are based almost entirely on the Liouville equation which describes the conservation of an ensemble of phase points as they move through phase space according to a specified Hamiltonian¹. Unfortunately, the Liouville equation is based on a fixed number of particles and cannot describe systems where the total number of particles varies. For example, the Liouville equation cannot describe the statistical mechanical, grand canonical ensemble used in the development of thermodynamic properties of open, equilibrium systems. In this letter, we extend the Liouville equation to systems where the number of particles varies. To our knowledge this has not been done before and has restricted the theoretical development of nonequilibrium thermodynamics. These systems include open systems, where the particle number can vary because of open boundaries, or closed or open systems where particles can be annihilated or created either classically by chemical reactions or due to quantum mechanical effects.

II. The Extended Liouville Equation

The Liouville equation describes the behavior of a collection or ensemble of phase points as they move through a multidimensional space, or phase-space. Each phase point represents the position (or generalized coordinates q_i) and momentum (or conjugate momentum p_i) of all N molecules or particles in the system. The phase points tend to be concentrated in regions of phase space where it is most likely to find the N particles with a certain momentum and position. Thus, the Liouville density function, ρ_L , can be interpreted (aside from a normalization constant) as a probability density function, i.e., $\rho_L d\mathbf{q}^N d\mathbf{p}^N$ is proportional to the probability of finding a phase point in a multidimensional region between $(\mathbf{q}^N, \mathbf{p}^N)$ and $(\mathbf{q}^N + d\mathbf{q}^N, \mathbf{p}^N + d\mathbf{p}^N)$ at any time, t .

Here and throughout we use the short-hand notation $\mathbf{q}^N = q_1, \dots, q_{3N}$ and $\mathbf{p}^N = p_1, \dots, p_{3N}$. It can be readily shown that the Liouville density function obeys the conservation equation¹

$$\frac{\partial \rho_L(\mathbf{p}^N, \mathbf{q}^N, t)}{\partial t} + \sum_{i=1}^{3N} \left[\frac{\partial \rho_L}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \rho_L}{\partial p_i} \frac{\partial H}{\partial q_i} \right] = 0 \quad (1)$$

where $H(\mathbf{p}^N, \mathbf{q}^N, t)$ is the Hamiltonian or total energy of the N -particle system.

It is to be noted that the Liouville equation describes the phase space behavior of the density function $\rho_L(\mathbf{p}^N, \mathbf{q}^N, t)$ under the conditions where the number of particles N is a constant. To extend the Liouville equation to a system where the number of particles or dimensionality can vary, consider the extended space $(\mathbf{p}^N, \mathbf{q}^N, N, t)$ shown in Fig.1. The total members of the ensemble in extended space is a constant denoted by M , and the density of phase points is denoted by $\rho_N(\mathbf{p}^N, \mathbf{q}^N, N, t)$. Thus, we have that

$$\sum_N \int \int \rho_N(\mathbf{p}^N, \mathbf{q}^N, N, t) d\mathbf{p}^N d\mathbf{q}^N = M \quad (2)$$

As shown in Fig.1, the extended space can be divided into “canonical” hyperplanes with the number of phase points at time t in each hyperplane denoted by M_{N_1}, M_{N_2} , etc., where

$$M = \sum_{N_i} M_{N_i} \quad (3)$$

The phase points in each hyperplane have the same number of particles or dimensionality and, hence, we call these “canonical” hyperplanes. As the system evolves in time, the phase points can move within the plane, if the dimensionality (N) does not change, or “jump” between

planes, if the dimensionality (N) of the phase point changes. In any event, the total number of phase points M remains constant with time.

Now, it is not possible to straightforwardly extend the Liouville equation by considering differential changes in \mathbf{q}^N , \mathbf{p}^N and N (the extended space), since as N changes so does the dimensionality of the phase space coordinates (\mathbf{q}^N , \mathbf{p}^N). However, if we define a new function

$$\bar{\rho}(N, t) = \int \int \rho_N(\mathbf{p}^N, \mathbf{q}^N, N, t) d\mathbf{p}^N d\mathbf{q}^N = M_N \quad (4)$$

then, in order to conserve the total number of phase points M , it can be readily shown (Appendix A) that $\bar{\rho}$ must obey the one-dimensional conservation equation

$$\frac{\partial \bar{\rho}}{\partial t} + \frac{\partial}{\partial N} [\bar{\rho} v(N, t)] = 0 \quad (5)$$

where we have introduced a “velocity” function, $v(N, t)$,

$$v(N, t) \equiv v^+(N, t) - v^-(N, t) \quad (6)$$

which physically represents the net fractional number of phase points leaving the N^{th} hyperplane per unit time (Appendix A); v^+ represents the fraction of phase points leaving the N^{th} hyperplane to the $(N + 1)^{th}$ hyperplane, and v^- are the fraction per unit time leaving to the $(N - 1)^{th}$ hyperplane. Now, within each hyperplane we introduce the Liouville density $\rho_L(\mathbf{p}^N, \mathbf{q}^N, t; N)$, defined as a conditional density function

$$\begin{aligned}\rho_L(\mathbf{p}^N, \mathbf{q}^N, t; N) &\equiv \frac{\rho_N(\mathbf{p}^N, \mathbf{q}^N, N, t)}{\int \rho_N d\mathbf{p}^N d\mathbf{q}^N} \\ &= \frac{\rho_N}{\bar{\rho}}\end{aligned}\tag{7}$$

The phase points that move within the hyperplane over a differential time, dt , obey the usual, Liouville equation, Eq.(1),

$$\frac{\partial \rho_L}{\partial t} + \sum_{i=1}^{3N} \left(\frac{\partial \rho_L}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \rho_L}{\partial p_i} \frac{\partial H}{\partial q_i} \right) = 0\tag{8}$$

Thus, the space-time behavior of the extended Liouville density function $\rho_N(\mathbf{p}^N, \mathbf{q}^N, N, t)$ is uniquely determined through Eqs.(5), (7) and (8). Physically, we can envision that phase points move in each hyperplane over a differential time, dt , according to Eq.(8). At the end of the differential time, some phase points “jump” into new canonical hyperplanes according to Eq.(5). Thus, the number of phase points in each canonical hyperplane with a fixed dimensionality continually change with time. Although both the dimensionality change and time change are discrete, the continuous representation given by Eq.(5) should approximately hold for N large and $dt \rightarrow 0$.

Multiplying Eq.(5) by ρ_L , Eq.(8) by $\bar{\rho}$, and combining using Eq.(7) we can write *an extended Liouville equation* as

$$\frac{\partial \rho_N(\mathbf{q}^N, \mathbf{p}^N, N, t)}{\partial t} + \bar{\rho}(N, t) \left\{ \sum_i \frac{\partial \rho_L}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \rho_L}{\partial p_i} \frac{\partial H}{\partial q_i} \right\}$$

$$+\rho_L(\mathbf{q}^N, \mathbf{p}^N, t; N) \frac{\partial}{\partial N} [\bar{\rho}v(N, t)] = 0 \quad (9)$$

or,

$$\begin{aligned} \frac{\partial \ln \rho_N}{\partial t} + \sum_i \left(\frac{\partial \ln \rho_L}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \ln \rho_L}{\partial p_i} \frac{\partial H}{\partial q_i} \right) \\ +v(N, t) \frac{\partial \ln \bar{\rho}}{\partial N} + \frac{\partial v}{\partial N} = 0 \end{aligned} \quad (10)$$

In the next section, we derive a general conservation (transport) equation for an arbitrary dynamic variable where the total number of particles can vary. From this we obtain a particle number conservation equation which includes a general term accounting for the annihilation or creation of particles.

III. General Conservation Equation for Variable Particle Number Systems

In Cartesian coordinates, the extended Liouville equation can be written

$$\begin{aligned} \frac{\partial \rho_N(\mathbf{r}^N, \mathbf{p}^N, N, t)}{\partial t} + \bar{\rho}(N, t) \left\{ \sum_i \left[\frac{\partial}{\partial \mathbf{r}_i} \cdot \left(\frac{\mathbf{p}_i}{m} \rho_L \right) + \frac{\partial}{\partial \mathbf{p}_i} \cdot (\mathbf{F}_i \rho_L) \right] \right\} \\ +\rho_L(\mathbf{r}^N, \mathbf{p}^N, t; N) \frac{\partial}{\partial N} [\bar{\rho}v(N, t)] = 0 \end{aligned} \quad (11)$$

where we will now normalize the density function $\bar{\rho}$ by the total ensemble number M (fixed), i.e.,

$$\sum_N \bar{\rho} = 1 \quad (12)$$

$$\int \int \rho_L d\mathbf{r}^N d\mathbf{p}^N = 1 \quad (13)$$

$$\sum_N \int \int \rho_N d\mathbf{r}^N d\mathbf{p}^N = 1 \quad (14)$$

Extending Irving and Kirkwood's formalism², the "conservation equations" for variable particle number systems can be obtained by first considering an arbitrary quantity $\alpha(\mathbf{r}^N, \mathbf{p}^N, N)$ that does not depend explicitly on time. The average or expectation value of α is defined as

$$\langle \alpha \rangle \equiv \sum_N \int \int \alpha(\mathbf{r}^N, \mathbf{p}^N, N) \rho_N d\mathbf{r}^N d\mathbf{p}^N \quad (15)$$

Now, multiplying the extended Liouville equation by α and integrating over all $(\mathbf{r}^N, \mathbf{p}^N)$ and summing over all N yields

$$\begin{aligned} \frac{\partial \langle \alpha \rangle}{\partial t} - \sum_N \bar{\rho} \int \int \sum_i \rho_L \frac{\mathbf{p}_i}{m} \cdot \frac{\partial \alpha}{\partial \mathbf{r}_i} d\mathbf{r}^N d\mathbf{p}^N \\ - \sum_N \bar{\rho} \int \int \sum_i \rho_L \mathbf{F}_i \cdot \frac{\partial \alpha}{\partial \mathbf{p}_i} d\mathbf{r}^N d\mathbf{p}^N \\ + \sum_N \frac{\partial}{\partial N} (\bar{\rho} v) \int \int \alpha \rho_L d\mathbf{r}^N d\mathbf{p}^N \end{aligned} \quad (16)$$

where we have used the conditions

$$\left. \begin{array}{l} \rho_L \rightarrow 0 \\ \mathbf{p}_i \rho_L \rightarrow 0 \\ \alpha \mathbf{p}_i \rho_L \rightarrow 0 \end{array} \right\} \text{as } \mathbf{r}_i, \mathbf{p}_i \rightarrow \infty \quad (17)$$

Equation (16) is the general conservation equation for α .

Particle Number Conservation

Again, extending Irving and Kirkwood's formalism² for variable particle number systems, the equation for particle number conservation (ordinary number density) can be obtained from Eq.(16) by setting

$$\alpha(\mathbf{r}_N, \mathbf{p}_N, N) = \sum_{k=1}^N \delta(\mathbf{r}_k - \mathbf{r}) \quad (18)$$

Thus,

$$\begin{aligned} \langle \alpha \rangle &= n(\mathbf{r}, t) \equiv \sum_N \bar{\rho} \int \int \left[\sum_{k=1}^N \delta(\mathbf{r}_k - \mathbf{r}) \right] \rho_L d\mathbf{r}^N d\mathbf{p}^N \\ &= \sum_N \bar{\rho} N \int \int \rho_1(\mathbf{r}, \mathbf{p}, t; N) d\mathbf{p} \\ &= \sum_N \bar{\rho} N \psi_1(\mathbf{r}, t; N) \end{aligned} \quad (19)$$

where $\psi_1(\mathbf{r}, t; N)d\mathbf{r}$ is the conditional probability of finding a particle between \mathbf{r} and $\mathbf{r} + d\mathbf{r}$, given a system of N total particles at time t , i.e.,

$$\psi_1(\mathbf{r}, t; N) \equiv \int \rho_1(\mathbf{r}, \mathbf{p}, t; N)d\mathbf{p} \quad (20)$$

Note that for a system at equilibrium

$$\psi_1 = \frac{1}{V} \quad , \quad \text{equilibrium} \quad (21)$$

where V is the volume of the system; thus

$$n(\mathbf{r}, t) = \frac{1}{V} \sum_N \bar{\rho} N = \frac{\bar{N}}{V} \quad , \quad \text{equilibrium} \quad (22)$$

where \bar{N} is the equilibrium average number of particles in the volume V .

Analyzing the remaining terms in Eq.(16)

$$\begin{aligned} & - \sum_N \bar{\rho} \int \int \left[\sum_i \rho_L \frac{\mathbf{p}_i}{m} \cdot \frac{\partial \alpha}{\partial \mathbf{r}_i} \right] d\mathbf{r}^N d\mathbf{p}^N \\ & = \frac{\partial}{\partial \mathbf{r}} \cdot \left[\sum_N \bar{\rho} N \int \rho_1 \frac{\mathbf{p}}{m} d\mathbf{p} \right] \\ & = \frac{\partial}{\partial \mathbf{r}} \cdot [n(\mathbf{r}, t) \mathbf{v}_0(\mathbf{r}, t)] \end{aligned} \quad (23)$$

where we have defined the number average velocity as

$$\mathbf{v}_0(\mathbf{r}, t) \equiv \frac{\sum_N \bar{\rho} N \int \rho_1 \frac{\mathbf{p}}{m} d\mathbf{p}}{\sum_N \bar{\rho} N \psi_1} \quad (24)$$

and, for the final term,

$$\begin{aligned} & \sum_N \left[\frac{\partial}{\partial N} (\bar{\rho} v) \right] \int \int \alpha \rho_L d\mathbf{r}^N d\mathbf{p}^N \\ &= \sum_N \left[\frac{\partial}{\partial N} (\bar{\rho} v) \right] N \psi_1(\mathbf{r}, t; N) \end{aligned} \quad (25)$$

Summarizing, the particle number conservation equation becomes

$$\frac{\partial n(\mathbf{r}, t)}{\partial t} + \frac{\partial}{\partial \mathbf{r}} \cdot (n \mathbf{v}_0) + \sum_N \left[\frac{\partial}{\partial N} (\bar{\rho} v) \right] N \psi_1(\mathbf{r}, t; N) = 0 \quad (26)$$

The third term on the left-hand side accounts for the creation or annihilation of particles in the system as considered in more detail in the example below.

Example. First-Order Particle Annihilation

As a simple example of the application of Eq.(26) consider the problem of particle annihilation where the fractional number of particles leaving the N^{th} hyperplane to the $(N - 1)^{\text{th}}$ hyperplane is a constant k . Thus, Eq.(5) becomes

$$\frac{\partial \bar{\rho}}{\partial t} - k \frac{\partial \bar{\rho}}{\partial N} = 0 \quad (27)$$

or, in “finite N ” form (Appendix A)

$$\frac{\partial \bar{\rho}(N, t)}{\partial t} - k[\bar{\rho}(N + 1, t) - \bar{\rho}(N, t)] = 0 \quad (28)$$

Consider that initially all phase points are in the N_0 hyperplane, i.e.,

$$\bar{\rho}(N_0, 0) = 1 \quad (29)$$

and

$$\bar{\rho}(N, 0) = 0 \quad , \quad N \in [0, N_0 - 1] \quad (30)$$

Around the N_0 hyperplane, the population balance equation is

$$\frac{\partial \bar{\rho}(N_0, t)}{\partial t} + k\bar{\rho}(N_0, t) = 0 \quad (31)$$

Thus, using the initial state Eq.(29), we obtain

$$\bar{\rho}(N_0, t) = e^{-kt} \quad (32)$$

Now, performing a balance around the $N_0 - 1$ hyperplane using Eq.(32) leads to

$$\frac{\partial \bar{\rho}(N_0 - 1, t)}{\partial t} + k\bar{\rho}(N_0 - 1, t) = ke^{-kt} \quad (33)$$

and, using the initial condition Eq.(30), we obtain

$$\bar{\rho}(N_0 - 1, t) = kte^{-kt} \quad (34)$$

Continuing this process, for the N^{th} hyperplane we must have

$$\bar{\rho}(N, t) = \frac{(kt)^m}{m!} e^{-kt} \quad (35)$$

where $m = N_0 - N$, $m \in [0, N_0 - 1]$. We note that for $N_0 \rightarrow \infty$,

$$\sum \bar{\rho}(N, t) = e^{-kt} \sum_{m=0}^{\infty} \frac{(kt)^m}{m!} = 1 \quad (36)$$

as required. Note that for small or finite values of N_0 , we must consider a population balance around the $N = 0$ hyperplane (total particle annihilation plane) in order to satisfy the normalization condition.

Now consider the annihilation/creation term in the particle number conservation equation, Eq.(26). We have that

$$\frac{\partial}{\partial N}(\bar{\rho}v) = -\frac{\partial \bar{\rho}}{\partial t} = k \left[1 - \frac{(N_0 - N)}{kt} \right] \bar{\rho} \quad (37)$$

Substituting into the third term on the l.h.s of Eq.(26) leads to the first-order decay kinetic expression, under the conditions $kt \gg (N_0 - N)$, as

$$\frac{\partial n(\mathbf{r}, t)}{\partial t} + \frac{\partial}{\partial \mathbf{r}} \cdot (n\mathbf{v}_0) + \sum_N k\bar{\rho}N\psi_1 = 0 \quad , \quad kt \gg (N_0 - N) \quad (38)$$

or

$$\frac{\partial n(\mathbf{r}, t)}{\partial t} + \frac{\partial}{\partial \mathbf{r}} \cdot (n\mathbf{v}_0) + kn = 0 \quad , \quad kt \gg (N_0 - N) \quad (39)$$

Note that for small times compared to $(N_0 - N)/k$ the decay kinetics are not first-order, even for a constant rate of phase point hopping, k . This example helps illustrate the generality of the extended Liouville equation. Next, we address the question as to whether the extended Liouville equation leads to an equilibrium behavior consistent with the grand canonical ensemble.

IV. Equilibrium Behavior. Grand Canonical Ensemble

Under equilibrium conditions, $\partial\bar{\rho}/\partial t = 0$, the general stationary solution to Eq.(5) can be written as

$$\bar{\rho}(N, t) = C_1 f(N) \quad (40)$$

$$v(N, t) = C_2 [f(N)]^{-1} \quad (41)$$

where C_1 and C_2 are constants to be determined and $f(N)$ is an arbitrary function of N . Now,

it is already well-known that a *particular stationary solution* to Eq.(8), under the conditions $\partial\rho_L/\partial t = 0$, can be written in dimensionally correct form as³

$$\rho_L(\mathbf{p}^N, \mathbf{q}^N; N) = \frac{C_3}{h^{3N}N!} \exp \left\{ -\beta H(\mathbf{p}^N, \mathbf{q}^N) \right\} \quad (42)$$

where h is Planck's constant that has the units of action (length \times momentum), $H(\mathbf{p}^N, \mathbf{q}^N)$ is the Hamiltonian or total energy, and C_3 and β are constants to be determined.

The constant β can be determined by requiring that the average kinetic energy on each hyperplane be a uniform constant (canonical ensemble on the hyperplane), i.e., in Cartesian coordinates

$$\frac{(1/2m) \int p_k^2 \rho_L(\mathbf{r}^N, \mathbf{p}^N) d\mathbf{p}^N}{\int \rho_L d\mathbf{p}^N} = \frac{3}{2} kT \quad (43)$$

and

$$H(\mathbf{r}^N, \mathbf{p}^N) = \sum_{i=1}^{3N} \frac{1}{2} \frac{p_i^2}{m} + \Phi(\mathbf{r}^N) \quad (44)$$

where $\Phi(\mathbf{r}^N)$ is the total potential. Thus, we obtain

$$\beta = \frac{1}{kT} \quad (45)$$

Combining Eqs.(40) and (42) gives the extended equilibrium density function as ($C_4 \equiv C_1 C_3$)

$$\rho_N(\mathbf{p}^N, \mathbf{q}^N, N) = \frac{C_4}{h^{3N} N!} \exp \left\{ -\beta H(\mathbf{p}^N, \mathbf{q}^N) \right\} f(N) \quad (46)$$

The constant C_4 can be determined from Eq.(2) as

$$C_4 = \frac{M}{\sum_N \frac{1}{h^{3N} N!} f(N) \int \int \exp \left\{ -\beta H(\mathbf{p}^N, \mathbf{q}^N) \right\} d\mathbf{p}^N d\mathbf{q}^N} \quad (47)$$

Finally, normalizing ρ_N with the total number of phase points M leads to the well-known form for the (classical) probability distribution function for the grand canonical ensemble as⁵,

$$\rho_N(\mathbf{p}^N, \mathbf{q}^N, N) = [E^{Grand} N! h^{3N}]^{-1} f(N) \exp \left\{ -\frac{H(\mathbf{p}^N, \mathbf{q}^N)}{kT} \right\} \quad (48)$$

where the (classical) grand partition function E_N^{Grand} is given by

$$E^{Grand} = \sum_N E_N^{can} f(N) \quad (49)$$

where E_N^{can} is the classical canonical partition function

$$E_N^{can} = [N! h^{3N}]^{-1} \int \int \exp \left\{ -\beta H(\mathbf{p}^N, \mathbf{q}^N) \right\} d\mathbf{p}^N d\mathbf{q}^N \quad (50)$$

Now, it is well-known from most probable distribution methods⁶ that

$$f(N) = e^{\lambda N} \quad (51)$$

for the equilibrium grand canonical ensemble. All that we can state from the extended Liouville equation is that Eq.(51) is a *particular stationary solution* to Eq.(5)⁴. We are presented with the same difficulty as in the equilibrium solution to the ordinary Liouville equation, Eq.(29), i.e., the lack of proof of the uniqueness of Eq.(29) under equilibrium conditions - a well-known problem in statistical mechanics.

Finally, we note that to complete the grand canonical ensemble description, we must still specify λ . This procedure is already well-known and is generally accomplished by comparison to the thermodynamic relation⁶.

$$TdS = dU + pdV - \mu dN \quad (52)$$

where μ is the chemical potential

$$\mu \equiv \left(\frac{\partial U}{\partial N} \right)_{S,V} \quad (53)$$

which leads to

$$\lambda = + \frac{\mu}{kT} \quad (54)$$

and an expression for the entropy, S , in the grand canonical ensemble as

$$S = \frac{\bar{U}}{T} - \frac{\bar{N}\mu}{T} + k \ln E^{Grand} \quad (55)$$

where \bar{U} and \bar{N} are the average internal energy and particle number over the grand canonical ensemble

$$\bar{U} = \sum_N \int \int H \rho_N d\mathbf{p}^N d\mathbf{q}^N \quad (56)$$

$$\bar{N} = \sum_N \int \int N \rho_N d\mathbf{p}^N d\mathbf{q}^N \quad (57)$$

The results and procedures given in Eqs.(52) - (57) are already well established, and we present them only for completeness.

IV. Conclusions

The importance of the analysis given here lies in the evolution equations, Eqs.(5) - (8), or Eqs.(9) or (10), that describe, in general, the nonequilibrium behavior of the extended Liouville density $\rho_N(\mathbf{p}^N, \mathbf{q}^N, N, t)$ for systems where the number of particles is not fixed. A general conservation equation can be derived, following Irving and Kirkwood's paradigm², for systems where the total number of particles can vary. As an example, the general particle number conservation equation was obtained that was shown to include a general term for particle annihilation or creation. It was also demonstrated that the grand canonical ensemble distribution is a particular stationary solution to the extended Liouville equation. As with the equilibrium solution to the ordinary Liouville equation, *uniqueness* of the variable particle number solution under equilibrium conditions remains to be demonstrated. It is to be noted that the general stationary solution to the extended Liouville equation, Eqs.(40) and (41), demonstrates that $v(N)$ is smallest where $\bar{\rho}$ is largest, i.e., near the most probable state. Conversely, phase points move

or change their dimensionality more rapidly from regions where $\bar{\rho}$ is small, i.e., near the least probable state. Further treatment of the extended Liouville equation in nonequilibrium systems is under current investigation.

-
- ¹ Liouville, *Journ. de Math.* **3**, 349 (1838), as cited by R.C. Tolman, *The Principles of Statistical Mechanics*, Oxford University Press, 1938 (p.49).
- ² J.H. Irving and J.G. Kirkwood, *J. Chemical Phys.*, **18**, 817 (1950).
- ³ See J.O. Hirschfelder, C.F. Curtiss and R.B. Bird, *Molecular Theory of Gases and Liquids*, Wiley, NY, 1960 (Eq.2.1 - 20, p.86).
- ⁴ From Eq.(5), we have that $\partial \ln \bar{\rho} / \partial N = \partial \ln v / \partial N = \lambda$ which gives Eq.(51) as a particular solution.
- ⁵ See, e.g., Ref. 3 (Eq.2.1 - 22, p.87).
- ⁶ D.A. McQuarrie, *Statistical Mechanics*, Harper and Row, NY, 1976 (Ch.3, p.54).

Acknowledgements

I want to thank Professor I. Bitsanis for suggesting this problem to me.

Appendix A. Conservation Equation for $\bar{\rho}(N, t)$

Consider a simple population balance around the N^{th} hyperplane (see Fig.1). Since phase points cannot be created or destroyed, we must have

$$\frac{\partial \bar{\rho}(N, t)}{\partial t} + [\bar{\rho}(N, t)v^+(N, t) - \bar{\rho}(N - 1, t)v^+(N - 1, t)]$$

$$-[\bar{\rho}(N+1, t)v^-(N+1, t) - \bar{\rho}(N, t)v^-(N, t)] = 0 \quad (A1)$$

where $v^+(N, t)$ represents the fraction of phase points in the N^{th} hyperplane leaving to the $(N+1)^{\text{th}}$ plane per unit time, $v^-(N, t)$ represents the fraction of phase points in the N^{th} hyperplane leaving to the $(N-1)^{\text{th}}$ plane per unit time, etc. Now, using a finite difference representation of a derivative, we can write

$$\frac{\partial(\bar{\rho}v^+)}{\partial N} \cong \frac{\bar{\rho}(N, t)v^+(N, t) - \bar{\rho}(N-1, t)v^+(N-1, t)}{N - (N-1)} \quad (A2)$$

Thus, Eq.(A1) becomes in the limit of large N

$$\frac{\partial \bar{\rho}}{\partial t} + \frac{\partial(\bar{\rho}v^+)}{\partial N} - \frac{\partial(\bar{\rho}v^-)}{\partial N} = 0 \quad (A3)$$

or,

$$\frac{\partial \bar{\rho}}{\partial t} + \frac{\partial(\bar{\rho}v)}{\partial N} = 0 \quad (A4)$$

where

$$v \equiv v^+ - v^- \quad (A5)$$

This figure "Figure1.GIF" is available in "GIF" format from:

<http://arxiv.org/ps/physics/9809039v2>