

Some FRW models with variable G and Λ

José Antonio Belinchón*

Grupo Interuniversitario de Análisis Dimensional

Dept. Física ETS Arquitectura UPM Juan de Herrera 4 28040 España

January 30, 2018

Abstract

We consider several models with metrics type FRW, with $k = 0$ and with a generic equation of state, $p = \omega\rho$, furthermore, we take into account the conservation principle of the energy-momentum tensor, $div(T_{ij}) = 0$, but with G and Λ variable. We find trivially, a set of solutions through dimensional analysis. We stand out that, $G \propto t^{1+3\omega}$, *i.e.* its behaviour depends on the state equation and $\Lambda \propto t^{-2}$ in any case. The scale factor f varies as $f \propto t$ in the studied models *i.e.* have no horizon problem.

1 Introduction.

Recently have been studied several models with metrics FRW in those which are considered the "constants" G and Λ as dependent functions of the time t ([1]). In this paper we want to emphasize as the use of the Dimensional Analysis (D.A.) permits us to find in a trivial way a set of solutions to this type of models, but with $k = 0$, and taking into account the conservation principle. To stand out also, that the variation of G that is obtained in this type of solutions is always proportional to the time (depending on the state equation) and that Λ always show a behaviour inversely proportional to t^2 *i.e.* $\Lambda \propto t^{-2}$ independently of the state equation that we impose ($p = \omega\rho / \omega = const.$).

The paper is organized as follows: In the second paragraph we present the equations of the model and are made some small considerations on the followed dimensional method (address to reader to the classic literature on the topic ([2])). In the third paragraph we make use of the dimensional analysis (Pi theorem) to obtain a solution to the principal quantities that appear in the model and finally in the fourth paragraph we end with a short recapitulation and succinct conclusions.

*E-mail: abelinchon@caminos.recol.es

2 The model.

The modified field equations are:

$$R_{ij} - \frac{1}{2}g_{ij}R - \Lambda(t)g_{ij} = \frac{8\pi G(t)}{c^4}T_{ij} \quad (1)$$

and we impose that

$$\text{div}(T_{ij}) = 0$$

where $\Lambda(t)$ represents the cosmological constant. The basic ingredients of the model are:

1. The line element is defined by:

$$ds^2 = -c^2 dt^2 + f^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right]$$

here only we will consider the case $k = 0$

2. The energy-momentum tensor is defined by:

$$T_{ij} = (\rho + p)u_i u_j - pg_{ij} \quad p = \omega\rho$$

The field equations, already developed, in function of the metric $g \in T_2^0(M)$ of type FRW are:

$$2\frac{f''}{f} + \frac{(f')^2}{f^2} = -\frac{8\pi G(t)}{c^2}p + c^2\Lambda(t) \quad (2)$$

$$\frac{(f')^2}{f^2} = \frac{8\pi G(t)}{3c^2}\rho + c^2\Lambda(t) \quad (3)$$

$$\text{div}(T_{ij}) = 0 \Leftrightarrow \rho' + 3(\omega + 1)\rho\frac{f'}{f} = 0 \quad (4)$$

integrating equation (4), we obtain the following equation.

$$\rho = A_\omega f^{-3(\omega+1)} \quad (5)$$

where f represents the scale factor that appears in the metrics and A_ω is the constant of integration that depends on the state equation that is imposed.

The Dimensional Analysis that we apply needs to make the following distinctions. We need to know beforehand the set of fundamental quantities, in this case it is solely the cosmic time t as is detached of the homogeneity and isotropy of the model and to distinguish the set of constants, universal and unavoidable or characteristic, that in this case, they are respectively the speed of the light c

and the constant of integration A_ω that depending on the state equation that is imposed, it will have different dimensions and physical meaning.

In a previous paper ([3]) was calculated the dimensional base of this type of models, being this $B = \{L, M, T, \theta\}$ where θ represents the dimension of the temperature. The corresponding dimensions of each quantity (with respect to this base B) are

$$[t] = T \quad [c] = LT^{-1} \quad [A_\omega] = L^{2+3\omega} MT^{-2}$$

All the quantities that we are going to calculate we will make it exclusively in function of the cosmic time t and of the unavoidable constants, c and A_ω with respect to the dimensional base $B = \{L, M, T, \theta\}$.

3 Solutions through D.A.

We are going to calculate through dimensional analysis D.A. the variation of $G(t)$ in function so much of t as of the temperature θ , $G(\theta)$ ([4]), the energy density $\rho(t)$, the radius of the universe $f(t)$, the temperature $\theta(t)$, as well as to entropy $s(t)$ and the entropy density $S(t)$ and finally $\Lambda(t)$.

3.1 Calculation of $G(t)$

As we have indicated above, we are going to accomplish the calculate applying the Pi theorem. The quantities that we consider are $G = G(t, c, A_\omega)$ with respect to the dimensional base $B = \{L, M, T, \theta\}$. We know that $[G] = L^3 M^{-1} T^{-2}$ them:

$$\begin{pmatrix} G & t & c & A_\omega \\ L & 3 & 0 & 1 & 2 + 3\omega \\ M & -1 & 0 & 0 & 1 \\ T & -2 & 1 & -1 & -2 \end{pmatrix}$$

$$G \propto \frac{t^{1+3\omega} c^{5+3\omega}}{A_\omega} \quad (6)$$

the cases more well-known are:

1. Radiation predominance $\omega = \frac{1}{3}$

$$G \propto \frac{t^2 c^6}{A_\omega} \quad G \propto t^2 \quad (7)$$

In this expression, if we want to calculate the value of the constant A_ω we can take into account the numerical values of the rest of the quantities, that is to say: $G \approx 10^{-10.1757} m^3 kg^{-1} s^{-2}$, $t \approx 10^{20} s$ and $c \approx 10^{8.5} ms^{-1}$ with this values we obtain that: $A_\omega \approx 10^{100.5} m^3 kg s^{-2}$

2. Matter predominance $\omega = 0$

$$G \propto \frac{tc^5}{A_\omega} \quad G \propto t \quad (8)$$

This result was obtained for Milne in 1935. If we consider ρ as mass density then A_ω represents the mass of the universe, the equation (6) is in this case to:

$$G \propto \frac{tc^3}{A_\omega} \quad G \propto t \quad (9)$$

this relation verify the kwon Sciama's formulate: $\rho G t^2 \approx 1$ (about the Mach's principle).

- Furthermore if we take into account the numerical values of the constant and the quantity t we obtain the nowadays value of $G \approx 10^{-10.1757} m^3 kg^{-1} s^{-2}$ i.e. $t \approx 10^{20} s$, $c \approx 10^{8.5} ms^{-1}$ and $A_\omega \approx 10^{56} kg$.
- If $\omega = -\frac{1}{3}$ then $G = const.$ though this possibility is physically unrealistic. If $\omega < -\frac{1}{3}$ then G will vary in a way inversely proportional to time.

If we want to relate G with θ (ver [4]) the solution that the D.A. provide is: $G = G(t, c, A_\omega, a, \theta)$. We need to introduce a new constant, in this case thermodynamic, to relate the temperature to the rest of quantities. The same result is obtained if we consider k_B .

$$G \propto \frac{c^4 (a\theta^4)^{\frac{\omega-1}{3(\omega+1)}}}{A_\omega^{\frac{1}{3(\omega+1)}}} \quad (10)$$

if $\omega = \frac{1}{3}$

$$G \propto \frac{c^4}{(A_\omega a)^{\frac{1}{2}} \theta^2} \quad G \propto \theta^{-2}$$

3.2 Calculation of energy density $\rho(t)$.

$\rho = \rho(t, c, A_\omega)$ that with respect to the base $B : [\rho] = L^{-1} M T^{-2}$

$$\rho \propto \frac{A_\omega}{(ct)^{3(\omega+1)}} \quad (11)$$

1. If $\omega = 1/3$ radiation predominance

$$\rho \propto \frac{A_\omega}{(ct)^4} \quad i.e. \quad \rho \propto t^{-4} \quad (12)$$

If we take into account the value of A_ω from equat. (7) we can see easily that the value of the energy density correspond with the nowadays accepted i.e. $\rho \approx 10^{-13.379} J m^{-3}$

2. If $\omega = 0$ matter predominance

$$\rho \propto \frac{A_\omega}{(ct)^3} \quad i.e. \quad \rho \propto t^{-3} \quad (13)$$

3.3 Calculation of the radius of the Universe $f(t)$.

$f = f(t, c, A_\omega)$ where $[f] = L \implies$

$$f \propto ct \quad \quad \quad f \propto t \quad (14)$$

it does not depend on A_ω i.e. it does not depend on the state equation, in both models, radiation and matter predominance, the radius of the Universe is the same. We can observe that:

$$q = -\frac{f''f}{(f')^2} = 0$$

$$H = \frac{f'}{f} = \frac{1}{t}$$

$$d_H = ct \lim_{t_0 \rightarrow 0} \int_{t_0}^t \frac{dt'}{f(t')} = \infty$$

Thus, the model has **no horizon problem** because d_H diverges for $t_0 \rightarrow 0$.

3.4 Calculation of the temperature $\theta(t)$.

$\theta = \theta(t, c, A_\omega, a)$ where the dimensional equation of the temperature is $[\theta] = \theta$ and a is the radiation constant.

$$a^{\frac{1}{4}}\theta \propto \frac{A_\omega^{\frac{1}{4}}}{(ct)^{\frac{3}{4}(1+\omega)}} \quad (15)$$

if $\omega = \frac{1}{3} \implies$

$$a^{\frac{1}{4}}\theta \propto \frac{A_\omega^{\frac{1}{4}}}{(ct)} \quad \quad \quad \theta \propto t^{-1} \quad (16)$$

If we take into account the value of A_ω from eqt. (7) we can see easily that the value of the temperature correspond with the nowadays accepted i.e. $\theta \approx 10^{0.4361}K$ or $\theta \approx 2.73^\circ K$ where the radiation constant take the value $a \approx 10^{-15.1211} Jm^{-3}K^{-4}$. We can show also that with this behaviour we obtain the well-known result:

$$\rho = a\theta^4$$

3.5 Calculation of the Entropy

$s = s(c, A_\omega, a)$ here we do not consider the time since upon imposing $div(T_{ij}) = 0$ the equations that we are considering are adiabatics i.e. they do not depend on the time. $[s] = L^2 MT^{-2} \theta^{-1}$

$$s \propto \left(A_\omega^3 a c^{3(1-3\omega)} \right)^{\frac{1}{4}} \quad (17)$$

if $\omega = \frac{1}{3} \Rightarrow$

$$s \propto (A_\omega^3 a)^{\frac{1}{4}} \quad s = const. \quad (18)$$

3.6 Entropy Density $S(t)$.

$$S = S(t, c, A_\omega, a) \quad [S] = L^{-1} MT^{-2} \theta^{-1}$$

$$S \propto \frac{(A_\omega^3 a)^{\frac{1}{4}}}{(ct)^{\frac{3}{4}(1+\omega)}} \quad (19)$$

if $\omega = \frac{1}{3} \Rightarrow$

$$S \propto \frac{(A_\omega^3 a)^{\frac{1}{4}}}{(ct)^3} \quad S \propto t^{-3} \quad (20)$$

3.7 Calculation of the cosmological constant $\Lambda(t)$

$$\Lambda = \Lambda(t, c, A_\omega); [\Lambda] = L^{-2}$$

$$\Lambda \propto \frac{1}{c^2 t^2} \quad (21)$$

it does not depend on A_ω i.e. it does not depend on the state equation, for this reason this quantity show the same behavior in both models. We can show that if $t \approx 10^{20} s$ and $c \approx 10^{8.4} ms^{-1}$ we obtain the at present admitted value of the so called cosmological constant i.e. $\Lambda \approx 10^{-56} m$.

4 Summary and conclusions.

We have solved through dimensional analysis a FRW model with $k = 0$ and with conservation of the energy-momentum tensor where G and Λ are variables with respect to the time t . We have found that $G \propto t^{1+3\omega}$ i.e. depend on the equation of state. If $\omega = \frac{1}{3}$ we obtain that $G \propto t^2$ and if $\omega = 0$, then, $G \propto t$ while, $\Lambda \propto t^{-2}$ independently from the state equation that we impose. Only it appears $G = const.$ when $\omega = -\frac{1}{3}$ state equation this physically unrealistic. The radius of the universe either depends on the state equation and it is directly proportional to the time $f \propto t$, with this radius, the model does not has the problem of the horizon and appears with $q = 0$ i.e. an always expansive universe. This result are not in disagreement with the values nowadays accepted.

ACKNOWLEDGEMENTS. I wish to thank Prof. M. Castaños for suggestions and enlightening discussions

References

- [1] **A-M. M. Abdel-Rahman.** Gen. Rel. Grav. **22**, 655,(1990). **M. S. Bermann.** Gen. Rel. Grav. **23**, 465,(1991). **Abdussaltar and R. G. Vishwakarma.** Class. Quan. Grav. **14**, 945,(1997)
- [2] **Barenblatt.**Scaling, self-similarity and intermediate asymptotics. Cambridge texts in applied mathematics N 14 1996 CUP. **Palacios, J.** Dimensional Analysis. Macmillan 1964 London.
- [3] **Belinchón, J. A.** physic/9811016.
- [4] **Zee.** Phys. Rev. Lett. **42**, 417,(1979)