

*Dedicated to the 90-anniversary of Moscow State University Professor Anatolii A. Vlasov (1908-1975)*

**STABILITY OF UNIFORM RECTILINER MOTION  
OF CHARGED SHELL  
IN SOMMERFELD MODEL WITH SELF-ACTION**

**Alexander A. Vlasov**

High Energy and Quantum Theory  
Department of Physics  
Moscow State University  
Moscow, 119899  
Russia

*It is proved that in classical electrodynamics the uniform rectilinear motion of charged "rigid" sphere with self-action (Sommerfeld model) is stable.*  
03.50.De

In classical electrodynamics long time ago in Sommerfeld works [1, see also 2] was derived the expression of self-force acting on "nonrelativistically rigid charged sphere", i.e sphere with radius  $a$ , its center moving along trajectory  $\vec{R}(t)$ , with total charge  $Q$  and charge density (in laboratory reference frame)

$$\rho(t, \vec{r}) = \frac{Q}{4\pi a^2} \delta(|\vec{r} - \vec{R}| - a)$$

In the case of shell rectilinear motion this force has the form

$$F_{self} = \frac{Q^2}{4a^2} \left[ -c \int_{T^-}^{T^+} dT \frac{cT - 2a}{L^2} + \ln \frac{L^+}{L^-} + \left( \frac{1}{\beta^2} - 1 \right) \ln \frac{1 + \beta}{1 - \beta} - \frac{2}{\beta} \right] \quad (1)$$

here  $cT^\pm = 2a \pm L^\pm$ ,  $L^\pm = |R(t) - R(t - T^\pm)|$ ,  $L = |R(t) - R(t - T)|$ .

The total shell equation of motion then will be

$$m \frac{d}{dt}(\gamma v) = F_{self} \quad (2)$$

This equation has one trivial solution - the uniform motion without radiation:

$$R(t) = R_0 + vt \quad (3)$$

For linear analysis of stability of this solution let us take

$$R(t) = R_0 + vt + \delta(t), \quad R(t-T) = R_0 + v(t-T) + \delta(t-T), \quad L = vT + \delta(t) - \delta(t-T), \quad (4)$$

here  $\delta(t)$  - small perturbation.

Then in linear approximation the self-force (1) reads

$$F_{self} = \frac{Q^2}{4a^2} \left[ \frac{4\dot{\delta}}{\beta^2} \left( 1 - \frac{1}{2\beta} \ln \frac{(1+\beta)}{(1-\beta)} \right) - \frac{2c}{v^3} \int_{T^-}^{T^+} dT \frac{cT - 2a}{T^3} \delta(t-T) \right] \quad (5)$$

here  $T^\pm = \frac{2a}{c(1 \mp \beta)}$ , and the equation of shell motion is

$$m\gamma^3 \ddot{\delta} = F_{self} \quad (6)$$

Solution of eq. (6) we take as  $\delta \sim \exp(pt)$ . Then the parameter  $p$  must obey the following equation [3]

$$p^2 = \frac{Q^2}{a^2 m \gamma^3} \left[ \frac{p}{c\beta^2} \left( 1 - \frac{1}{2\beta} \ln \frac{(1+\beta)}{(1-\beta)} \right) - \frac{c}{2v^3} \int_{T^-}^{T^+} dT \frac{cT - 2a}{T^3} \exp(-pT) \right] \quad (7)$$

This equation one can rewrite in the form:

$$q^2 = -A^2 \int_{x_-}^{x_+} dx \cdot \frac{x-2}{x^3} \cdot [qx + \exp(-qx)] \quad (8)$$

here  $q = pa/c$ ,  $x_\pm = \frac{2}{1 \mp \beta}$ ,  $A^2 = (r_{cl}/a) \cdot (1/(2\gamma^3\beta^3))$ ,  $\beta \neq 1$ .  
Dividing  $q$  into real and imaginary parts:

$$q = \eta + i \cdot w$$

we split eq.(8) into two:

$$\eta^2 - w^2 = -A^2 \int_{x_-}^{x_+} dx \cdot \frac{x-2}{x^3} \cdot [\eta x + \exp(-\eta x) \cos(wx)] \quad (9a)$$

$$2\eta w = -A^2 \int_{x_-}^{x_+} dx \cdot \frac{x-2}{x^3} \cdot [xw - \exp(-\eta x) \sin(wx)] \quad (9b)$$

Equations (9) are invariant under transformations  $w \rightarrow -w$ , so both signs of  $w$  are solutions, thus further we can take  $w$  as positive quantity.

Integrand in (9b) one can divide into two functions:

$$f_1 = \frac{x-2}{x^3}$$

and

$$f_2 = [xw - \exp(-\eta x) \sin(wx)]$$

Integral of  $f_1$  over  $x$  between limits  $x_{\pm}$  has zero value;  $f_2$  - is positive function with positive derivative for  $\eta, w > 0$  and  $x > 0$ . Thus for these values of  $\eta$  and  $w$  the integral of  $f_1 \cdot f_2$  is positive and the sign of R.H.S. is negative. But this result contradicts with the sign of L.H.S., which is positive for  $\eta > 0, w > 0$ . Consequently the equations (9) have solutions only for negative values of the real part  $\eta$  of  $q$ .

This proves that the uniform rectilinear motion of Sommerfeld charged sphere with  $\beta < 1$  is stable!

## REFERENCES

1. A.Sommerfeld, Gottingen Nachrichten, 29 (1904), 363 (1904), 201 (1905).
2. P.Pearle, in *Electromagnetism*, ed. D.Tepliz, Plenum, NY, 1982, p.211.
3. Alexander A.Vlasov, physics/9804001.