$\mathcal{U}$ Ruhr-Universität<br>Bochum Institut für Neuroinformatik

Internal Report 98–09

# Evolution in time-dependent fitness landscapes

by Claus O. Wilke

Ruhr-Universität Bochum Institut für Neuroinformatik 44780 Bochum



IR-INI 98–09 November 1998 ISSN 0943-2752

 $\odot$  1998 Institut für Neuroinformatik, Ruhr-Universität Bochum, FRG

# Evolution in time-dependent fitness landscapes

Claus O. Wilke<sup>∗</sup>

#### Abstract

Evolution in changing environments is an important, but little studied aspect of the theory of evolution. The idea of adaptive walks in fitness landscapes has triggered a vast amount of research and has led to many important insights about the progress of evolution. Nevertheless, the small step to time-dependent fitness landscapes has most of the time not been taken. In this work, some elements of a theory of adaptive walks on changing fitness landscapes are proposed, and are subsequently applied to and tested on a simple family of time-dependent fitness landscapes, the oscillating NK landscapes, also introduced here. For these landscapes, the parameter governing the evolutionary dynamics is the fraction of static fitness contributions  $f_s$ . For small  $f_s$ , local optima are virtually non-existent, and the adaptive walk constantly encounters new genotypes, whereas for large  $f<sub>S</sub>$ , the evolutionary dynamics reduces to the one on static fitness landscapes. Evidence is presented that the transition between the two regimes is a 2nd order phase transition akin a percolation transition. For  $f<sub>S</sub>$ close to the critical point, a rich dynamics can be observed. The adaptive walk gets trapped in noisy limit cycles, and transitions from one noisy limit cycle to another occur sporadically.

## 1 Introduction

The key element of Darwinian evolution is natural selection, which prefers advantageous mutations and rejects deleterious ones. It is possible to separate any evolutionary system under study into two components: a single species (or population of individuals) and an environment. The individuals in the population change because of mutations, and the environment discriminates between better adapted ones (the ones generating more offspring, i.e., the "fitter" ones) and worse adapted ones. The difference in the number of offspring suffices to remove efficiently the worse adapted individuals from the population. Clearly, the environment does not necessarily remain constant in time. It consists of all the influences acting on the species, and it therefore contains a-biotic and biotic components. Changes may have an a-biotic origin (e.g., change in mean temperature, change in sea level) or a biotic origin (e.g., other species in the same ecosystem changing because of adaption). The latter case is usually referred to as coevolution. From the viewpoint of a single species that adapts in an environment, all evolutionary scenarios can be divided into three distinct classes.

These classes are, as depicted in Fig. [1,](#page-2-0) i) evolution in a constant environment, ii) evolution in a variable environment without feedback to the environment, and iii) evolution with feedback to the environment. The last class is the coevolutionary case mentioned above. The current knowledge about evolutionary dynamics is very unevenly distributed over these three classes. An overwhelming amount of theoretical results is known

<sup>∗</sup> email claus.wilke@ruhr-uni-bochum.de

<span id="page-2-0"></span>for class i). Most of the work in the field of population genetics has been done with constant environment  $(e.g. [6, 9, 17])$  $(e.g. [6, 9, 17])$  $(e.g. [6, 9, 17])$  $(e.g. [6, 9, 17])$ , and also the idea of adaptive walks in fitness landscapes, introduced by Wright [\[27](#page-16-0)], and intensively studied for example by Kauffman[[14\]](#page-16-0), has usually only been applied to constant fitness landscapes. Class iii) has also been studied intensively. One the one hand, there exists a rich theoretical ecology literature (e.g. [\[8, 11\]](#page-15-0)), dealing with predator-prey interactions such as Lotka-Volterra systems, or with host-parasite interactions. On the other hand, over the last decade, coevolutionary scenarios have been studied through direct competition of two or more species in Artificial Life style computer simulations (e.g.[[12,](#page-15-0) [15](#page-16-0), [22](#page-16-0)]). The evolutionary patterns observed there are often very complex, and a clear theoretical understanding of these patterns has not been obtained yet. Finally, Class ii), which lies between classes i) and iii), and which could therefore yield many new



insights about the connection between adaptive walks and coevolution, has nevertheless widely been neglected. There exists some work on evolution in randomly fluctuating environments[\[10](#page-15-0)], but this work obviously cannot cover the wide range of possible scenarios within class ii). In particular, a theory for the natural generalization of class i) adaptive walks, i.e., adaptive walks on a changing fitness landscape, is still missing.

## 2 Environmentally guided drift

Evolutionary processes in static fitness landscapes tend to get stuck quickly in local optima, often far away from some globally optimal solution. This hinders evolution's progress, and severely restricts the amount of possible solutions that can be found during the adaptive progress. Nevertheless, Nature's rich diversity of extremely clever solutions to a large array of different problems indicates that some mechanisms are driving evolution out of local optima. Probably, the most common proposal for such a mechanism is the idea of neutral evolution, introduced by Kimura[[17\]](#page-16-0), and based on the observation that the genotype-phenotype mapping of living beings discards information. Certain mutations can appear without change in the organism's fitness, and therefore, without being constrained by natural selection. The accumulation of such neutral mutations is called neutral drift. In principle, the neutral drift could allow a certain gene to drift far away from its original position in genome space, and then to change to a very much advantageous state by

<span id="page-3-0"></span>means of a single point mutation. There exists a lot of interesting theoretical results on this topic[[7,](#page-15-0) [13](#page-16-0), [20](#page-16-0), [21\]](#page-16-0). However, on the level of proteins, there is not much evidence in favor of this viewpoint [\[2,](#page-15-0) [19\]](#page-16-0). Clearly, there are neutral mutations on the level of DNA (different codons code for the same amino acid) or in the RNA folding problem, but on the level of amino acid sequences almost every change seems to have some selective impact.

One out of the numerous investigations demonstrating the absence of neutrality on the protein level is the following study of the  $Fast/ Slow$  (Adh-F/Adh-S) polymorphism at the alcohol dehydrogenase locus (Adh) of Drosophila melanogaster. The Fast/Slow polymorphism is probably the most intensively studied polymorphism at all, with a number of 364 scientific research papers written on it already back in 1988 [\[4\]](#page-15-0). Kreitman [\[18](#page-16-0)] sequenced eleven cloned *Drosophila m. Adh* genes from five natural populations. Forty-two out of 43 differences in the DNA sequence were silent, only one base-pair substitution actually caused a change in the amino acid sequence. This base-pair substitution constituted exactly the  $Adh-F/Adh-S$  polymorphism. The immediate conclusion that could be drawn was that most amino acid changes in alcohol dehydrogenase would be selectively deleterious for the wild-type *Drosophila m.* Other studies demonstrate that the Fast/Slow polymorphism in itself is sensitive to some selective pressure. The population's composition of  $Adh-F$  and  $Adh-S$  mutants was found to change with the geographical latitude (latitudinal cline) [\[24](#page-16-0), [26\]](#page-16-0). In a recent study, Berry and Kreitman[[3](#page-15-0)] found a cline along the East Coast of North America that could only be observed in the amino acid responsible for the  $Adh-F/Adh-S$  polymorphism, but not in the accompanying silent mutations. This finding rejects neutral drift as the origin of the cline.

Comprehensive reviews on the subject of protein evolution's sensitivity to selective pressures can be found in[[2](#page-15-0), [19](#page-16-0)].

Another mechanism capable of driving evolution away from local optima is a slowly changing environment, as proposed by Benner and Ellington [\[2\]](#page-15-0). If most changes in the amino acid sequence of a certain protein are not selectively neutral, then, on the other side, most changes in the selective force will have some impact on the amino acid sequence of this protein. A slowly changing environment will then (on an evolutionary time scale) drag genes around in genotype space. This results in another form of genetic drift, which can happen under complete absence of neutrality. However, this drift is not a diffusion process such as neutral drift. Instead, if the environmental changes happen slowly, the population as a whole will move through the genotype space, since transitions to a selectively advantageous state happen very fast, as first order phase transitions[[1](#page-15-0)]. Such environmental influences on the evolutionary dynamics have rarely been considered, and the kind of dynamics they will produce is not at all understood. In the reminder of this paper, we will investigate the influence of a slowly changing environment for a simple example of a time-dependent fitness landscape. Note that we will only be interested in environmental changes happening on time-scales much larger than the population's typical transition times to states with higher fitness. Therefore, we can restrict our investigations to the study of adaptive walks[[14\]](#page-16-0). But, before we embark upon this endeavor, we have to define some useful notions.

First of all, we are going to define the concept of environmentally guided drift of a gene. This concept will be useful in the further study of evolution in time-dependent fitness landscapes.

**Definition 1 (Environmentally guided drift)** A gene is said to drift environmentally guided if in every jump from one point in genotype space to another, the gene's fitness in this time step remains either constant or increases.

<span id="page-4-0"></span>This definition includes neutral drift as a special case, and it also includes those adaptive walks in which transitions to lower fitnesses are forbidden. Note that the fitness can nevertheless decrease during environmentally guided drift because of environmental changes. However, this decrease is completely due to a change in the structure of the underlying fitness landscape, and not to a movement in the fitness landscape.

Neutral drift has been included in the Definition [1](#page-3-0) to make the term environmentally guided drift sufficiently general. However, our main interest in this paper lies in fitness landscapes that do not contain any neutrality.

Definition [1](#page-3-0) reads very much like the definition of standard hill-climbing, and is of course nothing else in essence. Nevertheless, it seems appropriate to introduce a new notion to remind ourselves that the dynamics changes drastically in a time-dependent fitness landscape. In particular, a gene may appear to be in a local optimum for almost every single time step, and nonetheless it may travel constantly due to the changing positions of the local optima.

As in the case of pure neutral drift, the environmentally guided drift could allow a gene to arrive at a point in genotype space that would not have been accessible by means of simple hill-climbing on a fixed fitness landscape. Since environmental changes are often oscillatory (seasons, for example, happen on an evolutionary time scale for microorganisms) or pseudo-oscillatory (for example, cf. Fig. 2 in [\[23](#page-16-0)]), a species most likely encounters the same environmental state several times, and can use the intermediate times of environmental guided drift for further adaption.

## 3 Environmentally linked networks

An important ingredient of the neutral theory of evolution is the notion of neutral networks. A neutral network is formed by points in the genotype space that have identical fitnesses and that are connected by elementary mutations. The neutral drift is a diffusion process over these networks. In analogy to the situation of neutral evolution, we can define structures on which the environmentally guided drift naturally takes place. These structures will in the following be called environmentally linked networks or short EL networks.

EL networks can be defined on any time-dependent fitness landscape that has identical states, i.e., on fitness landscapes for which there exists a sequence of times  $t_0, t_1, t_2, \ldots$ , such that

$$
\mathcal{F}(t_0) = \mathcal{F}(t_1) = \mathcal{F}(t_2) = \dots \,, \tag{1}
$$

where  $\mathcal{F}(t) = \mathcal{F}(t')$  means that all points in genotype space have the same fitness at times  $t$  and  $t'$ .

Definition 2 (Environmentally linked networks) An environmentally linked network is the set of all those points in the genotype space that can be reached by means of environmentally guided drift at times  $t_i$ , for which  $\mathcal{F}(t_0) = \mathcal{F}(t_i)$ , starting from some fixed genome at time  $t_0$ .

The EL networks according to Definition 2 contain all the points in the genotype space that can in principle be reached by sole hill-climbing. The probability to reach some of these points may be very low. Therefore, these points will most certainly never be encountered. As a consequence, EL networks are too "large" for practical purposes, <span id="page-5-0"></span>they contain many points with no relevance for the evolutionary dynamics. To resolve this problem, we define smaller structures, in which the most unlikely points are removed.

Definition 3 ( $\epsilon$ -environmentally linked networks) The set of all the points  $x_i$  in an EL network for which a sequence of points  $x_0, x_1, \ldots, x_{i-1}, x_i$  in the EL network exists, such that every transition from  $x_{j-1}$  to  $x_j$  can occur with probability  $\epsilon^{t_j-t_{j-1}}$ , is called an  $\epsilon$ -environmentally linked network.

This definition immediately gives a trivial upper bound to the number of points  $n_t$  that can belong to the  $\epsilon$ -EL network after t evolution time steps:

$$
n_t \le \frac{1}{\epsilon^t} \,. \tag{2}
$$

A question that arises in conjunction with the above definitions is whether environmentally guided drift can produce, in absence of neutrality, analogues of percolating neutral networks, i.e., EL networks whose points are close in Hamming-distance to any arbitrary point in the genotype space. The existence, or non-existence, of percolating networks determines whether a mechanism is efficient in generating evolutionary innovation.

## 4 Oscillating NK-models

Environmental changes can be modelled in many different ways. For example, the fluctuations can be stochastic or deterministic, continuous or discontinuous, spatially resolved or unresolved. Clearly, each of the different possibilities has its reason of being. Gillespie has studied the effects of selection in a randomly fluctuating environment[[10](#page-15-0)]. In his models, the fluctuations' time scale is comparable to the duration of one generation of the organisms considered. As a result, the genotype with the largest geometric mean fitness dominates the population, and ultimately wins over all competing genotypes. In the present work, we are interested in a different situation. The subject of this work are environmental changes that happen so slowly that the population as a whole moves to a more advantageous genotype once it is discovered. In this situation, a randomly fluctuating environment does not seem to be appropriate. Rather, we are going to consider a continuous, deterministic model. For reasons of simplicity, we will also neglect any spatial structure. A changing environment of this kind can easily be included into the well-known NK family of fitness landscapes [\[14](#page-16-0), [16](#page-16-0)], standard landscapes commonly used for the study of adaptive walks. An  $NK$ -landscape is a fitness landscape for a bitstring of N bits. The fitness of a bitstring is the average over the fitness contributions of all bits in the string. The contribution of a single bit depends on its state and on the state of  $K$  other bits interacting with it. Mathematically, we can write the fitness  $f$  as:

$$
f = \frac{1}{N} \sum_{i=1}^{N} f_i(S_i, S_{\alpha_{i1}}, \dots, S_{\alpha_{iK}}),
$$
\n(3)

where  $S_i$  is the state of the *i*th bit (zero or one),  $\alpha_{ij}$  is the number of the *j*th bit interacting with bit i, and  $f_i(\cdot)$  is a function that maps the state of the ith bit and its interacting bits onto a real number between 0 and 1. In the following, we will write the state of the  $K+1$  bits  $S_i, S_{\alpha_{i1}}, \cdots, S_{\alpha_{iK}}$  as  $\{S\}_i$ . Usually, the functions  $f_i(\{S\}_i)$  are realized as tables containing a different fitness contribution  $C_{\alpha}$  for every possible arrangement of bits:

$$
f_i({S}_i) = C_{i,{S}_i}.
$$
\n(4)

A straightforward generalization of this concept is reached by making the functions  $f_i({S}_i)$  time dependent. Instead of the constants on the right-hand side of Eq. [\(4\)](#page-5-0) we then have to use time-dependent functions:

$$
f_i(\{S\}_i, t) = F_{i, \{S\}_i}(t). \tag{5}
$$

The functions  $F_{i,\{S\}_i}(t)$  could in principle be arbitrarily complicated. Here, we are going to consider a simple trigonometric time dependency, i.e.,

$$
F_{i,\{S\}_i}(t) = \frac{1}{2} [\sin(\omega_{i,\{S\}_i} t + \delta_{i,\{S\}_i}) + 1].
$$
 (6)

This introduces only a single additional constant per fitness contribution, since there are now two of them (the frequency  $\omega_{i,\{S\}_i}$ , and the phase  $\delta_{i,\{S\}_i}$ ). The frequencies  $\omega_\alpha$  and the phases  $\delta_{\alpha}$  are chosen randomly at the beginning of the evolutionary process, and then kept fixed for all times t. The phases have to be distributed uniformly in the interval  $[0, 2\pi)$ if the resulting fitness landscape is supposed to be homogeneous in time. The choice of the frequencies determines whether the fitness landscape is periodic in time. If we set the frequencies to  $\omega_{i,\{S\}_i} = 2\pi n_{i,\{S\}_i}/T$ , with  $n_{i,\{S\}_i}$  being integral, and T being arbitrary, but the same for all  $\omega_{i,\{S\}_i}$ , we obtain a periodic fitness landscape with oscillation period T. Throughout the rest of this paper, we will stick to this choice. Its advantage rests in the easy comparison of a bistring's evolution in different oscillation periods.

If we set a frequency  $\omega_{i,\{S\}_i}$  to zero, the according fitness contribution  $F_{i,\{S\}_i}(t)$  is constant in time. The fraction of these non-oscillatory (static) fitness contributions will be called  $f<sub>S</sub>$  in the following. The quantity  $f<sub>S</sub>$  is the total number of static fitness contributions, divided by the total number of all fitness contributions. Adaptive walks on oscillating  $NK$ -landscapes show several distinct classes of behavior, most strongly influenced by  $f_s$ .

#### 4.1 Numerical experiments

We have done several numerical experiments with oscillating  $NK$ -landscapes. In all simulations presented below, only one oscillating mode was used. This means, all frequencies  $\omega_i$  were set either to zero or to some fixed value  $\omega$ . If not indicated otherwise, the value  $\omega = 2\pi/200$  was used. In Figures [2-](#page-8-0)[4](#page-9-0) some typical runs of adaptive walks in oscillating NK-landscapes are presented. The adaptive walk was performed exactly as in Kauffman's original work: a random point mutation was accepted if it increased the bitstring's fitness. Otherwise, the mutation was rejected. Hence, the adaptive walk per definition took place on an EL network. Fig. [2](#page-8-0) shows an example of the evolutionary dynamics with a relatively low fraction of static fitness contributions. The resulting pattern is a chaotically changing fitness. With almost every accepted mutation, a new genotype is encountered. The environmentally guided drift resembles a random walk on the fitness landscape. Interestingly, this is the case even for small K.  $NK$ -landscapes with small K are highly correlated, which should bias the adaptive walk. Nevertheless, it seems the oscillations destroy the correlations' effect. The picture changes drastically with increasing  $f_S$ . The higher amount of static fitness contributions reduces the number of possible advantageous mutations in every time step. The sites connected to static contributions freeze out in a locally optimal state, and only the sites connected to oscillating contributions can still change. Hence, the dynamics gets confined in a small region of the genotype space. The same mutational patterns are seen over and over again in the different oscillation periods. In the fitness plots, we can identify this behavior with a periodic or almost periodic change of the fitness, as shown in Fig. [3.](#page-9-0) Using the language of dynamic systems, we can say that the attractor of an adaptive walk on an oscillating landscape with intermediate  $f<sub>S</sub>$  is a limit cycle. Since the dynamics we are investigating here is not deterministic, the limit cycles we observe will generally be noisy. Hence, the limit cycles have to be understood in the sense of the  $\epsilon$ -EL networks defined above. With a certain small probability  $p < \epsilon$ . the process can leave an attractor, and find a new one. A transition from one attractor to another is shown in Fig. [4.](#page-9-0) In this case, the mean fitness has significantly increased after the transition.

A detailed analysis of the transition in Fig. [4](#page-9-0) reveals the typical behavior of noisy limit cycles in oscillating NK-models. We begin our analysis at time  $t = 6000$ . The dynamics proceeded as follows. Underlined bits have been mutated in the corresponding time step.



We find a limit cycle with two states. At the beginning of each period, bit 5 is switched from 0 to 1. Towards the end of the period, the bit is switched back to 0. Remember we used  $T = 200$  in this run, hence oscillation periods started at  $t = 6000, 6200, 6400, \ldots$ The exact times in which bit 5 was switched differ from period to period, because of the nondeterministic dynamics. Further 33 mutations occurred in bit 5 until a mutation in bit 4 in time-step  $t = 10110$  generated a rapid movement towards a far away region in the genotype space:



Further 40 mutations occurred between  $t = 10114$  and  $t = 11301$ , until finally, in  $t = 11302$ , a new stable configuration was found:

<span id="page-8-0"></span>

Figure 2: The evolutionary dynamics is chaotic for small  $f_S$ .



This time, the dynamics had found a stable fix point. No further mutations occurred until the simulation was aborted in time-step  $t = 40000$ .

At this point, it is interesting to ask whether all limit cycles will be left for  $t \to \infty$ , or whether the dynamics inevitably encounters a stable limit cycle eventually. A possible answer to this question will be given in Sec. [4.3](#page-12-0), where we present evidence for the existence of a percolation transition in the region of intermediate  $f<sub>S</sub>$ . For  $f<sub>S</sub>$  above the critical value, the dynamics would sooner or later reach a stable fixed point, or a stable limit cycle, whereas for  $f<sub>S</sub>$  below the critical value, every limit cycle which is part of the infinite (in the limit  $N \to \infty$ ) EL network will be eventually left.

#### 4.2 Short oscillation periods

The oscillation period used in the previous section was relatively long, if compared to the length  $N$  of the bitstring. In such a situation, a single site can be hit repeatedly by mutations during one oscillation period. This generates complicated time correlations, which are hard to tackle theoretically. The correlations can be neglected in situations in which every site is, on average, hit at most once during one oscillation period. This is assured if the number of time-steps  $T$  in one oscillation period is small compared to the number of bits  $N$  in the string,

$$
T \ll N. \tag{7}
$$

<span id="page-9-0"></span>

Figure 3: With increasing  $f_s$ , some bits in the bitstring freeze out, and the evolutionary pattern becomes more and more oscillatory.



Figure 4: A transition from one oscillatory state to another, with higher mean fitness.

<span id="page-10-0"></span>The case  $f<sub>S</sub> = 0$  can be treated analytically in this regime. Again, we consider only a single frequency  $\omega$ , i.e.,  $\omega_i = \omega$  for all i. Let us begin with a bitstring of N bits without epistatic interactions, i.e., with  $K = 0$ . In this case, the mutation of a single bit will change a single fitness contribution  $f_i(t)$  to a new form  $f'_i(t)$ . With what probability will the new fitness contribution exceed the old one at time t? Since both  $f_i(t)$  and  $f'_i(t)$  can be considered being chosen randomly from a large set of possible functions, the probability equals the fraction of the interval  $[0; T)$  in which the function  $f_i'(t)$  exceeds the function  $f_i(t)$ . This fraction is easily calculated. We compute the intersecting points  $t_j^*$  from  $f_i(t^*) = f'_i(t^*)$ :

$$
\frac{1}{2} \left[ \sin(\omega t^* + \delta_i) + 1 \right] = \frac{1}{2} \left[ \sin(\omega t^* + \delta'_i) + 1 \right]. \tag{8}
$$

With the use of a trigonometric identity, we find

$$
2\cos\left(\omega t^* + \frac{\delta_i + \delta'_i}{2}\right)\sin\left(\frac{\delta_i - \delta'_i}{2}\right) = 0.
$$
\n(9)

Since  $\delta_i = \delta'_i$  with probability 0, we assume that  $\delta_i$  and  $\delta'_i$  are not equal, and find

$$
t_1^* = \left(\frac{1}{4} - \frac{\delta_i + \delta_i'}{4\pi}\right)T\,,\tag{10}
$$

$$
t_2^* = \left(\frac{3}{4} - \frac{\delta_i + \delta_i'}{4\pi}\right)T.
$$
\n(11)

Hence,  $f_i'(t)$  will exceed  $f_i(t)$  half of the period, and will be smaller than  $f_i(t)$  during the other half. Therefore, the probability  $p_{\text{climb}}$ , with which a random mutation will lead to a higher fitness, is  $p_{\text{climb}} = 1/2$ . The resulting bitstring's motion in the genotype space is a random walk, in which jumps to neighboring positions occur only with probability 1/2.

The above result can easily be extended to higher K. The probability  $1/2$  actually stems from the symmetry that the fitness contributions possess as trigonometric functions:

$$
f_i(t+T/2) - 1/2 = -\left(f_i(t) - 1/2\right). \tag{12}
$$

With Eq. (12), from  $f_i'(t) > f_i(t)$  follows  $f_i'(t+T/2) < f_i(t+T/2)$ . Half of the interval is dominated by  $f_i(t)$ , the other half by  $f'_i(t)$ . A similar equation exists for the case  $K > 0$ . For positive K, a single mutation affects the fitness contributions of  $K + 1$  bits.  $K + 1$ fitness contributions change. Without loss of generality, assume they are numbered from 1 to  $K + 1$ . Then, a mutation is accepted if the sum of the changed fitness contributions exceeds the sum of the original fitness contributions, i.e., if

$$
\frac{1}{K+1} \sum_{j=1}^{K+1} f'_j(t) \ge \frac{1}{K+1} \sum_{j=1}^{K+1} f_j(t).
$$
 (13)

The symmetry property Eq. (12) of the single fitness contributions can be transferred to the sum  $\sum_{j=1}^{K+1} f_j(t+T/2)/(K+1)$ . With Eq. (12), we find

$$
\frac{1}{K+1} \sum_{j=1}^{K+1} f_j(t+T/2) - 1/2 = \frac{1}{K+1} \sum_{j=1}^{K+1} \left( -f_j(t) + 1 \right) - 1/2
$$

$$
= -\left( \frac{1}{K+1} \sum_{j=1}^{K+1} f_j(t) - 1/2 \right), \tag{14}
$$



Figure 5: The movement in Hamming distance  $d_h$  away from the original string for 4 different experiments. The upper 3 graphs stem from oscillating  $NK$  models. The oscillation period was  $T = N = 50$  in all cases. The bottom graph stems from a random walk with a probability of 1/2 for resting one time-step.

which is the exact equivalent of Eq.  $(12)$  for positive K. Thus, for a given set of fitness contributions  $f_j(t)$ ,  $f'_j(t)$ ,  $j = 1, 2, ..., K + 1$ , the sum of the contributions  $f_j(t)$  will dominate half of the interval  $[0; T)$ , and it will be dominated on the other half. Clearly, in the case  $K > 0$  the interval  $[0; T)$  is divided into more than 2 subintervals. However, the measure associated with the subintervals in which  $\sum f_j'(t) > \sum f_j(t)$  equals the measure associated with the subintervals in which  $\sum f_j'(t) < \sum f_j(t)$ . The points for which both sums equal have measure zero, and can be neglected. The fact that we now have to deal with several subintervals, instead of just 2, does not defeat the argument given above for  $K = 0$ . Therefore, the dynamics for  $f_S = 0$  and  $T \ll N$  is a random walk, independent of the choice of K. For all K, the probability that the next step in the random walk is taken is 1/2.

We have tested our analytical results in numerical experiments. The adaptive walk behaves indeed very similar to a random walk for  $f_s = 0$ . In Fig. 5, we have displayed the trajectories generated during the first 1000 time-steps in 4 different computer experiments. Since it is hard to visualize a high-dimensional trajectory, we plotted the Hamming distance  $d_H$  between the bitstring at time t and the bitstring at time 0. Three of the experiments were done with oscillating NK-models, the forth one was a simple random walk, used as a comparison. In all 3 experiments with oscillating landscapes, we used  $T = N/2 = 50$ . The qualitative behavior is very similar in all 4 plots. In all cases, the

<span id="page-12-0"></span>

Figure 6: The probability  $p_{\text{climb}}$  versus the oscillation period T for different values of K. We used  $N = 100$  in all 4 cases. For small T, we observe a plateau at  $p_{\text{climb}} = 0.5$ , whereas for large  $T$ ,  $p_{\text{climb}}$  decays to zero. The decay happens the faster, the larger we set  $K$ .

walker quickly moves away from the initial position in genome space. For the rest of the run, the walker's Hamming distance from the initial position fluctuates about  $d_{\rm H} = 50$ . This is to be expected for a random walk with  $N = 100$ . On average, in two randomly chosen sequences, half of the bits are the same. Therefore,  $d_H$  fluctuates about  $N/2$ .

Some more insight can be gained from a direct measurement of  $p_{\text{climb}}$  for different period lengths  $T$ . The result of such a measurement is presented in Fig 6. For small  $T$ ,  $p_{\text{climb}}$  forms a plateau around 0.5, as it was predicted by our analysis above. For large times  $T$ , the probability  $p_{\text{climb}}$  decays to zero.

#### 4.3 EL networks and the percolation transition

In this section, we will study the question of the existence of a percolating regime for small  $f_S$ . In the following, we will only be interested in EL networks, i.e., we observe the dynamics only at the beginning of each oscillation period, and disregard what is going on in intermediate times. We will say an EL network percolates if it consists of infinitely many points. This definition is similar to the usual definition of the percolating cluster on the Bethe lattice, and is the appropriate way to define percolation in high-dimensional spaces[[25\]](#page-16-0). Clearly, it can be applied literally only in the limit  $N \to \infty$ . However, the genotype space grows so fast with increasing  $N$  that this restriction can be neglected.

The study of EL networks in oscillating NK landscapes is computationally very demanding, since we have to go through the full oscillation periods in the simulation. Hence, we restrict ourselves to a single case we study as an example. We use  $\omega_i = \omega = 2\pi/T$  with  $T = 200$ , as we have done in the previous sections. Moreover, we consider only the case  $N = 20.$ 

Figure [7](#page-13-0) shows the fraction  $\gamma$  of new genotypes among all the genotypes encountered at the beginning of each oscillation period. This is a measure for the size of an EL network.

<span id="page-13-0"></span>

Figure 7: The fraction of newly encountered genotypes at the beginning of each oscillation period in oscillating fitness landscapes. For small  $f<sub>S</sub>$ , a new genotype is encountered in almost every adaptive move. With increasing  $f_{\rm S}$ , the fraction of new genotypes decreases to zero. This happens the faster the larger the value of K.

A value near 1 means a new genotype has been encountered in almost every oscillation period. On the other hand, a value near 0 means the network's size is small, thus confining the adaptive walk in a limited region of the genotype space. In the limit of infinitely many oscillation periods, only percolating networks can have a positive  $\gamma$ , whereas finite networks yield  $\gamma = 0$ . Therefore,  $\gamma$  is a proper order parameter to indicate a percolation transition. Clearly, in numerical experiments the number of oscillation periods over which the measurement is taken is finite, and therefore we will observe a positive  $\gamma$  even in the non-percolating regime. In the case of Fig. 7, for example, the value  $\gamma$  was obtained from averaging over 60 adaptive walks, each on a different fitness landscape. Every adaptive walk endured 200 oscillation periods. The error bars present the standard deviations of the single measurements.

Let us begin the discussion of Fig. 7 with the two graphs in the bottom row, for  $K = 8$ and  $K = 14$ . In both cases,  $\gamma = 1$  for small  $f_S$ , and  $\gamma = 0$  for  $f_S \approx 1$ . The standard deviations are very small in both limiting regimes. In the region about  $f_s \approx 0.5$ , a sharp drop in  $\gamma$  can be observed, accompanied with an enormous increase in the error bars. The large error bars are a sign for critical fluctuations, which are observed in 2nd order phase transitions. The picture is not as clear in the upper row of Fig. 7, for  $K = 0$  and  $K = 2$ . The error bars are large for the whole range of  $f_s$ , and  $\gamma$  does not reach the value 1 for  $f_{\rm S} = 0$ . This implies a percolating El network, if it exists for small K (below we will give an argument why this is probably not the case for  $K = 0$ , does not span the whole genotype space in this case, even in the limit  $f_s = 0$ . The transition from the percolating to the non-percolating regime has to be much softer, and the critical value  $f_S^*$  cannot be estimated from our measurements.

Measurements with an extremely increased accuracy are necessary to determine the

critical point  $f_S^*$  exactly, and to decide whether a percolation transition does exist at all for small K. This work is currently being undertaken, and will be presented elsewhere.

The measurements in Fig. [7](#page-13-0) reveal another surprising fact. For small  $f_S$ ,  $\gamma$  is increased with increased ruggedness of the landscape. The "Fujiyama" landscape  $K = 0$  has a much smaller  $\gamma$  than the relatively rugged landscape with  $K = 8$ . This means ruggedness promotes the movement in the genotype space for the oscillating regime, a situation completely opposite to the case of fixed fitness landscapes, where ruggedness is regarded as an impediment to the movement in the genotype space. We can understand this observation in the "adiabatic" limit, where changes in the fitness landscape happen so slowly that the adaptive walk can always reach the next local optimum before another change happens. In this limit, the EL network of the Fujiyama landscape degenerates to a single point. On the otherhand, the completely random Derrida landscape [[5](#page-15-0)] which we obtain for  $K = N - 1$ presents a multitude of local optima, and the changes in the fitness landscape provide the opportunity to hop from one local optimum to another in a random fashion during the oscillation periods.

## 5 Conclusions

Evolution in a slowly changing environment follows a dynamics very different from the situation in a fixed environment. Environmentally guided drift drives genes out of local optima, and drags them around in the genotype space. The evolution generally proceeds on  $\epsilon$ -EL networks. However, transitions from one  $\epsilon$ -EL network to another occur sporadically. In this work, we have presented evidence for the existence of a giant  $\epsilon$ -EL network in the limit of a vanishing amount of static fitness contributions. Consequently, in this limit the whole genotype space can be transversed by means of environmentally guided drift. Hence, the guided drift can provide – in absence of any neutral pathways in the fitness landscape – an efficient mechanism to generate constantly new genotypes, albeit at every single point in time the system seems to be trapped in a local optimum. In Sec. [4.3](#page-12-0), we could show that the efficiency of the environmentally guided drift is related to the ruggedness of the landscape. A more rugged landscape provides more opportunities to move around under environmental changes than a landscape with only a few peaks. Consequently, the rugged landscapes observed in protein evolution [\[2\]](#page-15-0) could accelerate protein evolution in an ever changing environment, instead of hindering it.

Above the percolation transition, the  $\epsilon$ -EL networks are relatively small. The dynamics in this regime is dominated by noisy limit cycles. The system goes through several noisy limit cycles until a stable limit cycle, or a stable fixed point, is reached.

The model studied in the present paper, i.e., the family of oscillating  $NK$ -landscapes, is certainly too simplistic to be accounted for as a realistic model of the evolution of a species in a changing environment. In particular, it can be argued whether oscillating fitness contributions, taking on all values between zero and one during one oscillation period, are justified. Nevertheless, such a simple model can be used to get a first impression of how evolution in a slowly changing environment could behave. Furthermore, a simple model can guide the development of tools and concepts that can then be applied to more complicated situations. The idea of environmentally linked networks is such a concept. EL networks can be defined on any fitness landscape with recurrent states. The restriction to such fitness landscapes may seem narrowing on the first sight, but it is justified in a number of reasonable situations, like the evolution of microorganisms under the change of seasons, as mentioned in Sec. [2](#page-2-0). Generally, two states of a system at different points <span id="page-15-0"></span>in time can only be compared under similar boundary conditions. Therefore, a condition like Eq. [\(1](#page-4-0)) is necessary to draw meaningful comparisons. In particular, in evolution two different adaptive solutions can only be compared if the problem to solve is identical.

In future work, it should be interesting to study the percolation transition in more detail, and to determine for what  $K$  a percolating regime actually exists. Furthermore, higher oscillation modes should be considered and the influence of the string length N and the period length  $T$  should be investigated. Investigations similar to the ones we have presented in this work can also be done for other time-periodic fitness landscapes.

## 6 Acknowledgments

I would like to thank Steven Benner, who made me aware of the strong constraints on the evolution of proteins imposed by natural selection, and of the environmental changes' consequent importance to innovation in evolution.

### References

- [1] Christoph Adami. Introduction to Artificial Life. Telos, Springer-Verlag Publishers, Santa Clara, 1998.
- [2] Steven Benner and Andrew D. Ellington. Interpreting the behavior of enzymes: purpose or pedigree? CRC Critical Reviews in Biochemistry, 23:369–425, 1988.
- [3] A. Berry and M. Kreitman. Molecular analysis of an allozyme cline: Alcohol dehydrogenase in Drosophila melanogaster on the East Coast of North America. Genetics, 134:869–893, 1993.
- [4] G. K. Chambers. The Drosophila alcohol dehydrogenase gene-enzyme system. Adv. Genet., 25:39–107, 1988.
- [5] Bernard Derrida. Random energy model: An exactly solvable model of disordered systems. Phys. Rev. B, 24:2613–2626, 1981.
- [6] Warren J. Ewens. Mathematical Population Genetics. Springer-Verlag, New York, 1979.
- [7] Walter Fontana and Peter Schuster. Continuity in evolution: On the nature of transitions. Nature, 280:1451–1455, 1998.
- [8] H. I. Freedman. Deterministic Mathematical Models in Population Ecology. Marcel Dekker, New York, 1980.
- [9] J. S. Gale. Theoretical Population Genetics. Unwin Hyman, London, 1990.
- [10] John H. Gillespie. The Causes of Molecular Evolution. Oxford University Press, 1991.
- [11] Thomas G. Hallam and Simon A. Levin, editors. Mathematical Ecology. An Introduction. Springer-Verlag, New York, 1986.
- [12] W. Daniel Hillis. Co-evolution parasites improve simulated evolution as an optimization procedure. In Christopher G. Langton, Charles Taylor, J. Doyne Farmer, and Steen Rasmussen, editors, Artificial Life II, pages 313–324. Addison-Wesley, 1991.
- <span id="page-16-0"></span>[13] Martijn A. Huynen, Peter F. Stadler, and Walter Fontana. Smoothness within ruggedness: The role of neutrality in adaption. *Proc. Natl. Acad. Sci. USA*, 93:397-401, 1996.
- [14] Stuart A. Kauffman. The Origins of Order. Oxford University Press, Oxford, 1992.
- [15] Stuart A. Kauffman and Sonke Johnsen. Coevolution to the edge of chaos: Coupled fitness landscapes, poised states, and coevolutionary avalanches. J. theor. Biol., 149:467–505, 1991.
- [16] Stuart A. Kauffman and Edward D. Weinberger. The NK model of rugged fitness landscapes and its application to maturation of the immune response. J. theor. Biol., 141:211–245, 1989.
- [17] Motoo Kimura. The neutral theory of molecular evolution. Cambridge Univ. Press, 1984.
- [18] Martin Kreitman. Nucleotide polymorphism at the alcohol dehyrogenase locus of Drosophila melanogaster. Nature, 304:412–417, 1983.
- [19] Martin Kreitman and Hiroshi Akashi. Molecular evidence for natural selection. Annu. Rev. Ecol. Syst., 26:403–422, 1995.
- [20] M. E. J. Newman and Robin Engelhardt. Effects of selective neutrality on the evolution of molecular species. Proc. R. Soc. London B, 265:1333–1338, 1998.
- [21] Christian Reidys, Peter F. Stadler, and Peter Schuster. Generic properties of combinatory maps. Neutral networks of RNA secondary structure. Bull. Math. Biol., 59:339–397, 1997. Also published as: Preprint No. 95-07-058, Santa Fe Institute, Santa Fe, NM 1995.
- [22] Craig W. Reynolds. Competition, coevolution and the game of tag. In Rodney A. Brooks and Pattie Maes, editors, Artificial Life IV, pages 59–69. MIT Press, 1994.
- [23] M. Rietti-Shati, A. Shemesh, and W. Karlen. A 3000-year climatic record from biogenic silica oxygen isotopes in an equatorial high-altitude lake. Science, 281:980– 982, 1998.
- [24] Bonnie Sampsell. Isolation and genetic characterization of alcohol dehydrogenase thermostability variants occuring in natural populations of Drosophila melanogaster. Biochemical Genetics, 15:971–988, 1977.
- [25] Dietrich Stauffer and Amnon Aharony. Introduction to Percolation Theory. Taylor & Francis, Basingstoke Hauts, UK, 2nd edition, 1992.
- [26] A. V. Wilks, J. B. Gibson, J. G. Oakeshott, and G. K. Chambers. An electrophoretically cryptic alcohol dehydrogenase variant in Drosophila melanogaster. II. Postelectrophoresis heat-treatment screening of natural populations. Aust. J. Biol. Sci., 33:575–585, 1980.
- [27] S. Wright. The roles of mutation, inbreeding, crossbreeding, and selection in evolution. In Proceedings of the Sixth International Congress on Genetics, volume 1, pages 355– 366, 1932.