

## New Physics Discovery Potential in Future Experiments

S.I. Bitjukov<sup>1,2</sup>, N.V. Krasnikov<sup>3</sup>  
 Institute for Nuclear Research, Moscow, Russia

**Abstract**

We propose a method to estimate the probability of new physics discovery in future high energy physics experiments. Physics simulation gives both the average numbers  $\langle N_b \rangle$  of background and  $\langle N_s \rangle$  of signal events. We find that the proper definition of the significance for  $\langle N_b \rangle, \langle N_s \rangle \gg 1$  is  $S_{12} = \sqrt{\langle N_s \rangle + \langle N_b \rangle} - \sqrt{\langle N_b \rangle}$  in comparison with often used significances  $S_1 = \frac{\langle N_s \rangle}{\sqrt{\langle N_b \rangle}}$  and  $S_2 = \frac{\langle N_s \rangle}{\sqrt{\langle N_s \rangle + \langle N_b \rangle}}$ . We propose a method for taking into account the systematical errors related to nonexact knowledge of background and signal cross sections. An account of such systematics is very essential in the search for supersymmetry at LHC. We also propose a method for estimation of exclusion limits on new physics in future experiments.

---

<sup>1</sup>Institute for High Energy Physics, Protvino, Moscow region, Russia

<sup>2</sup>E-mails: bitjukov@mx.ihep.su, Serguei.Bitoukov@cern.ch

<sup>3</sup>E-mails: krasniko@ms2.inr.ac.ru, Nikolai.Krasnikov@cern.ch

# 1 Introduction

One of the common goals in the forthcoming experiments is the search for new phenomena. In the forthcoming high energy physics experiments (LHC, TEV22, NLC, ...) the main goal is the search for physics beyond the Standard Model (supersymmetry,  $Z'$ -,  $W'$ -bosons, ...) and the Higgs boson discovery as a final confirmation of the Standard Model. In estimation of the discovery potential of the future experiments (to be specific in this paper we shall use as an example CMS experiment at LHC [1]) the background cross section is calculated and for the given integrated luminosity  $L$  the average number of background events is  $\langle N_b \rangle = \sigma_b \cdot L$ . Suppose the existence of a new physics leads to the nonzero signal cross section  $\sigma_s$  with the same signature as for the background cross section that results in the prediction of the additional average number of signal events  $\langle N_s \rangle = \sigma_s \cdot L$  for the integrated luminosity  $L$ .

The total average number of the events is  $\langle N_{ev} \rangle = \langle N_s \rangle + \langle N_b \rangle = (\sigma_s + \sigma_b) \cdot L$ . So, as a result of new physics existence, we expect an excess of the average number of events. In real experiments the probability of the realization of  $n$  events is described by Poisson distribution [2]

$$f(n, \langle n \rangle) = \frac{\langle n \rangle^n}{n!} e^{-\langle n \rangle}. \quad (1)$$

Here  $\langle n \rangle$  is the average number of events.

Remember that the Poisson distribution  $f(n, \langle n \rangle)$  gives [3] the probability of finding exactly  $n$  events in the given interval of (e.g. space and time) when the events occur independently of one another and of  $x$  at an average rate of  $\langle n \rangle$  per the given interval. For the Poisson distribution the variance  $\sigma^2$  equals to  $\langle n \rangle$ . So, to estimate the probability of the new physics discovery we have to compare the Poisson statistics with  $\langle n \rangle = \langle N_b \rangle$  and  $\langle n \rangle = \langle N_b \rangle + \langle N_s \rangle$ . Usually, high energy physicists use the following “significances” for testing the possibility to discover new physics in an experiment:

- (a) “significance”  $S_1 = \frac{N_s}{\sqrt{N_b}}$  [4],
- (b) “significance”  $S_2 = \frac{N_s}{\sqrt{N_s + N_b}}$  [5, 6].

A conventional claim is that for  $S_1$  ( $S_2$ )  $\geq 5$  we shall discover new physics (here, of course, the systematical errors are ignored). For  $N_b \gg N_s$  the significances  $S_1$  and  $S_2$  coincide (the search for Higgs boson through the  $H \rightarrow \gamma\gamma$  signature). For

the case when  $N_s \sim N_b$ ,  $S_1$  and  $S_2$  differ. Therefore, a natural question arises: what is the correct definition for the significance  $S_1$ ,  $S_2$  or anything else ?

It should be noted that there is a crucial difference between “future” experiment and the “real” experiment. In the “real” experiment the total number of events  $N_{ev}$  is a given number (already has been measured) and we compare it with  $\langle N_b \rangle$  when we test the validity of the standard physics. So, the number of possible signal events is determined as  $N_s = N_{ev} - \langle N_b \rangle$  and it is compared with the average number of background events  $\langle N_b \rangle$ . The fluctuation of the background is  $\sigma_{fb} = \sqrt{\langle N_b \rangle}$ , therefore, we come to the  $S_1$  significance as the measure of the distinction from the standard physics. In the conditions of the “future” experiment when we want to search for new physics, we know only the average number of the background events and the average number of the signal events, so we have to compare the Poisson distributions  $P(n, \langle N_b \rangle)$  and  $P(n, \langle N_b \rangle + \langle N_s \rangle)$  to determine the probability to find new physics in the future experiment.

In this paper we estimate the probability to discover new physics in future experiments. We show that for  $\langle N_s \rangle, \langle N_b \rangle \gg 1$  the proper determination of the significance is  $S = \sqrt{\langle N_b \rangle + \langle N_s \rangle} - \sqrt{\langle N_b \rangle}$ . We suggest a method which takes into account systematic errors related to nonexact knowledge of the signal and background cross sections. We also propose a method for the estimation of exclusion limits on new physics in future experiments. Some of presented results has been published in our early paper [8].

The organization of the paper is the following. In the next section we give a method for the determination of the probability to find new physics in the future experiment and calculate the probability to discover new physics for the given  $(\langle N_b \rangle, \langle N_s \rangle)$  numbers of background and signal events under the assumption that there are no systematic errors. In section 3 we estimate the influence of the systematics related to the nonexact knowledge of the signal and background cross sections on the probability to discover new physics in future experiments. In Section 4 we describe a method for the estimation of exclusion limits on new physics in future experiments. In section 5 we estimate the probability of new physics discovery in future experiments. Section 6 contains the concluding remarks.

## 2 An analysis of statistical fluctuations

Suppose that for some future experiment we know the average number of the background and signal event  $\langle N_b \rangle, \langle N_s \rangle$ . As it has been mentioned in the Introduction, the probability of realization of  $n$  events in an experiment is given by the Poisson distribution

$$P(n, \langle n \rangle) = \frac{\langle n \rangle^n}{n!} e^{-\langle n \rangle}, \quad (2)$$

where  $\langle n \rangle = \langle N_b \rangle$  for the case of the absence of new physics and  $\langle n \rangle = \langle N_b \rangle + \langle N_s \rangle$  for the case when new physics exists. So, to determine the probability to discover new physics in future experiment, we have to compare the Poisson distributions with  $\langle n \rangle = \langle N_b \rangle$  (standard physics) and  $\langle n \rangle = \langle N_b \rangle + \langle N_s \rangle$  (new physics).

Consider, at first, the case when  $\langle N_b \rangle \gg 1$ ,  $\langle N_s \rangle \gg 1$ . In this case the Poisson distributions approach the Gaussian distributions

$$P_G(n, \mu, \sigma^2) = \frac{1}{\sigma\sqrt{2\pi}} \cdot e^{-\frac{(n-\mu)^2}{2\sigma^2}}, \quad (3)$$

with  $\mu = \sigma^2$  and  $\mu = \langle N_b \rangle$  or  $\mu = \langle N_b \rangle + \langle N_s \rangle$ . Here  $n$  is a real number. Note that for the Poisson distribution the mean equals to the variance.

The Gaussian distribution describes the probability density to realize  $n$  events in the future experiment provided the average number of events  $\langle n \rangle$  is a given number. In Fig.1 we show two Gaussian distributions  $P_G$  with  $\langle n \rangle = \langle N_b \rangle = 53$  and  $\langle n \rangle = \langle N_b \rangle + \langle N_s \rangle = 104$  ([6], Table.13, cut 6). As is clear from Fig.1 the common area for these two curves (the first curve shows the “standard physics” events distribution and the second one gives the “new physics” events distribution) is the probability that “new physics” can be described by the “standard physics”. In other words, suppose we know for sure that new physics takes place and the probability density of the events realization is described by curve II ( $f_2(x) = P_G(x, \langle N_b \rangle + \langle N_s \rangle, \langle N_b \rangle + \langle N_s \rangle)$ ). The probability  $\kappa$  that the “standard physics” (curve I ( $f_1(x) = P_G(x, \langle N_b \rangle, \langle N_b \rangle)$ )) can imitate new physics (i.e. the probability that we measure “new physics” but we think that it is described by the “standard physics”) is described by common area of curve I and II.

Numerically, we find that

$$\begin{aligned} \kappa &= \frac{1}{\sqrt{2\pi}\sigma_2} \int_{-\infty}^{\sigma_1\sigma_2} \exp\left[-\frac{(x-\sigma_2^2)^2}{2\sigma_2^2}\right] dx + \frac{1}{\sqrt{2\pi}\sigma_1} \int_{\sigma_1\sigma_2}^{\infty} \exp\left[-\frac{(x-\sigma_1^2)^2}{2\sigma_1^2}\right] dx \\ &= \frac{1}{\sqrt{2\pi}} \left[ \int_{-\infty}^{\sigma_1-\sigma_2} \exp\left[-\frac{y^2}{2}\right] dy + \int_{\sigma_2-\sigma_1}^{\infty} \exp\left[-\frac{y^2}{2}\right] dy \right] \\ &= 1 - \operatorname{erf}\left(\frac{\sigma_2 - \sigma_1}{\sqrt{2}}\right). \end{aligned} \quad (4)$$

Here  $\sigma_1 = \sqrt{N_b}$  and  $\sigma_2 = \sqrt{N_b + N_s}$ . The transformation of the distributions to standard normal distribution and the exploitation of the equality

$$\frac{x_0 - \sigma_1^2}{\sigma_1} = -\frac{x_0 - \sigma_2^2}{\sigma_2}$$

allows one to find the point  $x_0$  of the intersection of the curves I and II.

Let us discuss the meaning of our definition (4). For  $x \leq x_0 = \sigma_1\sigma_2$  we have  $f_1(x) \geq f_2(x)$ , i.e. the probability density of the standard physics realization is higher than the probability density of new physics realization. Therefore for  $x \leq x_0$  we do not have any indication in favour of new physics. The probability that the number of events is less than  $x_0$  is  $\alpha = \int_{-\infty}^{x_0} f_2(x)dx$ . For  $x > x_0$   $f_2(x) > f_1(x)$  that gives evidence in favour of new physics existence. However the probability of the background events with  $x > x_0$  is different from zero and is equal to  $\beta = \int_{x_0}^{\infty} f_1(x)dx$ . So we have two types of the errors. For  $x \leq x_0$  we do not have any evidence in favour of new physics (even in this case the probability of new physics realization is different from zero). For  $x > x_0$  we have evidence in favour of new physics. However for  $x > x_0$  the fluctuations of the background can imitate new physics. So the probability that standard physics can imitate new physics has two components  $\alpha$  and  $\beta$  and it is equal to  $\kappa = \alpha + \beta$ . If  $\kappa$  equals to 1 the new physics will never to be found in the experiment, if  $\kappa$  equals to 0 the first measurement with probability one has to answer to the question about presence or absence of the new physics (this case is impossible for two Poisson distributions). Note that the other estimator (normalized  $\kappa$ )  $\tilde{\kappa} = \frac{\kappa}{2 - \kappa}$  has the same properties. It means that the area of intersection of the probability density functions of the pure background and the background plus signal is the measure of the future experiment undiscovery potential.

As follows from formula (4) the role of the significance  $S$  plays

$$S_{12} = \sigma_2 - \sigma_1 = \sqrt{N_b + N_s} - \sqrt{N_b}. \quad (5)$$

Note that in refs.[7] the following criterion of the signal discovery has been used. The signal was assumed to be observable if  $(1 - \epsilon) \cdot 100\%$  upper confidence level for the background event rate is equal to  $(1 - \epsilon) \cdot 100\%$  lower confidence level for background plus signal ( $\epsilon = 0.01 - 0.05$ ). The corresponding significance is similar to our significance  $S_{12}$ . The difference is that in our approach the probability density  $\kappa$  that new physics is described by standard physics is equal to  $2\epsilon$ .

It means that for  $S_{12} = 1, 2, 3, 4, 5, 6$  the probability  $\kappa$  is correspondingly  $\kappa = 0.31, 0.046, 0.0027, 6.3 \cdot 10^{-5}, 5.7 \cdot 10^{-7}, 2.0 \cdot 10^{-7}$  in accordance with a general picture. As it has been mentioned in the Introduction two definitions of the significance are mainly used in the literature:  $S_1 = \frac{N_s}{\sqrt{N_b}}$  [4] and  $S_2 =$

$\frac{N_s}{\sqrt{N_s + N_b}}$  [5]. The significance  $S_{12}$  is expressed in terms of the significances  $S_1$  and  $S_2$  as  $S_{12} = \frac{S_1 S_2}{S_1 + S_2}$ .

For  $N_b \gg N_s$  (the search for Higgs boson through  $H \rightarrow \gamma\gamma$  decay mode) we find that

$$S_{12} \approx 0.5 \quad S_1 \approx 0.5 \quad S_2. \quad (6)$$

It means that for  $S_1 = 5$  (according to a common convention the  $5\sigma$  confidence level means a new physics discovery) the real significance is  $S_{12} = 2.5$ , that corresponds to  $\kappa = 1.24\%$ .

For the case  $N_s = kN_b$ ,  $S_{12} = k_{12}S_2$ , where for  $k = 0.5, 1, 4, 10$  the value of  $k_{12}$  is  $k_{12} = 0.55, 0.59, 0.69, 0.77$ . For not too high values of  $\langle N_b \rangle$  and  $\langle N_b + N_s \rangle$ , we have to compare the Poisson distributions directly. Again for the Poisson distribution  $P(n, \langle n \rangle)$  with the area of definition for nonnegative integers we can define  $P(x, \langle n \rangle)$  for real  $x$  as

$$\tilde{P}(x, \langle n \rangle) = \begin{cases} 0, & x \leq 0, \\ P([x], \langle n \rangle), & x > 0. \end{cases} \quad (7)$$

It is evident that

$$\int_{-\infty}^{\infty} \tilde{P}(x, \langle n \rangle) dx = 1. \quad (8)$$

So, the generalization of the previous determination of  $\kappa$  in our case is straightforward, namely,  $\kappa$  is nothing but the common area of the curves described by  $\tilde{P}(x, \langle N_b \rangle)$  (curve I) and  $\tilde{P}(x, \langle N_b \rangle + \langle N_s \rangle)$  (curve II) (see, Fig.2).

One can find that <sup>1</sup>

$$\begin{aligned} \kappa &= \alpha + \beta, \\ \alpha &= \sum_{n=0}^{n_0} \frac{(\langle N_b \rangle + \langle N_s \rangle)^n}{n!} e^{-(\langle N_b \rangle + \langle N_s \rangle)} = \\ &= \frac{\Gamma(n_0 + 1, \langle N_s \rangle + \langle N_b \rangle)}{\Gamma(n_0 + 1)}, \\ \beta &= \sum_{n=n_0+1}^{\infty} \frac{(\langle N_b \rangle)^n}{n!} e^{-\langle N_b \rangle} = 1 - \frac{\Gamma(n_0 + 1, \langle N_b \rangle)}{\Gamma(n_0 + 1)}, \\ n_0 &= \left[ \frac{\langle N_s \rangle}{\ln(1 + \frac{\langle N_s \rangle}{\langle N_b \rangle})} \right]. \end{aligned}$$

Numerical results are presented in Tables 1-6.

---

<sup>1</sup>We are indebted to Igor Semeniouk for the help in the derivation of these formulae

As it follows from these Tables for finite values of  $\langle N_s \rangle$  and  $\langle N_b \rangle$  the deviation from asymptotic formula (4) is essential. For instance, for  $N_s = 5$ ,  $N_b = 1$  ( $S_1 = 5$ )  $\kappa = 14.2\%$ . For  $N_s = N_b = 25$  ( $S_1 = 5$ )  $\kappa = 3.8\%$ , whereas asymptotically for  $N_s \gg 1$  we find  $\kappa = 1.24\%$ . Similar situation takes place for  $N_s \sim N_b$ .

### 3 An account of systematic errors related to nonexact knowledge of background and signal cross sections

In the previous section we determined the statistical error  $\kappa$  (the probability that “new physics” is described by “standard physics”). In this section we investigate the influence of the systematical errors related to a nonexact knowledge of the background and signal cross sections on the probability  $\kappa$  not to confuse a new physics with the old one.

Denote the Born background and signal cross sections as  $\sigma_b^0$  and  $\sigma_s^0$ . An account of one loop corrections leads to  $\sigma_b^0 \rightarrow \sigma_b^0(1 + \delta_{1b})$  and  $\sigma_s^0 \rightarrow \sigma_s^0(1 + \delta_{1s})$ , where typically  $\delta_{1b}$  and  $\delta_{1s}$  are  $O(0.5)$ .

Two loop corrections at present are not known. So, we can assume that the uncertainty related with nonexact knowledge of cross sections is around  $\delta_{1b}$  and  $\delta_{1s}$  correspondingly. In other words, we assume that the exact cross sections lie in the intervals  $(\sigma_b^0, \sigma_b^0(1 + 2\delta_{1b}))$  and  $(\sigma_s^0, \sigma_s^0(1 + 2\delta_{1s}))$ . The average number of background and signal events lie in the intervals

$$(\langle N_b^0 \rangle, \langle N_b^0 \rangle (1 + 2\delta_{1b})) \quad (9)$$

and

$$(\langle N_s^0 \rangle, \langle N_s^0 \rangle (1 + 2\delta_{1s})), \quad (10)$$

where  $\langle N_b^0 \rangle = \sigma_b^0 \cdot L$ ,  $\langle N_s^0 \rangle = \sigma_s^0 \cdot L$ .

To determine the probability that the new physics is described by the old one, we again have to compare two Poisson distributions with and without new physics but in distinction from Section 2 we have to compare the Poisson distributions in which the average numbers lie in some intervals. So, a priori the only thing we know is that the average numbers of background and signal events lie in the intervals (9) and (10), but we do not know the exact values of  $\langle N_b \rangle$  and  $\langle N_s \rangle$ . To determine the probability that the new physics is described by the old, consider

the worst case when we think that new physics is described by the minimal number of average events

$$\langle N_b^{min} \rangle = \langle N_b^0 \rangle + \langle N_s^0 \rangle. \quad (11)$$

Due to the fact that we do not know the exact value of the background cross section, consider the worst case when the average number of background events is equal to  $\langle N_b^0 \rangle (1 + 2\delta_{1b})$ . So, we have to compare the Poisson distributions with  $\langle n \rangle = \langle N_b^0 \rangle + \langle N_s^0 \rangle =$

$\langle N_b^0 \rangle (1 + 2\delta_{1b}) + (\langle N_s^0 \rangle - 2\delta_{1b} \langle N_b^0 \rangle)$  and  $\langle n \rangle = \langle N_b^0 \rangle (1 + 2\delta_{1b})$ . Using the result of the previous Section, we find that for case  $\langle N_b^0 \rangle \gg 1$ ,  $\langle N_s^0 \rangle \gg 1$  the effective significance is

$$S_{12s} = \sqrt{\langle N_b^0 \rangle + \langle N_s^0 \rangle} - \sqrt{\langle N_b^0 \rangle (1 + 2\delta_{1b})}. \quad (12)$$

For the limiting case  $\delta_{1b} \rightarrow 0$ , we reproduce formula (5). For not too high values of  $\langle N_b^0 \rangle$  and  $\langle N_s^0 \rangle$ , we have to use the results of the previous section (Tables 1-6).

As an example consider the case when  $\delta_{1b} = 0.5$ ,  $\langle N_s \rangle = 100$ ,  $\langle N_b \rangle = 50$  (typical situation for sleptons search). In this case we find that

$$\begin{aligned} S_1 &= \frac{\langle N_s \rangle}{\sqrt{\langle N_b \rangle}} = 14.1, \\ S_2 &= \frac{\langle N_s \rangle}{\sqrt{\langle N_s \rangle + \langle N_b \rangle}} = 8.2 \\ S_{12} &= \sqrt{\langle N_b \rangle + \langle N_s \rangle} - \sqrt{\langle N_b \rangle} = 5.2, \\ S_{12s} &= \sqrt{\langle N_b \rangle + \langle N_s \rangle} - \sqrt{2 \langle N_b \rangle} = 2.25. \end{aligned}$$

The difference between CMS adopted significance  $S_2 = 8.2$  (that corresponds to the probability  $\kappa = 0.24 \cdot 10^{-15}$ ) and the significance  $S_{12s} = 2.25$  taking into account systematics related to nonexact knowledge of background cross section is factor 3.6 The direct comparison of the Poisson distributions with  $\langle N_b \rangle (1 + 2\delta_{1b}) = 100$  and  $\langle N_b \rangle (1 + 2\delta_{1b}) + \langle N_{s,eff} \rangle$  ( $\langle N_{s,eff} \rangle = \langle N_s \rangle - 2\delta_{1b} \langle N_b \rangle = 50$ ) gives  $\kappa_s = 0.0245$ .

Another example is with  $\langle N_s \rangle = 28$ ,  $\langle N_b \rangle = 8$  and  $\delta_{1b} = 0.5$ . For such example we have  $S_1 = 9.9$ ,  $S_2 = 4.7$ ,  $S_{12} = 3.2$ ,  $S_{12s} = 2.0$ ,  $\kappa_s = 0.045$ .

So, we see that an account of the systematics related to nonexact knowledge of background cross sections is very essential and it decreases the LHC SUSY discovery potential.



## 4 Estimation of exclusion limits on new physics

In this section we generalize the results of the previous sections to obtain exclusion limits on signal cross section (new physics).

Suppose we know the background cross section  $\sigma_b$  and we want to obtain bound on signal cross section  $\sigma_s$  which depends on some parameters (masses of new particles, coupling constants, ...) and describes some new physics beyond standard model. Again as in Section 2 we have to compare two Poisson distributions with and without new physics. The results of Section 2 are trivially generalized for the case of the estimation of exclusion limits on signal cross section and, hence, on parameters (masses, coupling constants, ...) of new physics.

Consider at first the case when  $\langle N_b \rangle = \sigma_b \cdot L \gg 1$ ,  $\langle N_s \rangle = \sigma_s \cdot L \gg 1$  and the Poisson distributions approach the Gaussian distributions. As it has been mentioned in Section 2 the common area of the Gaussian curves with background events and with background plus signal events is the probability that "new physics" can be described by the "standard physics". For instance, when we require the probability that "new physics" can be described by the "standard physics" is less or equal 10% ( $S_{12}$  in formula (5) is larger than 1.64) it means that the formula

$$\sqrt{\langle N_b \rangle + \langle N_s \rangle} - \sqrt{\langle N_b \rangle} \leq 1.64 \quad (13)$$

gives us 90% exclusion limit on the average number of signal events  $\langle N_s \rangle$ . In general case when we require the probability that "new physics" can be described by the "standard physics" is more or less  $\epsilon$  the formula

$$\sqrt{\langle N_b \rangle + \langle N_s \rangle} - \sqrt{\langle N_b \rangle} \leq S(\epsilon) \quad (14)$$

allows us to obtain  $1 - \epsilon$  exclusion limit on signal cross section. Here  $S(\epsilon)$  is determined by the formula (4)<sup>2</sup>, i.e. we suppose that  $\epsilon = \kappa$ . It should be stressed that in fact the requirement that "new physics" with the probability more or equal to  $\epsilon$  can be described by the "standard physics" is our definition of the exclusion limit at  $(1 - \epsilon)$  probability for signal cross section. From the formula (14) we find that

$$\sigma_s \leq \frac{S^2(\epsilon)}{L} + 2S(\epsilon)\sqrt{\frac{\sigma_b}{L}}. \quad (15)$$

For the case of not large values of  $\langle N_b \rangle$  and  $\langle N_s \rangle$  we have to compare the Poisson distributions directly and the corresponding method has been formulated

---

<sup>2</sup>Note that  $S(1\%) = 2.57$ ,  $S(2\%) = 2.33$ ,  $S(5\%) = 1.96$  and  $S(10\%) = 1.64$

in Section 2. As an example in Table 7 we give 90% exclusion limits on the signal cross section for  $L = 10^4 pb^{-1}$  and for different values of background cross sections.

Formulae (14), (15) do not take into account the influence of the systematic errors related to nonexact knowledge of the background cross sections on the exclusion limits for signal cross section. To take into account such systematics we have to use the results of Section 3. The corresponding generalization of the formulae (14) and (15) is straightforward, namely:

$$\sqrt{\langle N_b \rangle + \langle N_s \rangle} - \sqrt{\langle N_b \rangle (1 + 2\delta_{1b})} \leq S(\epsilon), \quad (16)$$

$$\sigma_s \leq \frac{S^2(\epsilon)}{L} + 2S(\epsilon)\sqrt{\frac{\sigma_b(1 + 2\delta_{1b})}{L}} + 2\delta_{1b}\sigma_b. \quad (17)$$

Remember that  $\delta_{1b}$  describes theoretical uncertainty in the calculation of the background cross section. As an example, in Table 8 we give 90% exclusion limits on the signal cross section for  $L = 10^4 pb^{-1}$ ,  $2\delta_{1b} = 0.25$  and for different values of background cross sections.

Note that in refs.[9, 10] different and strictly speaking "ad hoc" methods to derive exclusion limits in future experiments has been suggested. As is seen from Fig.3 the essential differences in values of the exclusion limits take place. Let us compare these methods by the use of the equal probability test [11].

In order to estimate the various approaches of the exclusion limit determination we suppose that the new physics exists, i.e. the value  $\langle N_s \rangle$  equals to one of the exclusion limits from Fig.3 and the value  $\langle N_b \rangle$  equals to the corresponding value of expected background. Then we apply the equal probability test to find critical value  $n_0$  for hypotheses testing in future measurements. Here a zero hypothesis is the statement that new physics exists and an alternative hypothesis is the statement that new physics is absent. After calculation of the Type I error  $\alpha$  (the probability that the number of observed events will be equal to or less than the critical value  $n_0$ ) and the Type II error  $\beta$  (the probability that in the case of absence of new physics the number of observed events will be greater than the critical value  $n_0$ ) we can compare the methods. In Table 9 the comparison result is shown. As is seen from this Table the "Typical experiment" approach [10] gives too small values of exclusion limit. The difference in the 90% CL definition is the main reason of the difference between our result and the exclusion limit from ref. [9]. In ref [9] is used the criterion for determination exclusion limits:  $\beta < \delta$  and  $\frac{\alpha}{1 - \beta} < \epsilon$ , where the experiment will observe with probability at least  $1 - \delta$  at most a number of events such that the limit obtained at the  $1 - \epsilon$  confidence level

excludes the corresponding signal. If we define  $\epsilon$  as normalized  $\kappa$  ( $\epsilon = \tilde{\kappa} = \frac{\kappa}{2 - \kappa}$ ) we have the result close to ref. [9], i.e.  $\kappa = 0.17$  corresponds to  $\epsilon = 0.0929$ .

## 5 The probability of new physics discovery

In section 2 we determined the probability  $\kappa$  that "new physics" can be described by the "standard physics". But it is also very important to determine the probability of new physics discovery in future experiment. According to common definition [1] the new physics discovery corresponds to the case when the probability that background can imitate signal is less than  $5\sigma$  or in terms of the probability less than  $5.7 \cdot 10^{-7}$  (here of course we neglect any possible systematical errors).

So we require that the probability of the background fluctuations for  $n > n_0(\delta)$   $\beta(\delta)$  is less than  $\delta$ , namely

$$\beta(\delta) = \sum_{n=n_0(\delta)+1}^{\infty} P(< N_b >, n) \leq \delta \quad (18)$$

The probability  $1 - \alpha(\delta)$  that the number of signal events will be bigger than  $n_0(\delta)$  is equal to

$$1 - \alpha(\delta) = \sum_{n=n_0(\delta)+1}^{\infty} P(< N_b > + < N_s >, n) \quad (19)$$

It should be stressed that  $\delta$  is a given number and  $\alpha(\delta)$  is a function of  $\delta$ . Usually physicists claim the discovery of phenomenon [1] if the probability of the background fluctuation is less than  $5\sigma$  that corresponds to  $\delta_{dis} = 5.7 \cdot 10^{-7}$ . So from the equation (18) we find  $n_0(\delta)$  and estimate the probability  $1 - \alpha(\delta)$  that an experiment will satisfy the discovery criterion.

As an example consider the search for standard Higgs boson with a mass  $m_h = 110 \text{ GeV}$  at the CMS detector. For total luminosity  $L = 3 \cdot 10^4 \text{ pb}^{-1} (2 \cdot 10^4 \text{ pb}^{-1})$  one can find [1] that  $< N_b > = 2893(1929)$ ,  $< N_s > = 357(238)$ ,  $S_1 = \frac{N_s}{\sqrt{N_b}} = 6.6(5.4)$ .

Using the formulae (18, 19) for  $\delta_{dis} = 5.7 \cdot 10^{-7}$  ( $5\sigma$  discovery criterium) we find that  $1 - \alpha(\delta_{dis}) = 0.96(0.73)$ . It means that for total luminosity  $L = 3 \cdot 10^4 \text{ pb}^{-1} (2 \cdot 10^4 \text{ pb}^{-1})$  the CMS experiment will discover at  $\geq 5\sigma$  level standard Higgs boson with a mass  $m_h = 110 \text{ GeV}$  with a probability 96(73) percent.

## 6 Conclusions

In this paper we determined the probability to discover the new physics in the future experiments when the average number of background  $\langle N_b \rangle$  and signal events  $\langle N_s \rangle$  is known. We have found that in this case for  $\langle N_s \rangle \gg 1$  and  $\langle N_b \rangle \gg 1$  the role of significance plays

$$S_{12} = \sqrt{\langle N_b \rangle + \langle N_s \rangle} - \sqrt{\langle N_b \rangle}$$

in comparison with often used expressions for the significances  $S_1 = \frac{\langle N_s \rangle}{\sqrt{\langle N_b \rangle}}$

and  $S_2 = \frac{\langle N_s \rangle}{\sqrt{\langle N_s \rangle + \langle N_b \rangle}}$ .

For  $\langle N_s \rangle \ll \langle N_b \rangle$  we have found that  $S_{12} = 0.5S_1 = 0.5S_2$ . For not too high values of  $\langle N_s \rangle$  and  $\langle N_b \rangle$ , when the deviations from the Gaussian distributions are essential, our results are presented in Tables 1-6. We proposed a method for taking into account systematical errors related to the nonexact knowledge of background and signal events. An account of such kind of systematics is very essential in the search for supersymmetry and leads to an essential decrease in the probability to discover the new physics in the future experiments. We also proposed a method for the estimation of exclusion limits on new physics in future experiments.

We are indebted to M.Dittmar for very hot discussions and useful questions which were one of the motivations to perform this study. We are grateful to V.A.Matveev for the interest and useful comments.

## References

- [1] *The Compact Muon Solenoid*. Technical Proposal, CERN/LHCC 94 -38, 1994.
- [2] N.L.Johnson, S.Kotz, *Distributions Statistics – Discrete Distributions*, Wiley, New York, 1969.
- [3] Particle Data Group, Phys.Rev **D54** 1 (1996).
- [4] See as an example:  
V.Tisserand, *The Higgs to Two Photon Decay in the ATLAS Detector*, Talk given at the VI International Conference on Calorimetry in High Energy Physics, Frascati (Italy), June 8-14, 1996.  
S.I.Bitukov and N.V.Krasnikov, *The Search for New Physics by the Measurement of the Four-jet Cross Section at LHC and TEVATRON*, Modern Physics Letter **A12**(1997)2011, also hep-ph/9705338.

- M.Dittmar and H.Dreiner, *LHC Higgs Search with  $l^+\nu l^-\bar{\nu}$  final states*, CMS Note 97/083, October 1997.
- [5] See as an example:  
 D.Denegri, L.Rurua and N.Stepanov, *Detection of Sleptons in CMS, Mass Reach*, CMS Note CMS TN/96-059, October 1996.  
 F.Charles, *Inclusive Search for Light Gravitino with the CMS Detector*, CMS Note 97/079, September 1997.  
 S.Abdullin, *Search for SUSY at LHC: Discovery and Inclusive Studies*, Presented at International Europhysics Conference on High Energy Physics, Jerusalem, Israel, August 19-26, 1997, CMS Conference Report 97/019, November 1997.
- [6] S.I.Bityukov and N.V.Krasnikov, *The Search for Sleptons and Flavour Lepton Number Violation at LHC (CMS)*, Preprint IFVE 97-67, Protvino, 1997, also hep-ph/9712358.
- [7] N.Brown, *Degenerate Higgs and Z Boson at LEP200*, Z.Phys., **C49**, 1991, p.657.  
 H.Baer, M.Bisset, C.Kao and X.Tata, *Observability of  $\gamma\gamma$  decays of Higgs bosons from supersymmetry at hadron supercolliders*, Phys.Rev., **D46**, 1992, p.1067.
- [8] S.I.Bityukov and N.V.Krasnikov, *Towards the observation of signal over background in future experiments*, Preprint INR 0945a/98, Moscow, 1998, also physics/9808016.
- [9] J.J.Hernandez, S.Navas and P.Rebecchi, *Estimating exclusion limits in prospective studies of searches*, Nucl.Instr.&Meth. A **378**, 1996, p.301, also J.J.Hernandez and S.Navas, *JASP: a program to estimate discovery and exclusion limits in prospective studies of searches*, Comp.Phys.Comm. **100**, 1997 p.119.
- [10] T.Tabarelli de Fatis and A.Tonazzo, *Expectation values of exclusion limits in future experiments* (Comment), Nucl.Instr.&Meth. A **403**, 1998, p.151.
- [11] S.I.Bityukov and N.V.Krasnikov, *On observability of signal over background*, Preprint IFVE 98-48, Protvino, 1998, also physics/9809037.




Figure 1: The probability density functions  $f_{1,2}(x) \equiv P_G(x, \mu_{1,2}, \sigma^2)$  for  $\mu_1 = \langle N_b \rangle = 53$  and  $\mu_2 = \langle N_b \rangle + \langle N_s \rangle = 104$ .




Figure 2: The probability density functions  $f_{1,2}(x) \equiv \tilde{P}(x, \mu_{1,2})$  for  $\mu_1 = \langle N_b \rangle = 1$  and  $\mu_2 = \langle N_b \rangle + \langle N_s \rangle = 6$ .




Figure 3: Estimations of the 90% CL upper limit on the signal in a future experiment as a function of the expected background. The method proposed in ref. [10] gives the values of exclusion limit close to "Typical experiment" approach.



Table 1: The dependence of  $\kappa$  on  $\langle N_s \rangle$  and  $\langle N_b \rangle$  for  $S_1 = 5$

$\langle N_s \rangle$	$\langle N_b \rangle$	$\kappa$
5	1	0.1423
10	4	0.0828
15	9	0.0564
20	16	0.0448
25	25	0.0383
30	36	0.0333
35	49	0.0303
40	64	0.0278
45	81	0.0260
50	100	0.0245
55	121	0.0234
60	144	0.0224
65	169	0.0216
70	196	0.0209
75	225	0.0203
80	256	0.0198
85	289	0.0193
90	324	0.0189
95	361	0.0185
100	400	0.0182
150	900	0.0162
500	$10^4$	0.0136
5000	$10^6$	0.0125

Table 2: The dependence of  $\kappa$  on  $\langle N_s \rangle$  and  $\langle N_b \rangle$  for  $S_2 \approx 5$ .

$\langle N_s \rangle$	$\langle N_b \rangle$	$\kappa$
26	1	$0.15 \cdot 10^{-4}$
29	4	$0.14 \cdot 10^{-3}$
33	9	$0.44 \cdot 10^{-3}$
37	16	$0.99 \cdot 10^{-3}$
41	25	$0.17 \cdot 10^{-2}$
45	36	$0.26 \cdot 10^{-2}$
50	49	$0.31 \cdot 10^{-2}$
55	64	$0.36 \cdot 10^{-2}$
100	300	$0.74 \cdot 10^{-2}$
150	750	$0.89 \cdot 10^{-2}$

Table 3:  $\langle N_s \rangle = \frac{1}{5} \cdot \langle N_b \rangle$ . The dependence of  $\kappa$  on  $\langle N_s \rangle$  and  $\langle N_b \rangle$ .

$\langle N_s \rangle$	$\langle N_b \rangle$	$\kappa$
50	250	0.131
100	500	0.033
150	750	$0.89 \cdot 10^{-2}$
200	1000	$0.25 \cdot 10^{-2}$
250	1250	$0.74 \cdot 10^{-3}$
300	1500	$0.22 \cdot 10^{-3}$
350	1750	$0.65 \cdot 10^{-4}$
400	2000	$0.20 \cdot 10^{-4}$

Table 4:  $\langle N_s \rangle = \frac{1}{10} \cdot \langle N_b \rangle$ . The dependence of  $\kappa$  on  $\langle N_s \rangle$  and  $\langle N_b \rangle$ .

$\langle N_s \rangle$	$\langle N_b \rangle$	$\kappa$
50	500	0.275
100	1000	0.123
150	1500	0.059
200	2000	0.029
250	2500	0.015
300	3000	$0.75 \cdot 10^{-2}$
350	3500	$0.38 \cdot 10^{-2}$
400	4000	$0.20 \cdot 10^{-2}$
450	4500	$0.11 \cdot 10^{-2}$
500	5000	$0.56 \cdot 10^{-3}$

Table 5:  $\langle N_s \rangle = \langle N_b \rangle$ . The dependence of  $\kappa$  on  $\langle N_s \rangle$  and  $\langle N_b \rangle$ .

$\langle N_s \rangle$	$\langle N_b \rangle$	$\kappa$
2.	2.	0.561
4.	4.	0.406
6.	6.	0.308
8.	8.	0.239
10.	10.	0.188
12.	12.	0.150
14.	14.	0.121
16.	16.	0.098
18.	18.	0.079
20.	20.	0.064
24.	24.	0.042
28.	28.	0.028
32.	32.	0.019
36.	36.	0.013
40.	40.	$0.87 \cdot 10^{-2}$
50.	50.	$0.34 \cdot 10^{-2}$
60.	60.	$0.13 \cdot 10^{-2}$
70.	70.	$0.52 \cdot 10^{-3}$
80.	80.	$0.21 \cdot 10^{-3}$
100.	100.	$0.33 \cdot 10^{-4}$

Table 6:  $\langle N_s \rangle = 2 \cdot \langle N_b \rangle$ . The dependence of  $\kappa$  on  $\langle N_s \rangle$  and  $\langle N_b \rangle$ .

$\langle N_s \rangle$	$\langle N_b \rangle$	$\kappa$
2.	1.	0.463
4.	2.	0.294
6.	3.	0.200
8.	4.	0.141
10.	5.	0.102
12.	6.	0.073
14.	7.	0.052
16.	8.	0.037
18.	9.	0.027
20.	10.	0.020
24.	12.	0.011
28.	14.	$0.59 \cdot 10^{-2}$
32.	16.	$0.33 \cdot 10^{-2}$
36.	18.	$0.18 \cdot 10^{-2}$
40.	20.	$0.10 \cdot 10^{-2}$
50.	25.	$0.23 \cdot 10^{-3}$
60.	30.	$0.56 \cdot 10^{-4}$

Table 7: 90% exclusion limits on signal cross section for  $L = 10^4 pb^{-1}$  and for different background cross section (everything in pb). The third column gives exclusion limit according to formula (15).

$\sigma_b$	$\sigma_s$	$\sigma_s$ (continuous limit)
$10^3$	1.041	1.038
$10^2$	0.329	0.328
10	0.104	0.104
1	0.033	0.033
0.1	0.011	0.011
0.01	0.0036	0.0035
0.001	0.0013	0.0013
0.0001	0.00060	0.00060

Table 8: 90% exclusion limits on signal cross section for  $L = 10^4 pb^{-1}$ ,  $2\delta_{1b} = 0.25$  and for different background cross section (everything in pb). The third column gives exclusion limit according to formula (17).

$\sigma_b$	$\sigma_s$	$\sigma_s$ (continuous limit)
$10^3$	251.25	251.16
$10^2$	25.37	25.37
10	2.62	2.62
1	0.29	0.29
0.1	0.037	0.037
0.01	0.0064	0.0064
0.001	0.0017	0.0017
0.0001	0.00064	0.00066

Table 9: The comparison of the different approaches to determination of the exclusion limits. The  $\alpha$  and the  $\beta$  are the Type I and the Type II errors under the equal probability test. The  $\kappa$  equals to the sum of  $\alpha$  and  $\beta$ .

$N_b$	this paper				ref. [9]				ref. [10]			
	$N_s$	$\alpha$	$\beta$	$\kappa$	$N_s$	$\alpha$	$\beta$	$\kappa$	$N_s$	$\alpha$	$\beta$	$\kappa$
1	6.02	0.08	0.02	0.10	4.45	0.09	0.08	0.17	3.30	0.20	0.08	0.28
2	7.25	0.05	0.05	0.10	5.50	0.13	0.05	0.18	3.90	0.16	0.14	0.30
3	8.32	0.07	0.03	0.10	6.40	0.09	0.08	0.18	4.40	0.14	0.18	0.32
4	9.20	0.05	0.05	0.10	7.25	0.13	0.05	0.18	4.80	0.23	0.11	0.34
5	10.06	0.07	0.03	0.10	7.90	0.10	0.07	0.17	5.20	0.20	0.13	0.34
6	10.67	0.06	0.04	0.10	8.41	0.09	0.08	0.18	5.50	0.19	0.15	0.34
7	11.37	0.05	0.05	0.10	9.00	0.08	0.10	0.18	5.90	0.17	0.17	0.34
8	12.02	0.07	0.03	0.10	9.70	0.10	0.06	0.17	6.10	0.17	0.18	0.35
9	12.51	0.06	0.04	0.10	10.16	0.09	0.07	0.17	6.40	0.16	0.20	0.36
10	13.04	0.05	0.05	0.10	10.50	0.09	0.08	0.17	6.70	0.22	0.14	0.36
11	13.62	0.04	0.06	0.10	10.80	0.08	0.09	0.18	6.90	0.21	0.15	0.36