

**CLASSICAL TUNNELING
OF SOMMERFELD SPHERE
IN CLASSICAL ELECTRODYNAMICS**

Alexander A. Vlasov

High Energy and Quantum Theory
Department of Physics
Moscow State University
Moscow, 119899
Russia

One-dimensional motion of Sommerfeld sphere in the case of potential barrier is numerically investigated. The effect of classical tunneling is found out - Sommerfeld sphere overcomes the barrier and finds itself in the forbidden, from classical point of view, area

03.50.De

Here numerical investigations of one-dimensional motion of Sommerfeld sphere with total charge Q , mechanical mass m and radius a are continued (for prev. results - see [1,2]).

We consider one-dimensional motion for the case of potential barrier, produced by homogeneous static electric field E , stretched in z - direction for $0 < z < L$ (like in plane condenser):

$$E = \begin{cases} 0, & z < 0; \\ E, & 0 < z < L; \\ 0, & L < z; \end{cases}$$

For dimensionless variables $y = R/L$, $x = ct/L$, $a^* = 2a/L$ the equation of motion of Sommerfeld sphere for our problem is

$$\frac{d^2 y}{dx^2} = \left(1 - \left(\frac{dy}{dx}\right)^2\right)^{3/2} \left\{ k \cdot \left[-\int_{x^-}^{x^+} dz \frac{z - a^*}{L^2} + \ln \frac{L^+}{L^-} + \left(\frac{1}{\beta^2} - 1\right) \ln \frac{1 + \beta}{1 - \beta} \right] + \lambda \cdot F_Z \right\} \quad (1)$$

here

$$x^\pm = a^* \pm L^\pm, \quad L^\pm = y(x) - y(x - x^\pm), \quad L = y(x) - y(x - z),$$

$$\beta = dy/dx, \quad k = \frac{Q^2}{2mc^2 a}, \quad \lambda = \frac{LQE}{mc^2}$$

$$F_Z = \begin{cases} 0, & y < -a^*/2; \\ (2y/a^* + 1)/2, & -a^*/2 < y < a^*/2; \\ 1, & a^*/2 < y < 1 - a^*/2; \\ (-2y/a^* + 1 + 2/a^*)/2, & 1 - a^*/2 < y < 1 + a^*/2; \\ 0, & 1 + a^*/2 < y; \end{cases}$$

Later on we take $k = 1$.

It is useful to compare solutions of (1) with classical point charge motion in the same field, governed by the following relativistic equation without radiation force:

$$\frac{d^2y}{dx^2} = \left(1 - \left(\frac{dy}{dx}\right)^2\right)^{3/2} \lambda \cdot F_E \quad (2)$$

here

$$F_E = \lambda \begin{cases} 0, & y < 0; \\ 1, & 0 < y < 1; \\ 0, & 1 < y; \end{cases}$$

A.

Numerical calculations of eqs. (1-2) show that there is the effect of classical tunneling for Sommerfeld sphere.

Indeed, classical point particle motion, governed by eq. (2), is simple:

$$\gamma = \lambda + \gamma_0 \quad (3)$$

here $\gamma = 1/\sqrt{1 - (v/c)^2}$ - Lorentz factor, and γ_0 - Lorentz factor for initial velocity.

Thus for given initial velocity for $\lambda < 1 - \gamma_0$ there is the turning point - i.e. classical particle cannot overcome the potential barrier (for ex., for $(v/c)_0 = 0.3$ the crucial value of λ is approximately $-0.0482848\dots$) and the electric field turns the particle back.

For numerical calculations of eq.(1) we took $(v/c)_0 = 0.3$ and $\lambda = -0.04829$.

Numerical results for different values of a^* are shown on fig. (A.1-A.3) (vertical axis is velocity dy/dx , horizontal axis is coordinate y ; trajectories of Sommerfeld sphere are compared with trajectories of classical particle):

we see that for $a^* = 0.2$, (*Fig.A.1*) and for $a^* = 0.001$, (*Fig.A.2*) there is the effect of classical tunneling - Sommerfeld sphere overcomes the barrier and finds itself in the forbidden, from classical point of view, area with resulting velocities $(v/c)_r = 0.139630$ for $a^* = 0.2$ and $(v/c)_r = 0.037190$ for $a^* = 0.001$;

while for $a^* = 0.0002$, (*Fig.A.3*) - this effect is absent (the barrier is too high...).

These results confirm the results of the works [3,4] where the effects of "pumping" of extended object in the potential hole [4] (on the example of some model in classical field theory) and classical tunneling [3] for Lorentz-Dirac eq. were found.

We see that these unusual (from point of view of classical mechanics) results are not the consequences of "point-like" description of charged particles but reflect the general features of equations with retardation (difference-differential equations).

REFERENCES

1. Alexander A.Vlasov, physics/9901051.
2. Alexander A.Vlasov, physics/9902018.
3. W.Troost et al., preprint hep-th/9602066.
4. Alexander A.Vlasov, Theoretical and Mathematical Physics, 109, n.3, 1608(1996).




Fig.A.1




Fig.A.2




Fig.A.3