Proton structure effects in muonic hydrogen

Krzysztof Pachucki *

Institute of Theoretical Physics, Warsaw University, Hoża 69, 00-681 Warsaw, Poland.

Abstract

The proton structure effects, including finite size, polarizability and selfenergy is considered and their influence on energy levels of muonic hydrogen is recalculated. A new theoretical prediction for the Lamb shift is presented together with improved values of all known QED contributions.

PACS numbers 36.10 Dr, 12.20 Ds, 31.30 Jv

Typeset using REVT_{EX}

^{*}E-mail address: krp@fuw.edu.pl

The precision tests of quantum electrodynamics (QED) in atomic systems have reached the level, where nuclear structure effects become significant. Moreover, the lack of accurate data on low energy structure functions of the nucleus strongly limits theoretical predictions. The well known example is the hydrogen hyperfine structure splitting, where measurement is 6 orders of magnitude more precise than the current theoretical predictions. Other example is the Lamb shift in hydrogen, where inaccuracy in the proton charge radius dominate other theoretical uncertainties. Since many years it was the motivation to study the pure QED systems like muonium and positronium, where strong interaction effects are negligible or well estimated at the precision level of interest. A new possibilities in the improvement of QED test on bound systems appeared with the project of measurement of the Lamb shift in muonic hydrogen [1]. Being bound in the ground state, muon with its 200 heavier mass compared to the electron, penetrates the proton and become sensitive to the distribution of charge, magnetic moment or the polarizability. The main goal of the measurement of the Lamb shift in muonic hydrogen is a precise determination of mean square proton charge radius. It will verify older measurements of proton charge radius based on the elastic electronproton scattering. The results of these experiments had to be extrapolated to very low $q^2 = 0$ momentum transfer, and therefore might be not very reliable. In contrast the muonic Lamb shift measurement aims to improve the precision of the proton charge radius by at least 10 times. However it would require the calculation of all other contributions with the comparable or better precision. Few years ago, in the summary of all known up to date QED effects on muonic hydrogen energy levels [2], we concluded that three-loop vacuum polarization have limited the precision of theoretical predictions by 0.01 meV, and pointed attention to the proton polarizability effect, which was not calculated that time. The progress on the experimental side with the preparation of the measurement of muonic hydrogen Lamb shift has stimulated further theoretical works. The rather difficult threeloop vacuum polarization contribution has recently been calculated by Kinoshita and Nio [3] with the result

$$\Delta E(2P - 2S) = 0.0076 \text{ meV}.$$
 (1)

A new contribution coming from hadronic vacuum polarization has been studied by Friar and collaborators [4], who obtained the following result

$$\Delta E(2P - 2S) = 0.0113(3) \text{ meV}.$$
(2)

The proton polarizability correction has been analyzed by Rosenfelder in [5]. His estimate for this effect is

$$\Delta E(2P - 2S) = 0.017(4) \text{ meV}.$$
(3)

Another recent work [6] gives a similar result of ~ 0.018 meV. The calculation of proton polarizability effect is affected by the lack of precise data on low–energy proton structure functions. In this paper we present another estimate of this effect together with the complete review of all other proton structure contributions. We would like to emphasize the importance of the proton self–energy effect and the related problem with the meaning of mean square charge radius. In other words we present the relation of charge radius as obtained from atomic spectroscopy measurements with that based on the electron scattering data. It is well known, the shift of atomic energy levels due to finite charge distribution of atomic nucleus. It is given by the formula

$$E_{FS} = \frac{2}{3n^3} \alpha^4 \,\mu^3 \,\langle r^2 \rangle \,\delta_{l0} \,, \tag{4}$$

which gives -3.862(108) meV contribution to 2P–2S splitting, were we used r = 0.862(12) fm from [7]. From this one concludes, that the measurement of muonic hydrogen with the precision of 0.01 meV will lead to the tenfold improvement in the proton charge radius. This formula in (4) accounts for most of the proton structure effects. Any correction beyond that is much smaller and usually neglected for light (electronic) atoms. It is our aim to review them in the context of muonic hydrogen Lamb shift.

Since the leading contribution to the 2P-2S splitting comes from the electron vacuum polarization (e.v.p.), the second order corrections from combined finite size and e.v.p. effects are nonnegligible. They have been calculated in [8], here we present a little more accurate result

$$\Delta E(2P - 2S) = -r^2 \, 0.0282 = -0.0209(6) \text{ meV} \,. \tag{5}$$

Further corrections are due to pure photon exchange terms only. The $O(m \alpha^5)$ correction is given by two-photon scattering amplitude with external momenta on mass shell, see Eq. (7). It is the main contribution, which we analyze in this work. However, lets consider first the small corrections beyond this two-photon exchange approximation, namely that of order $m \alpha^6$. Since they are small, it is sufficient to calculate them in the external field approximation, or in other words in the limit of infinite nucleus mass. The energy levels are obtained then from the Dirac equation in the potential of the finite size nucleus. They have been considered in details by Friar in [9]. In the logarithmic approximation they are given by two terms

$$E = E_{FS} \alpha^2 \ln \alpha \left(\frac{2}{3} \mu^2 \langle r^2 \rangle - 1\right) \tag{6}$$

and contribute the amount of -0.0009 meV to the 2P-2S splitting and thus are almost negligible.

The correction given by the two–photon exchange is of the main interest. It is given by the following expression

$$E = -\frac{e^4}{2} \phi^2(0) \int \frac{d^4q}{(2\pi)^4 i} \frac{1}{q^4} \left[T^{\mu\nu} - t^{\mu\nu}(M) \right] t_{\mu\nu}(m)$$

= $-2 e^4 \phi^2(0) \frac{m}{M} \int \frac{d^4q}{(2\pi)^4 i} \frac{(T_2 - t_2)(q^2 - \nu^2) - (T_1 - t_1)(q^2 + 2\nu^2)}{q^4 (q^4 - 4m^2\nu^2)},$ (7)

where

$$T^{\mu\nu} = -i \int d^4 q \, e^{i \, q \, (x-x')} \, \langle P | T \, j^\mu(x) \, j^\nu(x') | P \rangle = -\left(g^{\mu\nu} - \frac{q^\mu \, q^\nu}{q^2}\right) \frac{T_1}{M} + \left(t^\mu - \frac{\nu}{q^2} \, q^\mu\right) \left(t^\nu - \frac{\nu}{q^2} \, q^\nu\right) \frac{T_2}{M} \,, \tag{8}$$

t = (1, 0, 0, 0), P = Mt is a proton momentum at rest and $\nu = q^0$. For a point–like proton $T^{\mu\nu} \equiv t^{\mu\nu}(M)$ and

$$t^{\mu\nu}(M) = \operatorname{Tr}\left[\gamma^{\mu} \frac{1}{\not p - M} \gamma^{\nu} \frac{\gamma^{0} + I}{4}\right] + (q \to -q), \qquad (9)$$

$$t_1 = -\frac{4 M^2 \nu^2}{q^4 - 4 M^2 \nu^2}, \qquad (10)$$

$$t_2 = \frac{4 M^2 q^2}{q^4 - 4 M^2 \nu^2}.$$
(11)

Since the amplitude $T^{\mu\nu}$ does not have any singularities at small q^2 the following holds:

$$T_2 = O(q^2) \,, \tag{12}$$

$$T_1 + \frac{\nu^2}{q^2} T_2 = O(q^2) \,. \tag{13}$$

The off-shell (spin averaged) forward Compton amplitude $T^{\mu\nu}$ of the proton is not directly measured. However, it could be expressed in terms of its imaginary part, through the dispersion relations. Before using them, one notices that the integral in Eq. (7) is infrared divergent. It requires subtraction of the leading finite size term, which has already been accounted for. It also requires an additional subtraction of the proton self-energy term. Since it changes the analytic behavior of T_1 and T_2 at small $p^2 - M^2$, see Eq. (20), this proton self-energy could be only partially accounted for in the proton formfactors, as a contribution to anomalous magnetic moment or the charge radius. Since, in our opinion this problem is not generally known, we repeat here the analysis from our former paper on radiative recoil corrections and correct some minor missprints.

If we assume a point-like proton the contribution of the proton self–energy to the Lamb shift of S-states is [10]

$$E = \frac{\alpha^5 \,\mu^3}{\pi \,n^3 \,M^2} \left[\left(\frac{10}{9} + \frac{4}{3} \,\ln \frac{M}{\mu \,\alpha^2} \right) - \frac{4}{3} \,\ln k_0(n) \right] \,. \tag{14}$$

For a *true* proton there is a finite size correction as given by Eq. (4). The problem is that the proton self-energy is modified by and modifies as well, the finite size effect. Therefore some corrections might be counted twice. To incorporate the correction (14) unambiguously we must precisely specify what is the nuclear mean square charge radius. Its usual definition through the Sachs formfactor

$$\frac{\langle r^2 \rangle}{6} = \frac{\partial G_E(q^2)}{\partial (q^2)} \Big|_{q^2 = 0} \tag{15}$$

is not correct at our precision level, because the radiative correction to G_E is infrared divergent or depend on spurious photon mass. Following [2] we propose thus a different definition using the forward scattering amplitude described by $T^{\mu\nu}$, or more precisely by its longitudinal component T_L defined by

$$T_L \equiv \left(1 - \frac{\nu^2}{q^2}\right) T_2 - T_1 \,. \tag{16}$$

For our purpose we consider a nonrelativistic limit $\nu \sim q^2$ and $p^2 - M^2 = (P+q)^2 - M^2 \approx 0$ of T_L . For a point-like particle without radiative corrections T_L is

$$T_L \approx M \operatorname{Tr} \left[\gamma^0 \frac{1}{\not p - M} \gamma^0 \frac{(\gamma^0 + I)}{4} \right] + (q \to -q) \approx \frac{2M^2}{p^2 - M^2} + (q \to -q) \,, \tag{17}$$

where p = P + q. With a finite size particle

$$\gamma^{\mu} \to \Gamma^{\mu} = \gamma^{\mu} F_1 + i \frac{\sigma^{\mu\nu}}{2M} q_{\nu} F_2 , \qquad (18)$$

 T_L acquires a correction

$$\Delta T_L \approx \frac{2M^2}{p^2 - M^2} [G_E^2(q^2) - 1] + (q \to -q) \approx \frac{2M^2}{p^2 - M^2} q^2 \frac{\langle r^2 \rangle_{bar}}{3} + (q \to -q), \qquad (19)$$

where $G_E = F_1 + \frac{q^2}{4M^2}F_2$ is an electric formfactor. The radiative corrections for a point-like particle [2] in the nonrelativistic limit are

$$\Delta T_L = \frac{\alpha}{\pi} \frac{q^2}{p^2 - M^2} \left(\frac{10}{9} + \frac{4}{3} \ln \frac{M^2}{M^2 - p^2} \right) + (q \to -q) \,. \tag{20}$$

We define $\langle r^2 \rangle$ by the following equation that describes the low-energy behavior of the forward scattering amplitude

$$T_L - t_L \approx \frac{q^2}{p^2 - M^2} \left(\frac{4\alpha}{3\pi} \ln \frac{M^2}{M^2 - p^2} + \frac{2}{3} M^2 \langle r^2 \rangle \right) + (q \to -q).$$
(21)

We expect that for any nucleus the logarithmic term above will be the same, since it is only related to the fact that nucleus has a charge, and does not depend on other details like the spin. The associated correction to the energy for S-states has the form

$$\Delta E = \frac{2}{3 n^3} \alpha^4 \mu^3 \langle r^2 \rangle + \frac{4 \alpha^5}{3 \pi n^3} \frac{\mu^3}{M^2} \left[\ln \left(\frac{M}{\mu \alpha^2} \right) - \ln k_0(n) \right].$$
(22)

The small second term in the above Eq. gives -0.0099 meV for 2S state in μ H. For P-states the proton self-energy contributes only through Bethe log. The correction to anomalous magnetic moment is already taken into account in the calculation of relativistic effects as given by the Breit Hamiltonian.

We now return to the forward scattering amplitude and its associated correction to energy of μ H. We can neglects here QED effects on the structure functions, since they are α times smaller. Otherwise dispersion relations for T_1 and T_2 as in Eq. (29,30) will not be correct. After neglection of QED effects one can expect a separated pole at $2M\nu = q^2$, that is due to the elastic contribution. It is obtained with the help of the proton elastic formfactors F_1 and F_2 , see Eq. (18). One derives for T_1 and T_2 the following expressions

$$T_1^B = -\frac{1}{(q^4 - 4M^2\nu^2)} \left(4M^2\nu^2F_1^2 + 2q^4F_1F_2 + q^4F_2^2\right)$$
(23)

$$T_2^B = \frac{1}{(q^4 - 4M^2\nu^2)} \left(4M^2 q^2 F_1^2 - q^4 F_2^2 \right).$$
(24)

Any contribution in the two photon exchange beyond this second order Born term is considered to be due to the proton polarizability. The correction to energy as given by Eq. (7) shows infrared singularity, which is due to the finite size effect, already accounted for at the lower order in α . One first subtracts this singularity and then integrate over q. With the commonly used dipole parametrization of proton formfactors and $\Lambda^2 = 0.71 \text{ GeV}^2$ one obtains $\Delta E = 0.018 \text{ meV}$. However, as pointed out in [1] this result is not correct, because this dipole parametrization corresponds to the proton radius r = 0.81 fm, which differs significantly from the value r = 0.862 fm. Instead, we use the parametrization by Simon *et al.* [7] with the result

$$\Delta E = 0.0232(15) \,\mathrm{meV} = r^3 \,0.0281 \,. \tag{25}$$

This parametrization in [7] is not correct for $Q^2 \equiv -q^2 \approx 1$ GeV or greater since for large Q^2 it behaves like Q^{-2} not Q^{-4} , however it does not influence the result at the presented precision level. This r^3 dependence in Eq. (25) is only an approximate dependence. This dependence become exact in the large nucleus mass limit with the dipole parametrization of proton formfactors. The result 0.0232(15) was obtained assuming r = 0.862(12) fm. If the proton radius r from μ H measurement will be significantly different, this result should be adjusted with the improved parametrization of proton formfactors.

Let us assume now, that the second order Born contribution is subtracted from T_1 and T_2 . The proton polarizability correction E_{POL} as given by Eq. (7) is split into two parts

$$E_{POL} = E_1 + E_2 \,, \tag{26}$$

$$E_1 = -2 e^4 \phi^2(0) \frac{m}{M} \int \frac{d^4 q}{(2\pi)^4 i} \frac{q^4 + 2\nu^4}{q^6(q^4 - 4m^2\nu^2)} T_2(\nu, q^2), \qquad (27)$$

$$E_2 = 2 e^4 \phi^2(0) \frac{m}{M} \int \frac{d^4q}{(2\pi)^4 i} \frac{q^2 + 2\nu^2}{q^4 (q^4 - 4m^2\nu^2)} \left(T_1(\nu, q^2) + \frac{\nu^2}{q^2} T_2(\nu, q^2)\right).$$
(28)

The first dominant part could be well calculated, while the calculation of the smaller second part will require additional assumptions. This second part was usually neglected in studies of polarizability effect in composite nuclei, see for example [11]. The imaginary part of T_i in variable ν at fixed q^2 is obtained from the inclusive cross section $\gamma^* + p \to X$. The real part could be restored from the imaginary one using the following dispersion relations:

$$T_2(\nu, q^2) = -\int_{\nu_{th}^2}^{\infty} d\nu'^2 \, \frac{W_2(\nu', q^2)}{\nu'^2 - \nu^2} \,, \tag{29}$$

$$T_1(\nu, q^2) = T_1(0, q^2) - \nu^2 \int_{\nu_{th}}^{\infty} \frac{d\nu'^2}{\nu'^2} \frac{W_1(\nu', q^2)}{\nu'^2 - \nu^2}, \qquad (30)$$

where ν_{th} is the threshold value of ν for the production of π mesons. This formulas are obtained from [12], by multiplying right hand side by -1/2, due to the different definition of T_i in our work. While W_2 for proton behaves like $1/\nu$ to large ν , W_1 goes like ν , and thus the dispersion relation in (30) contains a subtraction at $\nu = 0$. We have assumed here, that T_2 has similar asymptotic behavior as W_2 . i.e. vanishes in the large ν limit. However, there is no too much information on $T_1(0, q^2)$. One recognizes that in the limit of small q^2 it is given by the magnetic polarizability β_M

$$\lim_{q^2 \to 0} \frac{T_1(0, q^2)}{q^2} = \frac{M}{\alpha} \beta_M \,, \tag{31}$$

which amounts to $\beta_M = 1.56(57) \cdot 10^{-4}$ fm³, see [13]. Since there is no experimental data, we assume here, that the *q*-dependence is governed by the square of the elastic formfactor, namely

$$\beta_M(Q^2) = \beta_M \, \frac{\Lambda^8}{(\Lambda^2 + Q^2)^4} \tag{32}$$

with $\Lambda^2 = 0.71$ GeV². Using equations (27-31) one derives the following expressions for E_1 and E_2 .

$$E_1 = -\alpha^2 \frac{\phi^2(0)}{m M} \int_{\nu_{th}^2}^{\infty} d\nu^2 \int_0^{\infty} \frac{dt}{t^2} g(\nu, t) W_2(\nu, -t) , \qquad (33)$$

$$E_{2} = -\alpha^{2} \frac{\phi^{2}(0)}{m M} \int_{\nu_{th}}^{\infty} d\nu^{2} \int_{0}^{\infty} \frac{dt}{t^{2}} f(\nu, t) W_{2}(\nu, -t) \frac{1}{1+R} \left(1-R \frac{\nu^{2}}{t}\right) +\alpha^{2} \frac{\phi^{2}(0)}{m M} \int_{0}^{\infty} dt h(t) \frac{M}{\alpha} \beta_{M}(t) , \qquad (34)$$

where

$$g(\nu,t) = \frac{4}{\pi} \int_0^1 dx \sqrt{1-x^2} \, \frac{(1+2\,x^4)}{2\,(x^2+\frac{t}{4\,m^2})(x^2+\frac{\nu^2}{t})}\,,\tag{35}$$

$$f(\nu,t) = \frac{t}{2\nu^2} \frac{4}{\pi} \int_0^1 dx \sqrt{1-x^2} \frac{x^2(1+2x^2)}{(x^2+\frac{t}{4m^2})(x^2+\frac{\nu^2}{t})},$$
(36)

$$h(t) = \left[1 + \left(1 - \frac{t}{2m^2}\right) \left(\sqrt{\frac{4m^2}{t} + 1} - 1\right)\right], \qquad (37)$$

and R is equal to $R = W_L(\nu, q^2)/W_1(\nu, q^2) = \sigma_L/\sigma_T$, the ratio of longitudinal to transverse cross sections, and thus could be measured. A recent work [14] by E143 collaborations brings the most recent parametrization of R in the momentum range $Q^2 \ge 1$ GeV. What we find, is a weak dependence of R on ν for small Q^2 , of order few GeV, see for example Figure 3 of this work. Moreover R is between 0.3 - 0.4. However there is no precise data on the most important region for us of $Q^2 < 1$ GeV. One knows from conditions (12,13), that $R \sim Q^2$ at small Q^2 and constant ν . The only known paper by Brasse *et. al* [15] brings comments on R, that it is of order 10% at the resonance region. Since, what we find later on, E_2 is small and we do not have precise data on R at the resonance region we simply neglect R in the expression for E_2 in (34). We expect, this approximation will not alter significantly the final result for the polarizability correction. Various parametrization of structure functions W_2 were presented in the literature. We have chosen that, which match the $q^2 = 0$ limit known as a photoproduction and are correct for region of ν and Q^2 of order few GeV. We use the parametrization from [15] for the resonance region. It is given by a compact formula and a long table of numbers, see Table 1 of [15]. For ν close to the threshold for $\gamma^* + p^+ \rightarrow \pi^+ + n$ process, we assume a behavior $\sqrt{W - W_{th}}$ and match the value at W = 1.110 GeV, as given in [15]. The photoproduction of π^0 could be neglected close to this threshold value. Out of the resonance region, W > 1.990 GeV, we use an ALLM97 parametrization with the recent corrections [16]. With these parametrizations integrals in (33,34) are performed numerically with results

$$E_{POL} = E_1 + E_2 = -\alpha^2 \frac{\phi^2(0)}{m M} \left(1.82 + 0.12 - 0.26 \right) = -\alpha^2 \frac{\phi^2(0)}{m M} 1.68(33) , \qquad (38)$$

where the uncertainty of 0.33 forms 20% of final result and is an estimate due to approximations and assumptions performed during this derivation, namely: unknown R, unknown Q^2 dependence of β_M and inaccuracy of ALLM97 parametrization at low Q^2 and ν . The contribution to the (2P-2S) splitting of n = 2 states is

$$E_{POL} = 0.012(2) \,\mathrm{meV} \,.$$
 (39)

It is slightly lower than the result obtained by Rosenfelder in [5]. We think it is due to approximate treatment of the Q-dependence in his work. Other contributions to the Lamb shift in muonic hydrogen are presented in the Table I. It is an improved version of the former Table I in [8]. All QED corrections are recalculated with a better precision. The final result for the Lamb shift differs from that in [8] by more than the error estimate there. It is due to the mistake in the sum in Table I in [8] and due to the inclusion of new terms: proton polarizability, third order electron v.p. and the hadronic v.p. It should be noted here that the same effects should be included in hydrogenic Lamb shift, when having a new value of the proton charge radius. It was not necessary so far since in the electron scattering measurements of proton radius, these effects have not been excluded. The uncertainty in the final result has two main sources: the proton polarizability and the estimate of higher order QED effects. While the first one is pretty difficult to improve, the second source of uncertainty, higher order QED corrections could be well calculated. By these higher corrections we mean diagrams presented on Figure 1. Diagrams in the second raw, have been calculated in an approximate way in [8] and are named in the Table I as a muon self-energy with electron v.p. Some further theoretical work is necessary for obtaining the improved value of the hyperfine splitting of S-levels, since the proton polarizability effect is also present there. In summary, having now precise theoretical predictions, the measurement of the 2P-2S transition frequency in the muonic hydrogen will lead to the improved value of the proton charge radius.

ACKNOWLEDGMENTS

I wish to thank M. Krawczyk, E. Rondio for information on proton structure functions and F. Kottmann for pointing out a mistake in the sum of all contributions to the μ H Lamb shift in [8]. This work was supported by Polish Committee for Scientific Research under contract No. 2 P03B 024 11

REFERENCES

- F. Kottmann et al., Laser spectroscopy of the Lamb shift in muonic hydrogen, PSI Proposal R-98-03.1, 1998 (unpublished)
- [2] K. Pachucki, Phys. Rev. A **52**, 1079 (1997)
- [3] T. Kinoshita and M. Nio, Phys. Rev. Lett. 82, 3240 (1999)
- [4] J.L. Friar, J. Martorell, and D.W.L. Sprung, Phys. Rev. A 59, 4061 (1999)
- [5] R. Rosenfelder, hep-ph/9903352, Physics Letters to be published
- [6] R.N. Faustov and A.P. Martynienko, hep-ph/9904362
- [7] G.G. Simon, C.H. Schmidt, F. Borkowski, and V.H. Walther, Nucl. Phys. A 333, 381 (1980)
- [8] K. Pachucki, Phys. Rev. A 53, 2092 (1996)
- [9] J.L. Friar, Ann. Phys. (N.Y) **122**, 151 (1979)
- [10] J. Sapirstein and D. R. Yennie, in *Quantum Electrodynamics*, edited by T. Kinoshita (World Scientific, Singapore, 1990).
- [11] J. Bernabeu and T.E.O. Ericson, Z. Phys. A **309**, 213 (1983)
- [12] J. Bernabeu and C. Jarlskog, Nucl. Phys. B **75**, 59 (1974)
- [13] J. Tonnison, A.M. Sandorfi, S. Hoblit, and A.M. Nathan, Phys. Rev. Lett. 80, 4382 (1998)
- [14] K. Abe, et al., Phys. Lett. B **452**, 194 (1999)
- [15] F.W. Brasse, W. Flauger, J. Gayler, S.P. Goel, R. Haidan, M. Merkwitz, and H. Wriedt, Nucl. Phys. B 110, 413 (1976)
- [16] A. Abramowicz and A. Levy, hep-ph/9712415, there are few missprints in the values of several parameters describing W_2 structure function, correct values were obtained directly from the author (H.A.).
- [17] K. Pachucki, D. Leibfried, M. Weitz, A. Huber, W. König, and T.W. Hänsch, J. Phys. B. 29, 177 (1996)

TABLES

correction	value in meV
leading order e.v.p.	205.0074
rel. corr. to e.v.p .	0.0594
double e.v.p.	0.1509
two-loop e.v.p.	1.5079
three-loop e.v.p.	0.0076
muon self-energy + muon v.p.	-0.6677
muon self-energy with e.v.p.	-0.006(1)
recoil of order α^4	0.0575
recoil of order α^5	-0.0450
recoil of order α^6	0.0003
proton self energy	-0.0099
leading finite size of order α^4	$-r^2 5.1974 = -3.862(108)$
finite size of order α^5	$r^3 0.0363 = 0.0232(15)$
finite size of order α^6	-0.0009(3)
e.v.p. with finite size	$-r^2 0.0281 = -0.0209(6)$
hadronic v.p.	0.0113(3)
proton polarizability	0.012(2)
estimate for uncalculated terms	(0.002)
sum of corrections to Lamb	$206.085(3) - r^2 5.2255 + r^3 0.0363$
shift in μ H with $r = 0.862(12)$	= 202.225(108)

TABLE I. Summary of results for corrections to the Lamb shift in muonic hydrogen, e.v.p. denotes the electron vacuum polarization. QED corrections are calculated according to Ref. [17] and [8].

FIGURES



FIG. 1. Higher order diagrams contributing to muonic hydrogen Lamb shift. The upper and lower horizontal lines denote the muon and the proton respectively. A closed line denotes the electron loop.