Anharmonic force field and vibrational frequencies of tetrafluoromethane (CF_4) and tetrafluorosilane (SiF_4)

Xiao-Gang Wang*

Physical and Theoretical Chemistry Laboratory, University of Oxford, South Parks Road, Oxford OX1 3QZ, United Kingdom.

Edwin L. Sibert III

Department of Chemistry and Theoretical Chemistry Institute, University of Wisconsin — Madison, Madison, WI 53706, USA

Jan M. L. $Martin^{\dagger}$

Department of Organic Chemistry, Weizmann Institute of Science, 76100 Rehovot, Israel (Submitted to J. Chem. Phys. September 17, 1999)

Accurate quartic anharmonic force fields for CF_4 and SiF_4 have been calculated using the CCSD(T) method and basis sets of *spdf* quality. Based on the *ab initio* force field with a minor empirical adjustment, the vibrational energy levels of these two molecules and their isotopomers are calculated by means of high order Canonical Van Vleck Perturbation Theory(CVPT) based on curvilinear coordinates. The calculated energies agree very well with the experimental data. The full quadratic force field of CF_4 is further refined to the experimental data. The symmetrization of the Cartesian basis for any combination bands of T_d group molecules is discussed using the circular promotion operator for the doubly degenerate modes, together with tabulated vector coupling coefficients. The extraction of the spectroscopic constants from our second order transformed Hamiltonian in curvilinear coordinates is discussed, and compared to a similar procedure in rectilinear coordinates.

I. INTRODUCTION

Carbon tetrafluoride (also know as Freon-14) and tetrafluorosilane are molecules of wide ranging industrial, environmental, and economical interest. While the use of carbon tetrafluoride is being phased out together with that of other CFCs, it has a number of other important industrial and technological applications.¹ These include use as an isolator and extinguisher gas in high-voltage applications, and as an etching gas in the semiconductor industry.¹ The formation of $CF_4...O_2$ adducts was considered as a model system for the use of liquid perfluorochemicals in artificial blood.² Tetrafluorosilane, SiF₄, is a precursor (e.g. by glow discharge in SiF₄-H₂ mixtures) of amorphous Si-F-H semiconductors,¹ as well as for plasma deposition of low-dielectric Si-O-F thin solid films.³ It is the main by-product of beam etching of both semiconductors with fluorine, and silicon dioxide with fluorocarbons.⁴ In addition to being used to monitor the above processes [see e.g. Ref. 5], SiF₄ has been proposed⁶ as a remote volcano monitoring probe, since its presence can be measured using open-path FT-IR spectroscopy of $\nu_3(SiF_4)$. Finally, the isotopic separation of ³⁰Si (in ³⁰SiF₄) by infrared multi-photon dissociation (IRMPD) of natural-abundance Si₂F₆ using CO₂ lasers has been reported on a preparative scale.⁷

A key step for elucidating these processes is to understand the equilibrium structure, spectroscopy and energetics of these molecules. For these reasons the vibrational spectroscopy of the above molecules has been the subject of several theoretical and experimental studies. It is also the subject of the present paper in which we apply a combination of an accurate *ab initio* electronic structure treatment and a high-order CVPT treatment of the vibrational problem to the CF_4 and SiF_4 molecules.

Early gas-phase spectroscopic measurements of the CF₄ molecule^{8,9} have been reviewed and supplemented by Jones, Kennedy, and Ekberg (JKE).¹⁰ Earlier, Jeannotte *et al.*¹¹ obtained spectra in liquid argon solution, which as expected differ considerably from the JKE results due to solvent shifts. Early high-resolution work on the molecule was stimulated by the announcement¹² of mid-infrared CF₄ lasing based on the $\nu_2 + \nu_4 \rightarrow \nu_2$ transition. Esherick *et al.*¹³ determined ν_1 to high resolution by inverse Raman spectroscopy. High-resolution

^{*}Corresponding author for the vibrational calculation. Email: xgwang@emu.thchem.ox.ac.uk

[†]Corresponding author for the force field calculation. Email: comartin@wicc.weizmann.ac.il

Raman spectroscopy of ν_2 was carried out by Lolck,¹⁴ and of $2\nu_2$ and ν_1 by Tabyaoui *et al.*¹⁵ The $\nu_2 + \nu_4$ combination band was measured to high accuracy by Patterson *et al.*¹⁶ and by Poussigue *et al.*¹⁷ The ν_4 band was studied extensively by Jones *et al.*,⁹ by Tarrago *et al.*,¹⁸ and by McDowell *et al.*,¹⁹ the latter of whom determined ν_4 for all three isotopic species with Doppler-limited resolution. Doppler-limited measurements of the $2\nu_1 + \nu_4$ combination band were published by two different groups.^{20,21} A Fermi type 1 interaction exists between $2\nu_4$ and ν_3 ;²² detailed analyses of the ν_3 and $2\nu_3$ ²³ and of the $\nu_3/2\nu_4$ polyad²⁴ have been carried out.

The spectroscopic studies of CF₄ have been accompanied by several theoretical (spectroscopic) investigations. Harmonic valence force fields were first derived by by Duncan and Mills²⁵ and by Chalmers and McKean.²⁶ Jeannotte *et al.*²⁷ fitted a harmonic potential, supplemented by diagonal cubic and quartic stretching force constants only, to their earlier¹¹ liquid argon solution measurements. Brodersen²⁸ obtained a cubic force field from experimental rotation-vibration data. A quartic force field is said to be partly refined to rovibrational levels below 1400 cm⁻¹, but detailed information about the fitting is unavailable so far.²⁹ At higher energies, Boujut *et al.*³⁰ consider local and normal mode behavior in the ν_3 and ν_4 ladders of tetrahedral XY₄ and octahedral XY₆ molecules, with CF₄ being considered among the former. In their analysis they employ data from Ref. 24 as computerized into the STDS data bank.³¹ The $n\nu_3$ ladder has also been found to be a good testing ground for comparing experiment and theory. Using their results of high-resolution proton energy loss spectroscopy, Maring *et al.*,³² review earlier data for ν_3 , $2\nu_3$, and $3\nu_3$. They also summarize earlier unpublished data by Heenan³³ who derived two sets of Hecht³⁴-type anharmonic constants (to be denoted Heenan I and Heenan II in the remainder of the paper) from fitting Urey-Bradley type force fields to the Jeannotte¹¹ and JKE data, respectively. Maring *et al.* propose some alterations based on their measurements, including an apparently exceptionally large value for $X_{33} = -9.1$ cm⁻¹.

Early work on the spectroscopy of SiF₄ has been reviewed by McDowell *et al.*,³⁵ who note that research on the molecule was stimulated by the fact that the ν_3 fundamental overlaps the *P* branch of the 9400 nm band of the CO₂ laser. McDowell and coworkers reported a partial set of anharmonicity constants as well. In recent theoretical work, the low lying vibrational spectrum of SiF₄ has been modeled^{36,37} using an algebraic approach.³⁸

Patterson and Pine³⁹ determined $B_0=0.13676(3) \text{ cm}^{-1}$, whence $r_0=1.55982(17) \text{ Å}^{35}$. In later work, Takami and Kuze⁴⁰ and later Jörissen *et al.*⁴¹ substantially revised B_0 upward (to 0.137780439(92) cm⁻¹, consistent with a substantially shorter $r_0=1.55404 \text{ Å}$). As this paper was being finalized, we received a preprint by Demaison *et al.*⁴², in which $r_e=1.5524(8)$ is derived from a combination of *ab initio* and experimental results.

In the present paper we re-investigate the spectroscopy of these molecules using a combination of accurate *ab initio* anharmonic force fields and advanced techniques for solving the vibrational Schrödinger equation. In an earlier electronic structure study, Martin and Taylor⁴³ revised the heat of vaporization of silicon (which, among other things, is required for any *ab initio* or semiempirical calculation of the heat of formation of any silicon compound) from a benchmark *ab initio* calculation on SiF₄ from a very precise fluorine bomb calorimetric measurement⁴⁴ of the heat of formation of SiF₄(g) and a benchmark *ab initio* calculation. In part, we were motivated to study these molecules due to the success of a recent study in which, an *ab initio* quartic force field for methane⁴⁵ obtained using large basis sets and coupled cluster methods served as the starting point for several spectral refinement studies using variational methods,⁴⁶ low-order perturbation theory,⁴⁷ and high-order canonical Van Vleck perturbation theory (CVPT).⁴⁸ In general, we find that only the quadratic force constants and perhaps the geometry needs to be refined, and most of the remainder of the force field can be constrained to the *ab initio* values.

II. AB INITIO ANHARMONIC FORCE FIELD

All electronic structure calculations were carried out using MOLPRO 97⁴⁹ running on an SGI Origin 2000 minisupercomputer at the Weizmann Institute of Science.

Electronic correlation was treated at the CCSD(T) level [coupled cluster with all single and double substitutions (CCSD)⁵⁰ supplemented with a quasiperturbative estimate of the contribution of connected triple excitations⁵¹] method, using the implementation thereof in MOLPRO.⁵² This method is known⁵³ to yield correlation energies that are very close to the exact basis set correlation energy (i.e. the full configuration interaction result in the same basis set) as long as correlation is predominantly dynamical in character. As pointed out previously,⁴³ correlation in SiF₄ is essentially purely dynamical in character, and the same holds true for CF₄. For the record, values for the \mathcal{T}_1 diagnostic⁵⁴ (a measure for the importance of nondynamical correlation) are only 0.0118 for SiF₄ and 0.0119 for CF₄, respectively. (In all calculations reported here, only valence electrons were correlated.)

Geometry optimizations were carried out by univariate polynomial interpolation. At the reference geometry, a quartic force field in curvilinear symmetry coordinates was determined by recursive application of the central finite difference formula with step size 0.01 Å or radian. (Quartic contamination was removed from the quadratic force constants.) The symmetry coordinates are the same as those used in the previous study on CH_4 .⁴⁵

The resulting curvilinear internal coordinate force field was then transformed to rectilinear normal coordinates, then a standard second-order rovibrational perturbation theory⁵⁵ was carried out by means of the SPECTRO program.^{56,57} The alignment conventions for the anharmonic constants of a spherical top follow the work of Hecht³⁴ and general formulae for these constants were taken from the paper by Hodgkinson *et al.*.⁵⁸ (See Section III.B for a discussion about the definition of the ν_2 normal coordinates.) The implementation in SPECTRO was slightly modified to allow deperturbing for an arbitrary number of Fermi resonances,⁵⁹ and a simple routine implementing Hecht's³⁴ energy level formulas up to four vibrational quanta was added. Similar procedure has also been implemented in the curvilinear formalism, and the relationship between the two formalism will be discussed in Section III.B. As in past work (e.g. Ref. 45), correct alignment was verified by slightly (0.00001 a.m.u.) perturbing the masses of two atoms, then repeating the analysis in the asymmetric top formalism.

Given the n^3N^4 scaling (with *n* the number of electrons correlated and *N* the number of basis functions) of CPU time for a CCSD(T) calculation, the large number of valence electrons correlated (*n*=32) imposes certain restrictions on the size of the basis set for a complete quartic force field. We settled on the standard Dunning cc-pVTZ (correlation consistent polarized valence triple zeta) basis set^{60,61} (150 contracted Gaussian-type functions, CGTFs) for CF₄, while for SiF₄, we opted for the cc-pVTZ+1 basis set (159 CGTFs), where the suffix "+1" denotes the addition of a single high-exponent *d* function on second-row elements⁶² to accommodate inner polarization effects. The latter has been repeatedly shown to be essential for accurate computed properties of second-row molecules with polar bonds,⁶²⁻⁶⁴ and the addition of a mere five functions to the basis set increases CPU time insignificantly for our purposes. It should be pointed out that even with these relatively compact basis sets, the evaluation of each force field took the equivalent of six weeks of CPU time on the SGI Origin.

For purposes of validation, we in addition calculated geometries and harmonic frequencies at the CCSD(T) level with a number of other basis sets. For CF₄, we considered the (diffuse-function) augmented cc-pVTZ basis set aug-cc-pVTZ of Kendall *et al.*⁶⁵ (230 CGTFs), while for SiF₄, we considered both the standard cc-pVTZ basis set (154 CGTFs) and the aug-cc-pVTZ+1 basis set (239 CGTFs), in which the "+1" suffix again denotes the addition of an inner-polarization d function. The results are collected in Tables I and II.

For CF₄, the main deficiency appears to be that the asymmetric stretching frequency ω_3 is seriously overestimated. This is a known problem for polar bonds, particularly those involving fluorine .⁶⁶⁻⁶⁸ Increasing the basis set size to cc-pVQZ (275 CGTFs) or even cc-pV5Z (455 CGTFs) as a rule does not remedy this problem^{66,68} — and given the N⁴ scaling behavior would increase CPU time requirements by one and two orders of magnitude, respectively, and would therefore be essentially impossible. The use of anion basis set (i.e. aug-cc-pVnZ) on at least the electronegative atoms themselves generally leads to a dramatic improvement in basis set convergence,⁶⁶⁻⁶⁸ and here too we see that CCSD(T)/aug-cc-pVTZ frequencies for CF₄ are much closer to the final adjusted values than the CCSD(T)/cc-pVTZ values. While in HF, this sensitivity extends to the anharmonicities,⁶⁸ in heavy-atom systems like FNO and ClNO⁶⁷ the anharmonic force field at the CCSD(T)/aug-cc-pVTZ level, which would have increased CPU time requirements approximately fivefold.

Interestingly, the effect of adding the anion functions is much less pronounced for SiF_4 than it is for CF_4 . The effect of adding the inner polarization function is fairly pronounced on the bond distance (a decrease of almost 0.01 Å), but effects on the harmonic frequencies are fairly minor. Here too, we judged that the CCSD(T)/cc-pVTZ+1 level represented the best compromise between accuracy and computational cost for the entire anharmonic force field.

Purely *ab initio* calculation of the geometry and the harmonic part of the potential would presumably be feasible as far as basis set convergence and inner-shell correlation corrections are concerned, but we know from experience⁶⁸ that imperfections in the CCSD(T) method might still account for (usually positive) errors of several cm⁻¹ in the frequencies. Full configuration interaction calculations are out of the question for this system, and hence we have followed a different track similar to previous work on silane,⁶⁹ and refined the 'raw' CCSD(T)/cc-pVTZ and CCSD(T)/cc-pVTZ+1 force fields in the following way.

(1) Using standard second-order rovibrational perturbation theory, the four vibrational fundamentals for the most abundant isotopomer were obtained, together with the equilibrium and ground-state rotational constants B_e and B_0 , respectively. Hence (as $r_e\sqrt{B_e/B_0}$) the computed r_0 was obtained.

(2) Using the internal coordinate force field given above, the four diagonal quadratic force constants and r_e were then iteratively adjusted such that the fundamentals and r_0 obtained by second-order rovibrational perturbation theory coincide with the experimental values to the precision given. (F_{47} and all cubic and quartic force constants were held at their *ab initio* computed values.) This process converged after three iterations, and the final force fields are given in Table III. This force field was then used as input for the higher-order vibrational analysis.

As can be seen in Tables I and II, the adjustment only entails relatively minor changes in the geometry and harmonic frequencies. In addition, overall agreement between the available experimental energy levels and the values computed using standard second-order perturbation theory is such that we conclude that the CCSD(T)/VTZ(+1) quartic force fields are of sufficient quality for our purposes. (Since the computed energy levels are essentially identical to those obtained by means of second-order CVPT if no resonances are accounted for, the reader is referred to Tables VI and VIII and the relevant sections below for details.)

Concerning the equilibrium geometries, we note that our final r_e values for CF₄ and SiF₄ agree with the experimentally derived values of Brodersen²⁸ and Demaison *et al.*,⁴² respectively, to within their stated uncertainties.

III. CALCULATION OF VIBRATION ENERGY LEVELS

In this and the following two sections, in order to assess the accuracy of the *ab initio* force field, the vibrational energy levels of CF_4 and SiF_4 are calculated using high order canonical Van Vleck perturbation theory(CVPT) and compared to the experimental data. The calculation is similar to that which has been recently applied to the methane and its isotopomers. Details about this calculation can be found in Ref. 48, and only the major procedures will be summarized here.

Firstly the exact J = 0 vibrational Hamiltonian including the pseudo-potential V' term⁷⁰ is expanded in terms of the curvilinear normal coordinates based on the Morse coordinates for the stretch and angle extension coordinates for the bend. The redundancy relation between the six bond angles is properly taken into account in the expansion. As in the case of methane and its isotopomers,⁴⁸ the kinetic energy operators are expanded to fourth order, even though CVPT is carried out to a higher order. We choose this scheme because the kinetic contributions of order higher than four are believed to be small, and the number of terms rapidly increases with the order of the expansion. The quartic potential which is already in expanded form is used throughout this work. It contributes to the CVPT up to the second order. It is re-expressed in the coordinates described above so that the new expansion of the potential agrees with the original potential up to quartic terms.

The expanded Hamiltonian is separated in the form

$$H_v = H^{(0)} + \lambda H^{(1)} + \lambda^2 H^{(2)} + \dots + \lambda^n H^{(n)}, \tag{1}$$

where λ is the perturbation parameter. A succession of canonical transformations of the form

$$K_v = \exp\{i\lambda^n [S^{(n)},]\} \cdots \exp\{i\lambda^2 [S^{(2)},]\} \exp\{i\lambda [S^{(1)},]\} H_v$$
(2)

are then applied to the Hamiltonian of Eq. (1), where the $S^{(n)}$ are chosen such that the matrix representation of K_v is block diagonal through order n.^{71,72} The blocks are characterized by one or more polyad quantum numbers

$$N_t = \sum_i m_i v_i,\tag{3}$$

where the integers m_i define the block. These integers are chosen such that the resonance interactions are not perturbatively decoupled. Details regarding the choice of the polyad quantum numbers will be discussed later with respect to specific molecules.

Two points, regarding our analysis of this Hamiltonian are now considered in more detail. First, in order to obtain and analyze the eigenvalues of the individual blocks we diagonalize each block in a *symmetrized basis* to reduce the matrix size and to help assign the calculated levels. Second, the transformed (effective) Hamiltonian of Eq. (2) at second order is particularly interesting as its coefficients are the quartic anharmonicity constants and resonance constants, which have been conventionally used by spectroscopists to highlight the features of the vibrational spectra. In the following two subsections we discuss the basis symmetrization and extraction of spectroscopic constants as they have not been given in our previous work.⁴⁸

A. Symmetrization in the Cartesian basis for the T_d point group

There are generally two ways of choosing basis functions for the doubly and triply degenerated normal modes for the T_d group molecules. The first choice (angular momentum basis) are eigenfunctions of the vibrational angular momenta. The second choice (Cartesian basis) is the products of harmonic oscillator functions including different Cartesian components of the degenerate vibrations. In the angular momentum basis, the off-diagonal terms are the spherical tensor operators defined by Hecht³⁴ and their matrix elements are calculated using the Wigner-Eckart theorem. However it is more difficult to treat vibrational resonances in the angular momentum basis, and therefore the alternative Cartesian basis is more often used. Another advantage of the Cartesian basis is that the matrix elements are readily calculated since they all reduce to one-dimensional harmonic oscillator matrix elements. For both basis, the symmetrization is nonobvious and we are going to discuss the symmetrization in the Cartesian basis below.

The Cartesian basis denoted by $|n_1; n_{2a}, n_{2b}; n_{3x}, n_{3y}, n_{3z}; n_{4x}, n_{4y}, n_{4z}\rangle$ will be symmetrized into the irreducible representations of the T_d point group: A_1, A_2, E, F_1 and F_2 . Various equivalent approaches can be used, such as the projection operator technique,^{47,73} and the promotion operator technique with tabulated vector coupling coefficients.^{74,75} The vector coupling coefficients (also called Clebsch-Gordan coefficients) $\langle \Gamma_1 \Gamma_2 ab | \Gamma_1 \Gamma_2 \Gamma c \rangle$ defined in

$$|\Gamma_1 \Gamma_2 \Gamma c\rangle = \sum_{a,b} |\Gamma_1 \Gamma_2 ab\rangle \langle \Gamma_1 \Gamma_2 ab | \Gamma_1 \Gamma_2 \Gamma c\rangle \tag{4}$$

and tabulated in Table A20 of Ref. 76 (See also Table 1 of Ref. 77) essentially determine how the product basis from two basis $|\Gamma_1 a\rangle$ and $|\Gamma_2 b\rangle$ are further reduced according to the irreducible representation $|\Gamma_c\rangle$ of the point group. In this work, the latter method is adopted with some improvements to attain unique symmetrization.

First of all, the overtone of the $E(\nu_2)$ and $F_2(\nu_3$ and ν_4) normal modes are symmetrized separately. For the symmetrization of the overtone of the E normal modes, the promotion operators technique with tabulated vector coupling coefficients has been used in Ref. 75. The method is summarized as follows. The promotion operators with (E_a, E_b) symmetry are the familiar creation operators which act on the basis by

$$a_{a}^{\dagger}(E_{a})|n_{a},n_{b}\rangle = \sqrt{n_{a}+1}|n_{a}+1,n_{b}\rangle$$

$$a_{b}^{\dagger}(E_{b})|n_{a},n_{b}\rangle = \sqrt{n_{b}+1}|n_{a},n_{b}+1\rangle.$$
(5)

Taking the promotion operators to belong to (E_a, E_b) symmetries, the symmetry of the resultant promoted basis function can be identified according to the vector coupling coefficients. Starting from the v=1 basis function, one can derive all the symmetrized basis functions until v reaches a given number similar to climbing a v ladder. A computer code is desirable if one wants to symmetrize a high overtone. One problem is that spurious states with the same symmetry may arise if the promotion operator acts on every basis, and they are not orthogonal. For example, with promotion operators acting on the full $3\nu_2$ symmetrized basis $(A_1 + A_2 + E)$, three nonorthogonal E states are obtained while only two are allowed for $4\nu_2$. These spurious states are removed by using the Gram-Schmidt orthogonalization technique. The final orthogonal symmetrized basis is not unique, but all possible solutions lead to symmetrized Hamiltonian matrix blocks.

An alternative method, which yields unique symmetrization coefficients, is obtained by finding the exact decomposition of the angular momentum basis function into the Cartesian basis wavefunction. Once this is done, the symmetrization is trivial as the angular momentum basis function is readily symmetrized according to the value of l modulo 3. This task seems quite complicated since the angular momentum basis functions are basically the associated Laguerre polynomials and the Cartesian basis functions are essentially products of two Hermite polynomials.^{78,79} In fact, this complication can be circumvented by using the circular promotion operators defined by⁸⁰

$$a_d^{\dagger} = \frac{1}{\sqrt{2}} (a_a^{\dagger} + ia_b^{\dagger})$$

$$a_g^{\dagger} = \frac{1}{\sqrt{2}} (a_a^{\dagger} - ia_b^{\dagger})$$
(6)

They act on the angular momentum basis by

$$a_d^{\dagger} |v^l\rangle = \sqrt{\frac{(v+l)}{2} + 1} |(v+1)^{l+1}\rangle$$

$$a_{g}^{\dagger}|v^{l}\rangle = \sqrt{\frac{(v-l)}{2} + 1}|(v+1)^{l-1}\rangle \tag{7}$$

and on the Cartesian basis by

$$a_{d}^{\dagger}|n_{a},n_{b}\rangle = \sqrt{\frac{n_{a}+1}{2}}|n_{a}+1,n_{b}\rangle + i\sqrt{\frac{n_{b}+1}{2}}|n_{a},n_{b}+1\rangle$$

$$a_{g}^{\dagger}|n_{a},n_{b}\rangle = \sqrt{\frac{n_{a}+1}{2}}|n_{a}+1,n_{b}\rangle - i\sqrt{\frac{n_{b}+1}{2}}|n_{a},n_{b}+1\rangle$$
(8)

With the circular promotion operators acting on the v = 0 ground state $|0^0\rangle = |0,0\rangle$ in the two basis sets, respectively, one obtains the relationship for the v = 1 basis

$$|1^{\pm 1}\rangle = \frac{\sqrt{2}}{2}(|10\rangle \pm i|01\rangle) \tag{9}$$

Repeating this operation up to a given quantum number v leads to relationship of the two basis up to v. The next step is to transform the angular momentum basis from the (E^+, E^-) to (E_a, E_b) symmetries. It is well known that $|v^{\pm l}\rangle$ (l = 3p + 1, with p being any integer) form a basis for E^{\pm} irreducible representations. Following a convention which is learned from Eq. (9), the (E_a, E_b) symmetry basis functions are defined by

$$|v^{|l|}; E_a\rangle = \frac{\sqrt{2}}{2} (|v^{+l}\rangle + |v^{-l}\rangle) = \sqrt{2} \operatorname{Re}(|v^{+l}\rangle)$$
$$|v^{|l|}; E_b\rangle = \frac{\sqrt{2}}{2i} (|v^{+l}\rangle - |v^{-l}\rangle) = \sqrt{2} \operatorname{Im}(|v^{+l}\rangle)$$
(10)

where l = 3p + 1. The A_1/A_2 symmetry basis functions are defined similarly by

$$|v^{|l|}; A_1 \rangle = \frac{\sqrt{2}}{2} (|v^{+|l|}\rangle + |v^{-|l|}\rangle) = \sqrt{2} \operatorname{Re}(|v^{+|l|}\rangle)$$
$$|v^{|l|}; A_2 \rangle = \frac{\sqrt{2}}{2i} (|v^{+|l|}\rangle - |v^{-|l|}\rangle) = \sqrt{2} \operatorname{Im}(|v^{+|l|}\rangle)$$
(11)

where l = 3p. There is only A_1 symmetry if l is zero. Pak *et al.*⁸¹ have also discussed a similar scheme to transform the angular momentum basis functions to the real cosine and sine form, which are members of the irreducible representation of the D_{3h} group. It should be noted that the above results can be applied to other analogous doubly degenerated modes such as the E mode in the C_{3v} group and the E_g mode in the O_h group. The above procedure can be easily implemented by a computer code. Results up to v = 8 are listed in Table IV, which are obtained from a code in the Maple⁸² computer algebra language. These results may be useful for other applications. A fortran code is also written for the same purpose and is to be combined with the symmetrization of the F_2 normal mode to form a general symmetrization package.

Now we turn to the symmetrization of the overtone of the F_2 normal mode. This can also be achieved using the promotion operators together with the vector coupling coefficients. However there exist general rules on how a given overtone is symmetrized without resorting to promotion of lower states. These general rules can be derived using the projection operator methods. Roughly speaking, one obtains AE symmetry when the three Cartesian quantum numbers are all even or odd, and F symmetry when they are mixed even and odd. Detailed symmetrization coefficients have been given in Table 2 of Ref. 30 for the overtone of the triply degenerated modes (F_1 and F_2 modes of T_d group) of O_h and T_d group.

Finally given the fact that the E overtone and F_2 overtone have been symmetrized separately, the combination bands in the form $(n_1n_2n_3n_4)$ can be symmetrized using the vector coupling coefficients, as has been demonstrated in Ref. 75. In this process, no spurious states arise as the basis functions joining the product are from different normal modes. All the above steps have been implemented as a Fortran code which writes to a file the symmetrization coefficients for all the combination bands in a polyad with a given polyad number. This is highly desirable since one may encounter completely different polyad numbers such as those for CH_4 , CF_4 and SiF_4 . These symmetrization coefficients are further used in constructing the polyad block matrix of the transformed Hamiltonian.

It is interesting to compare with the basis symmetrization for the T_d group in the algebraic model.^{83–85} A key difference is that the basis functions symmetrized there are products of ten functions based on the ten

internal coordinates. Due to the redundancy between the six bond-angle coordinates considerable effort was required to remove the so-called spurious states.^{83,84} Their symmetrization procedure is basically the same as the promotion operator technique with tabulated vector coupling coefficients.^{74,75} Lemus and Frank⁸³ have also tabulated and derived part of the vector coupling coefficients in the T_d group, unaware of an earlier full tabulation.^{77,76}

B. Extraction of spectroscopic constants

The spectroscopic constants we want to extract from our second order transformed Hamiltonian are specifically the quartic anharmonicity constants X_{ij} , G_{ij} , the tensor constants T_{ij} and S_{ij} and the cubic $K_{i,jk}$ and quartic resonance constants $K_{i,jkl}$ and $K_{ij,kl}$. The quartic anharmonicity constants X_{ij} and G_{ij} have their usual definitions. The tensor constants were defined by Hecht³⁴ as coefficients of spherical tensor operators which cause splittings of the same symmetry components of the overtone and combination bands of spherical top molecules. This procedure has been used in earlier work on the acetylene molecule,⁸⁶ however the procedure used was not described. More importantly, the procedure has not been applied to molecules of T_d symmetry which present a special challenge. For these reasons we describe the procedure in full detail here.

The apparent difficulty of the extraction is that our CVPT results for the transformed Hamiltonian are expressed as expansions of creation-annihilation operators in normal form^{71,72} — the creation operators are put before the annihilation operators. These expansion terms need to be rearranged to obtain the traditional spectroscopic Hamiltonian. A simple example based on second order perturbation theory serves to illustrate this point. At second order the normal form expansion includes contributions, such as $(a_1^{\dagger})^2(a_1)^2 = \hat{n}_1(\hat{n}_1-1)$, $a_1^{\dagger}a_1 = \hat{n}_1$ plus a constant. It is found that by rearranging the normal form terms as powers of $(\hat{n}_1 + \frac{1}{2})$, all the linear contributions to second order of perturbation theory cancel, leaving the Hamiltonian in the standard spectroscopic form where only quadratic terms such as $(\hat{n}_1 + \frac{1}{2})^2$ contribute. With either choice of expansion the leading term will have the same coefficient which is X_{11} .

A similar problem has been studied by Hodgkinson *et al.*⁵⁸ who have suggested the same idea of extracting spectroscopic constants from the second order transformed Hamiltonian, although the Hamiltonian discussed there is in rectilinear normal coordinates and the creation-annihilation operators are not rearranged to normal form. Furthermore, the extraction of constants related with doubly and triply degenerated modes requires one more step than the above simple example of X_{11} . For example, the tensor constants T_{23} and anharmonicity constants X_{23} are determined simultaneously by

$$Coef(\hat{n}_{2a}\hat{n}_{2z}) = X_{23} + 8T_{23}$$

$$Coef(\hat{n}_{2b}\hat{n}_{2z}) = X_{23} - 8T_{23}$$
(12)

where $\operatorname{Coef}(\hat{o})$ denotes coefficient of operator \hat{o} , and $\hat{n}_{2a}\hat{n}_{2z}$ and $\hat{n}_{2b}\hat{n}_{2z}$ are the leading terms in the normal form Hamiltonian. These formulae are readily obtained by rewriting the O_{23} operator which was defined by Hecht [Eq. (8) of Ref. 34] as an operator whose coefficients are T_{23} . Such formulae for all the anharmonicity and tensor constants have been given in Table 5 of Ref. 58.

Here one has to take care of a convention problem regarding the definition of normal/symmetry coordinates for the ν_2 (*E*) mode of a tetrahedral molecule. We notice that Jahn defined the symmetry coordinate (Q_{2a}, Q_{2b}) on page 472 of Ref. 87 (See Fig. 1 of Ref. 88 for the numbering of the four peripheral atoms). His definition and the vector coupling coefficients for the *E* mode coupling other modes (Table II of Ref. 89) were adopted by Hecht in Ref. 34. Unfortunately the symmetry coordinates (S_{2a}, S_{2b}) for the ν_2 mode we and many other researchers^{90,45,75,47} adopted are those of Mills.^{91,90} After some manipulations, it is found that the two definitions are related such that $Q_{2a} = -S_{2b}, Q_{2b} = S_{2a}$. If one uses Mills' definition, all the Hecht formulae regarding the ν_2 mode should be modified according to the above relationship. The only significant change occurs with the O_{23} operator for which a minus sign needs to be attached to Hecht's definition. Only after this correction is considered, can the Table 5 of Hodgkinson *et al.* be reconciled with Hecht's definition. Eq. (12) is a pertinent example. Finally, it is noted that the vector coupling coefficients tabulated by Tababe and Sugano⁷⁷ and Griffith⁷⁶ agree with Mills' definition of the symmetry coordinates of ν_2 .

The extraction of the resonance constants is straightforward in the creation-annihilation form since they are usually defined in this form. One just needs to take care of constructing the correctly symmetrized Hamiltonian terms for the resonance. To this end, several equivalent approaches can be used. Other than the method by Hecht^{34,92} for constructing the correctly symmetrized potential tensor operators, we choose

to follow an elegant and more general approach by Halonen⁷⁵ who used the tabulated vector coupling coefficients to construct the correctly symmetrized Hamiltonian terms from any combination of operators such as coordinates, momenta and creation-annihilation operators. Methane has a variety of cubic and quartic resonances due to its approximate 2:1 ratio of the stretch and bend frequency. Halonen has given for the first time all the correctly symmetrized Darling-Dennison type quartic resonance terms in Ref. 75 using the vector coupling methods. The resonance constants encountered in CF_4 and SiF_4 are limited compared to methane. Up to second order resonance, they are (See also Eqs. (19) – (21) and (23) – (25))

$$H/hc = K_{1,22} \left[a_1^{\dagger} (a_{2a}^2 + a_{2b}^2) + \text{h.c.} \right]$$
(13)

$$H/hc = K_{3,44} \left[a_{3x}^{\dagger} a_{4y} a_{4z} + a_{3y}^{\dagger} a_{4z} a_{4x} + a_{3z}^{\dagger} a_{4x} a_{4y} + \text{h.c.} \right]$$
(14)

for CF_4 and

$$H/hc = K_{1,44} \left[a_1^{\dagger} (a_{4x}^2 + a_{4y}^2 + a_{4x}^2) + \text{h.c.} \right]$$
(15)

$$H/hc = K_{1,222} \left[a_1^{\dagger} a_{2a} (a_{2a}^2 - 3a_{2b}^2) + \text{h.c.} \right]$$
(16)

for SiF₄. The quartic resonance with coefficient $K_{1,222}$ is of a new type. We shall show how it is derived to illustrate the vector coupling method. The goal is to construct the totally symmetric (A_1 symmetry) quartic terms from one creation operator (a_1^{\dagger}) with A_1 symmetry and two annihilation operators (a_{2a}, a_{2b}) with E symmetry. Apparently there is only one such term. First the E_a/E_b symmetry quadratic terms are constructed from products of (a_{2a}, a_{2b}) and themselves as follows

$$E_{a} : \frac{1}{\sqrt{2}} (-a_{2a}^{2} + a_{2b}^{2})$$

$$E_{b} : \sqrt{2}a_{2a}a_{2b}$$
(17)

Then the A_1 symmetry cubic terms are constructed from products of (a_{2a}, a_{2b}) and the two terms of Eq. (17) as follows

$$A_{1} : \frac{1}{\sqrt{2}} \left[\frac{1}{\sqrt{2}} \left(-a_{2a}^{2} + a_{2b}^{2} \right) \cdot a_{2a} + \sqrt{2}a_{2a}a_{2b} \cdot a_{2b} \right]$$
$$= -\frac{1}{2}a_{2a}\left(a_{2a}^{2} - 3a_{2b}^{2}\right)$$
(18)

Finally the A_1 symmetry quartic term is obtained as product of a_1^{\dagger} and terms of Eq. (18) plus the Hermite conjugate of the product. The coefficient of the final resonance term in Eq. (16) has been redefined for convenience.

An interesting point is that the anharmonicity and tensor constants extracted using the method above agree exactly with standard second-order perturbation theory in rectilinear normal coordinates as implemented in SPECTRO, provided that the same force field is used and no resonances are considered. This seems to be a surprise, since the reported CVPT results are based on curvilinear coordinates. McCoy and Sibert⁹³ have explained this as follows. When using dimensionless normal coordinates the perturbation parameter in the expansion of Eq. (1) can be taken as $\hbar^{1/2}$. Here terms of order *n* in Eq. (1) are of order $\hbar^{(2+n)/2}$. This is true regardless of using rectilinear or curvilinear normal coordinates. Since both sets have identical zero-order Hamiltonians, the energies have to be identical order by order. If one uses perturbation theory to transform to block-diagonal Hamiltonian, which is subsequently diagonalized, the above argument breaks down, and notable differences have been found.^{94,95} In this work, a multi-resonance Hamiltonian has been considered for CF₄ and SiF₄. Therefore the spectroscopic constants are extracted from the second order CVPT transformed Hamiltonian, and some of them are different from those obtained using SPECTRO with resonances deperturbed. More details regarding this point are given with respect to CF₄ and SiF₄ molecules in the following section.

IV. RESULTS FOR CF₄

The choice of the polyad quantum numbers for CF₄ is based on the following considerations. The harmonic frequencies as calculated from the *ab initio* force fields ($\omega_1 = 921.57 \text{ cm}^{-1}$, $\omega_2 = 439.91 \text{ cm}^{-1}$, $\omega_3 = 1303.01 \text{ cm}^{-1}$, $\omega_4 = 637.89 \text{ cm}^{-1}$) suggest that three independent resonances are possible:

$$\omega_1 \approx 2\omega_2 \tag{19}$$

$$\omega_3 \approx 2\omega_4 \tag{20}$$

$$3\omega_2 \approx \omega_3$$
 (21)

The frequency differences between the above pairs are only 42 cm^{-1} , 27 cm^{-1} and 17 cm^{-1} , respectively. Since these small differences will go into the denominator of the expression of $S^{(n)}$, they will possibly lead to divergence of the perturbation theory when the coupling strength between the two pair states are not negligible. In practice we found that all the three resonances ought to be considered by keeping them in the K_v to ensure good convergence when the perturbation theory is carried up to the sixth order. It should be noted that the resonance in Eq. (21) is actually a fourth order resonance, because the symmetry is unmatched between $3\nu_2$ and ν_3 at second order. Due to the three independent resonances, only one good quantum number remains, which should be orthogonal to the three resonance vectors (1, -2, 0, 0), (0, 0, 1, -2)and (0, -3, 1, 0). It is then easy to find the remaining good quantum number to be

$$N = 4v_1 + 2v_2 + 6v_3 + 3v_4, \tag{22}$$

which will be used in the perturbation calculations reported below.

In Table VI, we collected all the reliable experimental band origins of the three isotopes. There are 14 high resolution data (with resolution better than 0.01 cm^{-1}). All the other data with typical 0.1 and 1 cm⁻¹ resolution are from the work of Jones *et al.*¹⁰ There are some other low-resolution data compiled in Ref. 10, which we found to be unreliable and were discarded. The calculated energy levels based on the adjusted *ab initio* force field of Table III agree well with the observations. We then further refined all five quadratic force constants to the observed levels with uncertainty for each level determined by the experimental precision, using a fast convergent second order least squares method where the Hessian is approximated as products of first derivatives.⁹⁶ The refined constants are given in Table V and the comparison of experimental and calculated energy levels is given in Table VI. The RMSD (root mean squares deviations) based on the refined potential is 0.179, 0.083 and 0.083 cm⁻¹ at second, fourth and sixth order CVPT, respectively, whereas the corresponding RMSD based on the *ab initio* potential in Table III is 0.095, 0.190 and 0.186 cm⁻¹. This shows that the *ab initio* potential is a rather good as an initial potential for the fitting. The following remarks can be made by examining the calculated results in Table VI.

(1) The CVPT calculation converges very well. The energy level changes typically on the order of 0.1 cm^{-1} by comparing the fourth and sixth order results. This contrasts with the case of CH₄ where the bending overtone $(4\nu_4)$ varies by about three wavenumbers at the eighth order.⁴⁸ Therefore the fourth order CVPT is used for the fitting. The contribution due to the V' terms is small. For example, the largest contribution to the fundamentals is 0.057 cm^{-1} for $(0001; F_2)$ band of ¹²CF₄ using sixth order CVPT.

(2) For ¹²CF₄, the resonance $\omega_3 \approx 2\omega_4$ is the most pronounced. Even the (0010; F_2) fundamental has 7 % admixture from (0002; F_2). The resonance $\omega_1 \approx 2\omega_2$ is less pronounced and the resonance $3\omega_2 \approx \omega_3$ can be ignored. The above remarks also apply for the ¹³CF₄ whereas all the three resonances seem to be less pronounced for ¹⁴CF₄.

The pronounced resonance between $\omega_3 \approx 2\omega_4$ means that previous treatment for the ν_3 ladder states where only the ν_3 overtones are considered are inadequate.^{23,30,32}

(3) The calculations for the three (0020) bands using either the *ab initio* or fitted force field cannot reproduce well the observations which were obtained through a rovibrational analysis of the isolated (0020) bands.²³ We cannot fully understand these discrepancies. But it is clear that the (0020) band is in Fermi resonance with the (0012) and (0004) bands, although the major perturbant (0012) is a little farther away. For example, the three (0012; F_2) bands are respectively at 2536.50 cm⁻¹, 2541.66 cm⁻¹, and 2544.45 cm⁻¹. We have also noted that near the experimental (0020; F_2) band origins (2561.91 cm⁻¹), there is a band calculated to be 2561.04 cm⁻¹ with 82%(0302; F_2) and 14%(0310; F_2). However it is very unlikely that this band should be as strong as the (0020; F_2) band. Further theoretical and experimental work on this band system appears to be desirable.

Finally, the vibrational spectroscopic constants for all the isotopes of CF_4 and SiF_4 obtained from the second order transformed CVPT Hamiltonian (curvilinear formalism) are given in Table VII. With these constants one can reproduce the second order CVPT calculation by constructing the block diagonal Hamiltonian matrices. The corresponding constants in the rectilinear formalism, where different, are given in footnotes to said table. The differences (for the Fermi resonance constants and for such anharmonic constants as are affected by the Fermi interactions) illustrate the points made in the previous section.

Discrepancies between the presently computed anharmonicity constants and the empirically derived sets of Heenan³³ mostly reflect the limitations of the latter in terms of available data. There is no indication

for the very high $X_{33} = -9.1 \text{ cm}^{-1}$ as suggested by Maring *et al.*³² We also note that both Heenan sets of harmonic frequencies differ quite substantially from our best values, and are confident that the latter are more reliable.

V. RESULTS FOR SIF₄

The choice of the polyad quantum numbers for SiF₄ is based on the following considerations. The harmonic frequencies as calculated from the *ab initio* force fields ($\omega_1 = 806.3 \text{ cm}^{-1}$, $\omega_2 = 265.2 \text{ cm}^{-1}$, $\omega_3 = 1044.0 \text{ cm}^{-1}$, $\omega_4 = 389.3 \text{ cm}^{-1}$) suggest that three independent resonances are possible:

$$\omega_1 \approx 2\omega_4 \tag{23}$$

$$\omega_1 + \omega_2 \approx \omega_3 \tag{24}$$

$$\omega_1 \approx 3\omega_2 \tag{25}$$

The differences between the above pairs are only 27 cm⁻¹, 27 cm⁻¹, and 11 cm⁻¹, respectively. Again all three resonances ought to be considered to ensure good convergence when the perturbation theory is carried up to sixth order. It should be noted that the resonance in Eq. (24) is actually a third order resonance, because the symmetry is unmatched between $\nu_1 + \nu_2$ (F_2) and ν_3 (E) at first order. Following the approach of the previous section the good quantum number, to be used in the perturbative calculations, is found to be

$$N = 6v_1 + 2v_2 + 8v_3 + 3v_4 \tag{26}$$

which we are going to use in the perturbation calculations reported below.

The sixteen observed band centers from Ref. 35 and the values of the second, fourth and sixth order CVPT calculations are given in Table VIII. The following remarks can be made by examining Table VIII.

(1) The calculated values converges to a few hundredths of a wavenumber by comparing the fourth and sixth order results. The convergence is better than in the case of CF₄. The contribution due to the V' terms is small. For example, the largest contribution to the fundamentals is 0.0016 cm⁻¹ for (0001; F_2) band of ²⁸SiF₄ using sixth order CVPT. This is smaller than that of the ¹²CF₄ since SiF₄ is heavier.

(2) The agreement between calculation and experiment is rather good with a RMSD of 0.59, 0.74 and 0.73 cm⁻¹ at second, fourth and sixth order CVPT, respectively. It should be noted that the slightly better agreement at the second order is fortuitous as the second order results still do not converge. On the whole, the CVPT results are remarkable in spite of the fact that the CVPT calculation uses the *ab initio* force field where only four diagonal quadratic force constants were optimized to the four experimental fundamentals. The success of the perturbation calculation is mainly due to the small anharmonicity for this relatively heavy-atom molecule (cfr. the $\omega_i - \nu_i$ values in Tables I and II).

(3) In Table VIII our calculations are compared with those of the U(2) algebraic model.³⁶ Although the U(2) algebraic model yields a similar RMSD (0.79 cm⁻¹), nine parameters were employed to fit the sixteen vibrational data. A fit of similar quality is also reported by the same authors³⁷ using a boson realization model. However, in both models the polyad number $N = v_1 + v_2 + v_3 + v_4$ is used and the Fermi resonance $\omega_1 \approx 2\omega_4$ is considered in neither.

(4) It can be seen from Table VIII that the resonance $\omega_1 \approx 2\omega_4$ is most prominent. The other two resonances are also considered to ensure good convergence at sixth order. To neglect these two resonances will considerably reduce the size of the matrix associated with a given value of the quantum number, so that high overtone states can be relatively easier to calculate. The lack of strong resonance in ν_3 and its overtone ensures that the isolated treatment of the ν_3 overtone is a good approximation.³⁹ Patterson and Pine has done such a calculation, and find the Cartesian basis is much inferior to the angular momentum basis due to the near degeneracy of the Cartesian basis. This near degeneracy is specific to SiF₄ since the relation $G_{33} = 8T_{33}$ is nearly satisfied.³⁹ However, as far as the calculation of the vibrational energy is concerned, both basis sets can be used.

The partial set of empirically derived anharmonicity constants (Table VII) of McDowell *et al.*³⁵ agrees fairly well with our computed values, while their derived harmonic frequencies agree to within their stated uncertainty with our own best values.

VI. CONCLUSIONS

The anharmonic vibrational spectra of the all-heavy atom spherical tops CF_4 and SiF_4 have been treated by the combination of an accurate *ab initio* anharmonic force field and high-order canonical Van Vleck perturbation theory (CVPT).

The anharmonic part of the potential energy surface is evidently very well described at the CCSD(T)/cc-pVTZ level for CF_4 , and at the CCSD(T)/cc-pVTZ+1 level for SiF_4 , where the '+1' notation refers to the addition of a high-exponent *d* function on second-row atoms. The harmonic frequencies of CF_4 exhibit substantial errors at the CCSD(T)/cc-pVTZ level, which disappear upon addition of diffuse functions to the basis set.

The force field is subsequently slightly refined by adjusting the equilibrium geometry and the diagonal quadratic force constants (in this case, five parameters) such that a standard second-order rovibrational perturbation theoretical treatment reproduces the experimental fundamentals and ground-state rotational constants of the molecules. The adjustments involved are fairly minor.

These force fields were then used as input for CVPT calculations up to sixth order inclusive. While agreement with experiment is fortuitously quite good at second order in CVPT, consistent convergence to 0.1 cm^{-1} in the energy level is only achieved at sixth order. However, the truncation error at second order is much less significant than in the case of hydride systems like CH₄ and SiH₄.

RMS deviation (RMSD) between computed (sixth order CVPT) and observed energy levels with the adjusted ab initio potentials is 0.19 cm^{-1} for CF₄ and 0.73 cm^{-1} for SiF₄, the latter to some extent reflecting the lesser accuracy of the experimental data. In the case of CF₄, improvement with experiment could be somewhat further improved (to RMSD= 0.08 cm^{-1}) by re-adjusting both diagonal and off-diagonal quadratic force constants to the complete set of experimental vibrational level information. Experimentation with refinement of additional force field parameters yielded no further improvement.

For CF₄, three resonances, Eqs. (23)–(25), were considered, of which only one ($\omega_3 \approx 2\omega_4$) is important. This shows that an isolated ν_3 overtone ladder model is inadequate. For SiF₄, three resonances, Eqs. (19)–(21), were considered, of which only one ($\omega_1 \approx 2\omega_4$) is important.

An improved approach for symmetrizing combination bands in the Cartesian basis for the T_d group is proposed. We also demonstrate how anharmonic spectroscopic constants X_{ij} , G_{ij} , T_{ij} , S_{ij} can be extracted from the second-order CVPT transformed Hamiltonian (in curvilinear internal coordinates) for T_d molecules. In the absence of resonances, the results are identical to those obtained by standard second-order perturbation theory in rectilinear normal coordinates. Differences occur when the constants are being deperturbed for Fermi resonances. Accurate sets of quartic spectroscopic constants for the isotomomers of CF₄ and SiF₄ are obtained. Agreement with previously published empirically derived sets of anharmonicity constants is fairly good for SiF₄, but less satisfactory for CF₄.

In order to stimulate further research on these molecules, sixth-order CVPT energy level predictions up to polyad number N = 24 for $\{12,13,14\}$ CF₄ and $\{28,29,30\}$ SiF₄ have been made available on the World Wide Web at the Uniform Resource Locator (URL) http://theochem.weizmann.ac.il/web/papers/cf4sif4.html. The force fields themselves are available in machine-readable form at the same URL.

ACKNOWLEDGMENTS

Prof. M. S. Child is thanked for helpful discussions on the basis set symmetrization. JM is a Yigal Allon Fellow and the incumbent of the Helen and Milton A. Kimmelman Career Development Chair, acknowledges support from the Minerva Foundation, Munich, Germany, and would like to thank Dr. Timothy J. Lee (NASA Ames Research Center) for helpful discussions and Prof. Jean Demaison (Université de Lille I) for a preprint of Ref. 42. XGW thanks the Royal Society KC Wong Fellowship for support.

¹ Kirk-Othmer encyclopedia of chemical technology, 4th Ed., R. E. Kirk, D. F. Othmer, J. I. Kroschwitz, Eds. (Wiley, New York, 1998)

² H.-G. Mack and H. Oberhammer, J. Chem. Phys. 87, 2158 (1987) and references therein

- ³ J.-H. Kim, S.-H. Seo, S.-M. Yun, H.-Y. Chang, K.-M. Lee, and C.-K. Choi, J. Electrochem. Soc. 143, 2990 (1996) and references therein.
- ⁴ J. Ding and N. Hershkowitz, Appl. Phys. Lett. **68**, 1619 (1996)
- ⁵ P. G. M. Sebel, L. J. F. Hermans, and H. C. W. Beijerinck, J. Vac. Sci. Tech. A **17**, 755 (1999); see also T. Lagarde, J. Pelletier, and Y. Arnal, J. Vac. Sci. Tech. A **17**, 118 (1999)
- ⁶ P. Francis, C. Chaffin, A. Maciejewski, and C. Oppenheimer, Geophys. Res. Lett. 23, 249 (1996)
- ⁷ K. Tanaka, S. Isomura, H. Kaetsu, Y. Yatsurugi, M. Hashimoto, K. Togashi, and S. Arai, Bull. Chem. Soc. Japan **69**, 493 (1996)
- ⁸ B. Monostori and A. Weber, J. Chem. Phys. **33**, 1867 (1960); A. Maki, E. K. Plyler, and R. Thibault, J. Chem. Phys. **37**, 1899 (1960); P. J. H. Woltz and A. H. Nielsen, J. Chem. Phys. **20**, 307 (1952);
- ⁹ L. H. Jones, B. J. Krohn, and R. C. Kennedy, J. Mol. Spectrosc. **70**, 288 (1978)
- ¹⁰ L. H. Jones, C. Kennedy, and S. Ekberg, J. Chem. Phys. **69**, 833 (1978).
- ¹¹ A. C. Jeannotte II, D. Legler, and J. Overend, Spectrochim. Acta A 29, 1915 (1973)
- ¹² J. J. Tiee and C. Wittig, Appl. Phys. Lett. **30**, 420 (1977); J. Appl. Phys. **49**, 61 (1978)
- ¹³ P. Esherick, A. Owyoung, and C. W. Patterson, J. Molec. Spectrosc. **86**, 250 (1981).
- ¹⁴ J. E. Lolck, J. Raman Spectrosc. **11**, 294 (1981).
- ¹⁵ A. Tabyaoui, B. Lavorel, R. Saint-Loup, and M. Rötger, J. Raman Spectrosc. **25**, 255 (1994).
- ¹⁶ C. W. Patterson, R. S. McDowell, N. G. Nereson, R. F. Begley, H. W. Galbraith, and B. J. Krohn, J. Molec. Spectrosc. 80, 71 (1980).
- ¹⁷ G. Poussigue, G. Tarrago, and A. Valentin, J. Phys. B. **47**, 1155 (1986)
- ¹⁸ G. Tarrago, G. Poussigue, M. Dang-Nhu, and J. Kauppinen, J. Mol. Spectrosc. **86**, 232 (1981)
- ¹⁹ R. S. McDowell, M. J. Reisfeld, H. W. Galbraith, B. J. Krohn, H. Flicker, R. C. Kennedy, J. P. Aldridge, and N. G. Nereson, J. Molec. Spectrosc. 83, 440 (1980).
- ²⁰ A. S. Pine, J. Molec. Spectrosc. **96**, 395 (1982)
- ²¹ M. Dang-Nhu, G. Graner, and G. Guelachvili, J. Mol. Spectrosc. **85**, 358 (1981)
- ²² M. Takami, J. Chem. Phys. **73**, 2665 (1980)
- ²³ T. Gabard, A. Nikitin, J. P. Champion, G. Pierre, and A. S. Pine, J. Molec. Spectrosc. **170**, 431 (1995).
- ²⁴ T. Gabard, L. Pierre, and M. Takami, Mol. Phys. **85**, 735 (1995)
- ²⁵ J. L. Duncan and I. M. Mills, Spectrochim. Acta **20**, 1089 (1964)
- ²⁶ A. A. Chalmers and D. C. McKean, Spectrochim. Acta **22**, 251 (1966)
- ²⁷ A. C. Jeannotte II, C. Marcott, and J. Overend, J. Chem. Phys. **65**, 2076 (1978)
- ²⁸ S. Brodersen, J. Mol. Spectrosc. **145**, 331 (1991)
- ²⁹ S. G. Larsen and S. Brodersen, J. Molec. Spectrosc. **157**, 220 (1993).
- ³⁰ V. Boujut, F. Michelot, and C. Leroy, Mol. Phys. **93**, 879 (1998).
- ³¹ V. G. Tyuterev, Y. L. Babikov, S. A. Tashkun, V. I. Perevalov, A. Nikitin, J. P. Champion, C. Wenger, G. Pierre, J. C. Hilico, and M. Loëte, J. Quant. Spec. Radiat. Transfer **52**, 459 (1994) based on Ref. 24
- ³² W. Maring, J. P. Toennies, R. G. Wang, and H. B. Levene, J. Chem. Phys. **103**, 1333 (1995).
- ³³ R. K. Heenan, Ph.D. Thesis (U. of Reading, UK, 1979)
- ³⁴ K. T. Hecht, J. Molec. Spectrosc. 5, 355 (1960).
- ³⁵ R. S. McDowell, M. J. Reisfeld, C. W. Patterson, B. J. Krohn, M. C. Vasquez, and G. A. Laguna, J. Chem. Phys. 77, 4337 (1982).
- ³⁶ X.-W. Hou, S.-H. Dong, M. Xie, and Z.-Q. Ma, Chem. Phys. Lett. **283**, 174 (1998).
- ³⁷ X.-W. Hou, M. Xie, S.-H. Dong, and Z.-Q. Ma, Ann. Phys. (NY) **203**, 340 (1998).
- ³⁸ F. Iachello, Chem. Phys. Lett. **78**, 581 (1981); F. Iachello and R. D. Levine, J. Chem. Phys. **77**, 3046 (1982)
- ³⁹ C. W. Patterson and A. S. Pine, J. Mol. Spectrosc. **96**, 404 (1982)
- ⁴⁰ M. Takami and H. Kuze, J. Chem. Phys. **78**, 2204 (1983).
- ⁴¹ L. Jörissen, H. Prinz, W. A. Kreiner, C. Wenger, G. Pierre, G. Magerl, W. Schupita, Can. J. Phys. 67, 532 (1989).
- ⁴² J. Breidung, J. Demaison, L. Margules, and W. Thiel, Chem. Phys. Lett., submitted ("Equilibrium Structure of SiF4"); preprint communicated to authors
- ⁴³ J. M. L. Martin and P. R. Taylor, J. Phys. Chem. A **103**, 4427 (1999).
- ⁴⁴ G. K. Johnson, J. Chem. Thermodyn. **18**, 801 (1986)
- ⁴⁵ T. J. Lee, J. M. L. Martin, and P. R. Taylor, J. Chem. Phys. **102**, 254 (1995)
- ⁴⁶ S. Carter, H. M. Shnider, and J. M. Bowman, J. Chem. Phys. **110**, 8417 (1999)
- ⁴⁷ E. Venuti, L. Halonen, and R. G. Della Valle, J. Chem. Phys. **110**, 7339 (1999).
- ⁴⁸ X.-G. Wang and E. L. Sibert, J. Chem. Phys. **111**, 4510 (1999).
- ⁴⁹ H.-J. Werner and P. J. Knowles, MOLPRO 97.3, a package of *ab initio* programs, with contributions from J. Almlöf, R. D. Amos, A. Berning, D. L. Cooper, M. J. O. Deegan, A. J. Dobbyn, F. Eckert, S. T. Elbert, C. Hampel, R. Lindh, A. W. Lloyd, W. Meyer, A. Nicklass, K. A. Peterson, R. M. Pitzer, A. J. Stone, P. R. Taylor, M. E. Mura, P. Pulay, M. Schütz, H. Stoll and T. Thorsteinsson.
- ⁵⁰ G. D. Purvis III and R. J. Bartlett, J. Chem. Phys. **76**, 1910 (1982)

- ⁵¹ K. Raghavachari, G. W. Trucks, J. A. Pople, and M. Head-Gordon, Chem. Phys. Lett. **157**, 479 (1989)
- ⁵² P. J. Knowles, C. Hampel, and H. J. Werner, J. Chem. Phys. **99**, 5219 (1993)
- ⁵³ T. J. Lee and G. E. Scuseria, in *Quantum mechanical electronic structure calculations with chemical accuracy*, Ed. S. R. Langhoff (Kluwer Academic Publishers, Dordrecht, The Netherlands, 1995).
- ⁵⁴ T. J. Lee and P. R. Taylor, Int. J. Quantum Chem. Symp. 23, 199 (1989)
- ⁵⁵ D. Papoušek and M. R. Aliev, Molecular Vibrational-Rotational Spectra (Elsevier Scientific Publishing Company,: New York 1982).
- ⁵⁶ A. Willetts, J. F. Gaw, W. H. Green Jr., and N. C. Handy, SPECTRO 1.0, a second-order rovibrational perturbation theory program (University Chemical Laboratory, Cambridge, UK, 1989)
- ⁵⁷ J. F. Gaw, A. Willetts, W. H. Green, and N. C. Handy, in Advances in molecular vibrations and collision dynamics (ed. J. M. Bowman), JAI Press, Greenwich, CT, 1990.
- ⁵⁸ D. P. Hodgkinson, R. K. Heenan, A. R. Hoy, and A. G. Robiette, Mol. Phys. 48, 193 (1983).
- ⁵⁹ This is fairly trivially done by taking apart the compound fractions in Eq.(3.20) of Ref. 58, then inserting a test that skips the appropriate one among the four affected terms if the respective denominator corresponds to the resonant interaction. Alternatively, the reciprocal denominators may be precomputed and stored in a 3-dimensional array, and any that correspond to resonant interactions set to zero before the actual spectroscopy routines are entered, similar to J. M. L. Martin and P. R. Taylor, Spectrochim. Acta A **53**, 1039 (1997).
- ⁶⁰ T. H. Dunning, Jr., J. Chem. Phys. **90**, 1007 (1989)
- ⁶¹ D. E. Woon and T. H. Dunning Jr., J. Chem. Phys. **98**, 1358 (1993).
- ⁶² J. M. L. Martin and O. Uzan, Chem. Phys. Lett. **282**, 16 (1998)
- ⁶³ J. M. L. Martin, J. Chem. Phys. **108**, 2791 (1998)
- ⁶⁴ J. M. L. Martin, Spectrochim. Acta A 55, 709 (1999) (special issue "Theoretical spectroscopy: state of the science")
- ⁶⁵ R. A. Kendall, T. H. Dunning Jr., and R. J. Harrison, J. Chem. Phys. 96, 6796 (1992).
- ⁶⁶ J. M. L. Martin and P. R. Taylor, Chem. Phys. Lett. **225**, 473 (1994).
- ⁶⁷ J. M. L. Martin, J. P. François, and R. Gijbels, J. Phys. Chem. **98**, 11394 (1994).
- ⁶⁸ J. M. L. Martin, Chem. Phys. Lett. **292**, 411 (1998)
- ⁶⁹ J. M. L. Martin, K. K. Baldridge, and T. J. Lee, Mol. Phys. **xx**, yyy (1999)
- ⁷⁰ H. M. Pickett, J. Chem. Phys. **56**, 1715 (1972).
- ⁷¹ E. L. Sibert, J. Chem. Phys. **88**, 4378 (1988).
- ⁷² E. L. Sibert, Comp. Phys. Comm. **51**, 149 (1988).
- ⁷³ R. G. Della Valle, L. Halonen, and E. Venuti, J. Comput. Chem., (in press).
- ⁷⁴ L. Halonen and M. S. Child, Comp. Phys. Comm. **51**, 173 (1988).
- ⁷⁵ L. Halonen, J. Chem. Phys. **106**, 831 (1997).
- ⁷⁶ J. S. Griffith, *The Theory of Transition-Metal Ions*, Cambridge University Press, Cambridge, (1961), Table A20 of Appendix 2.
- ⁷⁷ Y. Tanabe and S. Sugano, J. Phys. Soc. Jap. **9**, 753 (1954).
- ⁷⁸ L. Pauling and E. B.Wilson, Introduction to Quantum Mechanics with Applications to Chemistry (Dover Publications, Inc.: New York 1935).
- ⁷⁹ W. H. Shaffer, Rev. Mod. Phys. **16**, 245 (1944).
- ⁸⁰ C. Cohen-Tannoudji, B. Diu, and F. Laloë, Quantum Mechanics, vol 1, pp 727-741, Wiley, New York, (1977).
- ⁸¹ Y. Pak, E. L. Sibert III, and R. C. Woods, J. Chem. Phys. **107**, 1717 (1997).
- ⁸² MAPLE V Release 4, Waterloo Maple Software, Waterloo, Ontario, 1981-1994.
- ⁸³ R. Lemus and A. Frank, J. Chem. Phys. **101**, 8321 (1994).
- ⁸⁴ Z.-Q. Ma, X.-W. Hou, and M. Xie, Phys. Rev. A 53, 2173 (1996).
- ⁸⁵ A. Frank, R. Lemus, F. Pérez-Bernal, and R. Bijker, J. Mol. Spectrosc. **196**, 329 (1999).
- ⁸⁶ A. B. McCoy and E. L. Sibert III, J. Chem. Phys. **105**, 459 (1996); E. L. Sibert III and A. B. McCoy, J. Chem. Phys. **105**, 469 (1996).
- ⁸⁷ H. A. Jahn, Proc. Roy. Soc. A168, 469 (1938).
- ⁸⁸ H. A. Jahn, Ann. Phys. Lpz. **23**, 529 (1935).
- ⁸⁹ H. A. Jahn, Proc. Roy. Soc. **A168**, 495 (1938).
- ⁹⁰ D. L. Gray and A. G. Robiette, Mol. Phys. **37**, 1901 (1979).
- ⁹¹ I. M. Mills, Mol. Phys. 1, 107 (1958), Fig. 1 of this work is inconsistent with its definition of symmetry coordinates.
- ⁹² A. G. Robiette, D. L. Gray, and F. W. Birss, Mol. Phys. **32**, 1591 (1976).
- ⁹³ A. B. McCoy and E. L. Sibert, Mol. Phys. **77**, 697 (1992).
- ⁹⁴ A. B. McCoy and E. L. Sibert, J. Chem. Phys. **95**, 3476 (1991).
- ⁹⁵ A. B. McCoy, D. C. Burleigh, and E. L. Sibert, J. Chem. Phys. **95**, 7449 (1991).
- ⁹⁶ K. L. Mardis and E. L. Sibert, J. Molec. Spectrosc. **187**, 167 (1998).

	cc-pVTZ unadj.			aug-cc-pVTZ	cc-pVTZ adjusted			Expt. ³⁵		
	r_e	r_0	r_z	r_e	r_e	r_0	r_z	r_e	r_0	
	1.31919	1.32310	1.32389	1.32112	1.31526	1.31925	1.32004	$1.3151(17)^a$	$1.319247(1)^{b}$	
i	ω_i	$ u_i$	$\omega_i - \nu_i$	ω_i	ω_i	$ u_i$	$\omega_i - \nu_i$	ω_i	$ u_i$	
1	922.80	910.75	12.06	915.2	921.57	909.07	12.50		$909.0720(1)^{13}$	
2	440.00	435.59	4.40	435.2	439.91	435.40	4.51		$435.399(10)^{14}$	
3	1322.25	1303.15	19.10	1301.3	1303.01	1283.66	19.35		$1283.66429(12)^{24}$	
4	638.81	632.22	6.59	630.4	637.89	631.06	6.84		$631.05890(13)^{24}$	

TABLE I. Computed (CCSD(T)) and observed bond distances (Å), harmonic and fundamental frequencies (cm⁻¹) for ${}^{12}CF_4$

^a Ref. 28.

^b From $B_0 = 0.19118709(32) \text{ cm}^{-1}.^{23}$

TABLE II. Computed (CCSD(T)) and observed bond distances (Å), harmonic and fundamental frequencies (cm⁻¹) for ${}^{28}SiF_4$

	$^{\rm cc-}_{\rm pVTZ}$	aug-cc- pVTZ+1	cc-pVTZ+1 unadj.			cc-pV	/TZ+1 adjı	$Expt.^{35}$		
	r_e	r_e	r_e	r_0	r_z	r_e	r_0	r_z	r_e	r_0
	1.56949	1.56332	1.56134	1.56368	1.56453	1.55182	1.55404	1.55489	$1.5524(8)^a$	1.55404^{b}
i	ω_i	ω_i	ω_i	$ u_i$	$\omega_i - \nu_i$	ω_i	$ u_i$	$\omega_i - \nu_i$	ω_i	$ u_i $
1	794.9	794.1	797.86	792.19	5.67	806.10	800.60	5.50	807.1(12)	800.6
2	259.8	258.8	263.18	262.13	1.05	265.20	264.20	1.00	267(3)	264.2
3	1036.5	1029.8	1037.49	1024.31	13.18	1044.04	1031.40	12.64	1044.2(12)	1031.3968
4	384.3	382.7	387.61	386.67	0.94	389.31	388.44	0.87	389.8(9)	388.4448

^a Ref. 42.

^b From Ref. 41, $B_0=0.137780439(92)$ cm⁻¹. In older work^{35,39}, $B_0=0.13676(3)$ cm⁻¹ and hence $r_0=1.55982(17)$ Å.

TABLE III. Quadratic, cubic and quartic force constants $(aJ/Å^m radian^n)$ for SiF₄ and CF₄

	SiF_4	CF_4		SiF_4	CF_4		SiF_4	CF_4
F_{11}	7.27355	9.50679	F_{22}	0.63194	1.24913	F_{44}	6.40971	6.13519
F_{74}	-0.34328	-1.03126	F_{77}	1.06468	1.83767	F_{111}	-20.29450	-32.46123
F_{221}	-0.69851	-2.19918	F_{441}	-18.73201	-23.71962	F_{741}	0.56578	2.57073
F_{771}	-0.97378	-3.04180	F_{222}	-0.53376	-1.11062	F_{662}	-0.60404	-2.24497
F_{962}	0.61671	2.05526	F_{992}	-1.01287	-2.25675	F_{654}	-18.17256	-18.94290
F_{954}	-0.17620	-0.38638	F_{984}	-0.01658	0.04702	F_{987}	0.86987	1.30646
F_{1111}	50.47669	89.63168	F_{2211}	1.26945	4.44699	F_{4411}	48.94593	71.68647
F_{7411}	-0.80741	-5.46450	F_{7711}	1.53293	5.89090	F_{2221}	0.76478	2.29325
F_{6621}	1.16338	5.73913	F_{9621}	-1.08209	-4.08511	F_{9921}	1.46146	4.33621
F_{6541}	48.23448	60.63713	F_{9541}	0.40465	1.24159	F_{9841}	0.15861	0.26676
F_{9871}	-0.63502	-1.85550	F_{2222}	0.76756	2.44329	F_{6622}	0.76815	3.16961
F_{6633}	-0.46055	-1.86075	F_{9622}	-1.12641	-3.26978	F_{9633}	0.03132	0.14651
F_{9922}	2.11522	5.37994	F_{9933}	0.39929	0.77598	F_{9542}	0.43597	1.12997
F_{8762}	0.01923	0.04530	F_{4444}	50.01306	72.17993	F_{5544}	48.79060	59.28204
F_{7444}	-0.85975	-6.35131	F_{8544}	0.39771	4.59345	F_{7744}	1.37066	5.70508
F_{8754}	0.02280	-0.04550	F_{8844}	-0.53371	-1.07258	F_{7774}	-2.21128	-6.91109
F_{8874}	-0.14830	-0.36797	F_{7777}	4.58572	11.28388	F_{8877}	2.18342	3.67611

wavefunction.					
$ 2^0; A_1\rangle =$	$+\frac{\sqrt{2}}{2} 20\rangle$	$+\frac{\sqrt{2}}{2} 02\rangle$			
$ 2^2; E_a\rangle =$	$-\frac{\sqrt{2}}{\sqrt{2}} 20\rangle$	$+\frac{\sqrt{2}}{\sqrt{2}} 02\rangle$			
$ 2^2; E_b\rangle =$	$+1 11\rangle$	2 1 7			
$ 3^3; A_1\rangle =$	$+\frac{\sqrt{1}}{2} 30\rangle$	$-\frac{\sqrt{3}}{2} 12\rangle$			
$ 3^3; A_2\rangle =$	$-\frac{\sqrt{1}}{2} 03\rangle$	$+\frac{\sqrt{3}}{2} 21\rangle$			
$ 3^1; E_a\rangle =$	$+\frac{\sqrt{3}}{2} 30\rangle$	$+\frac{\sqrt{1}}{2} 12\rangle$			
$ 3^1; E_b\rangle =$	$+\frac{\sqrt{3}}{2} 03\rangle$	$+\frac{\sqrt{1}}{2} 21\rangle$			
$ 4^0; A_1\rangle =$	$+\frac{\sqrt{6}}{4} 40\rangle$	$+\frac{\sqrt{6}}{4} 04\rangle$	$+\frac{\sqrt{4}}{4} 22\rangle$		
$ 4^2; E_a\rangle =$	$+\frac{\sqrt{2}}{2} 40\rangle$	$-\frac{\sqrt{2}}{2} 04\rangle$			
$ 4^2; E_b\rangle =$	$-\frac{\sqrt{2}}{2} 31\rangle$	$-\frac{\sqrt{2}}{2} 13\rangle$	_		
$ 4^4; E_a\rangle =$	$+\frac{\sqrt{2}}{4} 40\rangle$	$+\frac{\sqrt{2}}{4} 04\rangle$	$-\frac{\sqrt{12}}{4} 22\rangle$		
$ 4^4; E_b\rangle =$	$+\frac{\sqrt{2}}{2} 31\rangle$	$-\frac{\sqrt{2}}{2} 13\rangle$			
$ 5^3; A_1\rangle =$	$+\frac{\sqrt{5}}{4} 50\rangle$	$-\frac{\sqrt{2}}{4} 32\rangle$	$-\frac{\sqrt{9}}{4} 14\rangle$		
$ 5^3; A_2\rangle =$	$-\frac{\sqrt{5}}{4} 05\rangle$	$+\frac{\sqrt{2}}{4} 23\rangle$	$+\frac{\sqrt{9}}{4} 41\rangle$		
$ 5^1; E_a\rangle =$	$\frac{\sqrt{10}}{4} 50\rangle$	$+\frac{\sqrt{4}}{4} 32\rangle$	$+\frac{\sqrt{2}}{4} 14\rangle$		
$ 5^1; E_b\rangle =$	$\frac{\sqrt{10}}{4} 05\rangle$	$+\frac{\sqrt{4}}{4} 23\rangle$	$+\frac{\sqrt{2}}{4} 41\rangle$		
$ 5^5; E_a\rangle =$	$\frac{\sqrt{1}}{4} 50\rangle$	$-\frac{\sqrt{10}}{4} 23\rangle$	$+\frac{\sqrt{5}}{4} 14\rangle$		
$ 5^5; E_b\rangle =$	$-\frac{\sqrt{1}}{4} 05\rangle$	$+\frac{\sqrt{10}}{4} 23\rangle$	$-\frac{\sqrt{5}}{4} 41\rangle$	-	
$ 6^0; A_1\rangle =$	$+\frac{\sqrt{5}}{4} 60\rangle$	$+\frac{\sqrt{5}}{4} 06\rangle$	$+\frac{\sqrt{3}}{4} 42\rangle$	$+\frac{\sqrt{3}}{4} 24\rangle$	
$ 6^3; A_1\rangle =$	$+\frac{\sqrt{2}}{8} 60\rangle$	$-\frac{\sqrt{2}}{8} 06\rangle$	$-\frac{\sqrt{30}}{8\pi} 42\rangle$	$+\frac{\sqrt{30}}{8} 24\rangle$	
$ 6^3; A_2\rangle =$	$+\frac{\sqrt{3}}{4} 51\rangle$	$-\frac{\sqrt{3}}{4} 15\rangle$	$+\frac{\sqrt{7}}{4} 33\rangle$	/2	
$ 6^2; E_a\rangle =$	$+\frac{\sqrt{30}}{8} 60\rangle$	$-\frac{\sqrt{30}}{8\pi} 06\rangle$	$+\frac{\sqrt{2}}{8} 42\rangle$	$-\frac{\sqrt{2}}{8} 24\rangle$	
$ 6^2; E_b\rangle =$	$-\frac{\sqrt{5}}{4} 51\rangle$	$-\frac{\sqrt{5}}{4} 15\rangle$	$-\frac{\sqrt{6}}{4} 33\rangle$		
$ 6^4; E_a\rangle =$	$+\frac{\sqrt{3}}{4} 60\rangle$	$+\frac{\sqrt{3}}{4} 06\rangle$	$-\frac{\sqrt{5}}{4} 42\rangle$	$-\frac{\sqrt{5}}{4} 24\rangle$	
$ 6^4; E_b\rangle =$	$+\frac{\sqrt{2}}{2} 51\rangle$	$\frac{-\frac{\sqrt{2}}{2} 15\rangle}{\frac{\sqrt{1}}{2}}$	15.	/27	
$ 7^{3}; A_{1}\rangle =$	$+\frac{\sqrt{21}}{8} 70\rangle$	$-\frac{\sqrt{1}}{8} 52\rangle$	$-\frac{\sqrt{15}}{8} 34\rangle$	$-\frac{\sqrt{27}}{8} 16\rangle$	
$ 7^3; A_2\rangle =$	$-\frac{\sqrt{21}}{8} 07\rangle$	$+\frac{\sqrt{1}}{8} 25\rangle$	$+\frac{\sqrt{15}}{8} 43\rangle$	$+\frac{\sqrt{27}}{8} 61\rangle$	
$ 7^{1}; E_{a}\rangle =$	$+\frac{\sqrt{35}}{8} 70\rangle$	$+\frac{\sqrt{15}}{8} 52\rangle$	$+\frac{\sqrt{3}}{8} 34\rangle$	$+\frac{\sqrt{5}}{8} 16\rangle$	
$ \mathcal{T}^{1}; E_{b}\rangle =$	$+\frac{\sqrt{33}}{8} 07\rangle$	$+\frac{\sqrt{15}}{8} 25\rangle$	$+\frac{\sqrt{3}}{8} 43\rangle$	$+\frac{\sqrt{3}}{8} 61\rangle$	
$ 7^{\circ}; E_a\rangle =$	$+\frac{\sqrt{3}}{8} 70\rangle$	$-\frac{\sqrt{27}}{8} 52\rangle$	$-\frac{\sqrt{5}}{8} 34\rangle$	$+\frac{\sqrt{25}}{8} 10\rangle$	
$ \mathcal{T}^{\circ}; E_b\rangle =$	$-\frac{\sqrt{3}}{8} 07\rangle$	$+\frac{\sqrt{21}}{8} 25\rangle$	$+\frac{\sqrt{3}}{8} 43\rangle$	$-\frac{\sqrt{23}}{8} 61\rangle$	
$ T; E_a\rangle =$	$+\frac{\sqrt{3}}{8} 10\rangle$	$-\frac{\sqrt{21}}{8} 52\rangle$	$+\frac{\sqrt{35}}{8} 34\rangle$	$-\frac{\sqrt{3}}{8} 10\rangle$	
$ T; E_b\rangle =$	$-\frac{1}{8} 07\rangle$	$+\frac{1}{8} 23\rangle$	$-\frac{\sqrt{8}}{8} 43\rangle$	$+\frac{1}{8} 01\rangle$	$\sqrt{36} 4 \rangle$
$ 8^{\circ}; A_1\rangle =$	$+\frac{16}{16} 80\rangle$	$+\frac{16}{16} 08\rangle$	$+\frac{16}{16} 02\rangle$	$+\frac{16}{16} 20\rangle$	$+\frac{16}{16} 44\rangle$
$ 0; A_1\rangle = 0^6, A_1\rangle$	$+\frac{1}{4} 80\rangle$	$-\frac{4}{4} 08\rangle$	$-\frac{4}{\sqrt{14}} 02\rangle$	$+\frac{1}{4} 20\rangle$	
$ 0, A_2\rangle = 8^2, F\rangle =$	$+\frac{8}{\sqrt{7}} (1) $	$+\frac{8}{\sqrt{7}}$	$-\frac{8}{\sqrt{1}} 00\rangle$	$-\frac{8}{\sqrt{1}}$	
$ 0\rangle, E_a/=$ $ 8^2, E_b\rangle =$	$+\frac{4}{4} 00\rangle$	$-\frac{4}{\sqrt{14}}$	$+\frac{4}{4} 02\rangle$	$-\frac{4}{\sqrt{18}}$	
$ 0\rangle, E_b/=$ $ 8^4 \cdot E\rangle -$	$-\frac{8}{8}$ (1) $\pm \sqrt{14}$ (80)	$-\frac{8}{8} 11\rangle$ $\pm \sqrt{14} 08\rangle$	$-\frac{8}{8}$ [55]	$-\frac{\sqrt{8}}{8}$	$-\sqrt{20} _{\Lambda\Lambda}$
$ 0\rangle, D_a/=$ $ 8^4 \cdot E_1\rangle =$	$\pm \frac{\sqrt{7}}{171}$	$-\frac{\sqrt{7}}{117}$	$\frac{-\frac{8}{8}}{153}$	$-\frac{1}{8}$ [20/ $-\frac{\sqrt{1}}{25}$]	
$ 0\rangle, E_b/=$ $ 8^8 \cdot E\rangle =$	$+\frac{\sqrt{2}}{4} 11\rangle$	$-\frac{1}{4}$ 17/ + $\frac{\sqrt{2}}{108}$	$-\frac{\sqrt{56}}{62}$	$-\frac{4}{56} _{26}$	$+ \sqrt{140} \Lambda \Lambda \rangle$
$ 0\rangle, D_a/=$ $ 8^8 \cdot E_1\rangle =$	$+\frac{16}{16} 00\rangle$ $-\frac{\sqrt{1}}{71} 71\rangle$	$\pm \frac{16}{16}$ 00/ $\pm \sqrt{1}$ 17	$-\frac{16}{16}$ 02/ $\pm \sqrt{7}$ 53\	$-\frac{16}{\sqrt{7}}$	$\pm \frac{16}{16}$ (44)
$ 0, L_{b} =$	4 / 1/	4 4 /	4-100/	4 00/	

TABLE IV. The relationships of the angular momentum basis $|v^l\rangle$ and the Cartesian basis $|n_x n_y\rangle$ for the overtone of the doubly degenerated modes (ν_2). See Eqs. (10) and (11) for definition of A_1/A_2 and E_a/E_b symmetry wavefunction.

Force Constant	ab initio	ab initio	\mathbf{Fitted}^{c}	σ^d
	raw^a	$\operatorname{adjusted}^{b}$		
r_e	1.31919	1.31526	1.31526	
F_{11}	9.54112	9.50679	9.50711	0.0%
F_{22}	1.25733	1.24913	1.12477	-10.0%
F_{33}	6.32376	6.13519	6.18043	+0.7%
F_{34}	-1.03160	-1.03126	-1.04559	-1.4%
F_{44}	1.84886	1.83767	1.82662	-0.6%

TABLE V. Comparison of *ab initio^a* and fitted quadratic force constants for CF_4 . (Units are consistent with aJ, Å, and radian.).

 a Unadjusted CCSD(T)/cc-pVTZ values

 b Bond distance and diagonal quadratics adjusted to reproduce experimental r_{0} and ν_{i}

 c bond distance held constants; all quadratic force constants refined in fit againt complete experimental data set.

 d Relative deviation of the fitted force constants from the adjusted *ab initio* force constants.

N^a	Sym	Obs^b	Uncertainty	Ref.	Obs-E(2)	Obs-E(4)	$Obs - E(\overline{6})$	$v_1 v_2 v_3 v_3 v_4$	$(c_{i}^{2})^{d}$
		$^{12}CF_4$							
2	E	435.399	1	13	.227	.100	.103	0100(100%)	
3	F_2	631.0593	1	30	.127	039	031	0001(100%)	
4	A_1	867.90588	1	15	.476	012	003	0200(97%)	1000(2%)
4	A_1	909.0720	1	14	151	044	045	1000(97%)	0200(2%)
5	F_1	1066.6977	1	16	.300	.052	.065	0101(100%)	
5	F_2	1066.1220	1	16	.320	135	112	0101(100%)	
6	A_1	1261.809	10	30	.303	184	148	0002(90%)	
6	E	1262.112	1	30	.219	244	221	0002(100%)	
6	F_2	1260.430	1	30	.186	.173	.188	0002(93%)	0010(7%)
6	F_2	1283.720	1	23	.361	005	002	0010(93%)	0002(7%)
7	F_2	1539.3	10	10	198	251	241	1001(97%)	0201(3%)
8	F_2	1715.8	10	10	225	963	960	0110(91%)	0102(9%)
9	F_2	1889.6^{e}	-	10	.623	.887	.912	0003(84%)	0011(16%)
9	F_2						1893.42	0003(100%)	
9	F_2	1913.2	10	10	2.031	1.050	1.100	0011(84%)	0003(16%)
10	F_2	2168.5	10	10	.674	.604	.621	1002(89%)	1010(9%)
10	F_2	2186.1	10	10	602	636	632	1010(86%)	1002(9%)
11	F_2	2445.59644	1	20	340	.025	.051	2001(94%)	1201(5%)
12	A_1	2553.24(858)	-	23	6.225	3.072	3.215	0020(50%)	0012(41%)
12	E	2570.013	_	23	1.991	1.652	1.654	0020(93%)	0012(7%)
12	F_2	2561.9124	-	23	-2.127	-3.061	-3.067	0020(86%)	0012(13%)
		$^{13}CF_4$							
3	F_2	629.2868	1	19	.270	.110	.118	0001(100%)	
5	F_2	1064.39	10	10	.408	033	011	0101(100%)	
6	F_2	1241.7	10	10	251	.014	.009	0010(91%)	0002(9%)
6	F_2	1259.75	10	10	.775	.138	.160	0002(91%)	0010(9%)
7	F_2	1537.4	10	10	078	116	107	1001(97%)	0201(3%)
8	F_2	1674.7	10	10	099	081	087	0110(93%)	0102(7%)
9	F_2	1867.	100	10	1.074	1.631	1.630	0011(89%)	0003(11%)
9	F_2	$1888.^{e}$	-	10	1.262	.279	.353	0003(100%)	
9	F_2						1889.37	0003(89%)	0011(11%)
10	F_2	2145.	100	10	698	518	530	1010(92%)	1002(5%)
10	F_2	2166.3	10	10	.017	232	200	1002(92%)	1010(5%)
11	F_2	2443.3	10	10	512	.011	.048	2001(94%)	1201(6%)
12	F_2	2477.5	10	10	-3.086	-1.556	-1.565	0020(71%)	0012(26%)
		$^{14}\mathrm{CF}_4$							
3	F_2	627.3490	1	19	.275	.121	.128	0001(100%)	
5	F_2	1062.57	10	10	.449	.019	.041	0101(100%)	
6	F_2	1208.7	10	10	107	055	059	0010(99%)	
6	F_2	1254.95	10	10	.800	.387	.407	0002(99%)	
7	F_2	1535.3	10	10	204	230	221	1001(97%)	0201(3%)
8	F_2	1641.6	10	10	038	239	246	0110(99%)	
9	F_2	1833.4	10	10	1.846	2.023	2.030	0011(98%)	0003(2%)
9	F_2	1881.4^{e}	—	10	.717	212	136	0003(99%)	0011(1%)
9	F_2	<i>.</i>					1882.11	0003(99%)	
10	F_2	2073.7^{f}	10	10	1.721	1.109	1.098	0210(97%)	1010(3%)
10	F_2						2075.77	0210(99%)	
10	F_2	2112.	100	10	683	538	547	1010(97%)	0210(3%)
10	F_2	2161.9	10	10	.398	.218	.246	1002(96%)	0202(3%)
12	F_2	2412.	100	10	-3.208	-2.645	-2.638	0020(97%)	0012(2%)
RMSD					0.0179	0.083	0.083		

TABLE VI. Comparison of experimental and CVPT^a band origins (cm^{-1}) for CF_4 . The fit quadratic force constants of Table V plus the *ab initio* cubic and quartic force contants of Table III are used for the CVPT calculation.

 $\overline{a} N = 4v_1 + 2v_2 + 6v_3 + 3v_4$. This polyad number results from three independent resonances: $\omega_1 \approx 2\omega_2$, $3\omega_2 \approx \omega_3$, and $\omega_3 \approx 2\omega_4$.

^b The last figure is significant unless uncertainty in parenthesis is given otherwise.

^c Uncertainty used in the fit corresponds approximately to the experimental precision. The states excluded from the fit are $3\nu_4$ and $2\nu_2 + \nu_3$ whose multiple F_2 components are unresolved, and the $2\nu_3$ bands of ${}^{12}CF_4$, which differs from experimental results.

^d The largest two components in terms of $v_1v_2v_3v_4$ and their percentage (c_i^2) in the basis, based on the sixth order calculation. Only the components with the percentage larger than 1% are listed.

^e There are two (0003) F_2 states.

 f There are two (0210) F_{2} states.

Constants	$^{12}\mathrm{CF}_4$	$^{12}\mathrm{CF}_4$	$^{12}\mathrm{CF}_4$	$^{13}\mathrm{CF}_4$	$^{14}\mathrm{CF}_4$	$^{28}\mathrm{SiF}_4$	$^{28}\mathrm{SiF}_4$	$^{29}\mathrm{SiF}_4$	$^{30}SiF_4$
		Heenan I ³³	Heenan II ³³				Ref. 35		
ω_1	921.596	910.52	913.25	921.596	921.596	806.101	807.1(12)	806.101	806.101
ω_2	439.665	437.15	435.37	439.665	439.665	265.199	267(3)	265.199	265.199
ω_3	1302.510	1292.96	1302.26	1262.587	1227.629	1044.044	1044.2(12)	1034.820	1026.177
ω_4	637.681	634.35	635.83	635.548	633.406	389.306	389.8(9)	387.793	386.320
X_{11}	-1.109	-0.279	-0.275	-1.109	-1.109	669	-0.57(5)	669	669
X_{12}	911^{a}	0.589	0.474	911	911	317	-0.6(11)	317	317
X_{13}	-6.253	-4.896	-4.954	-5.908	-5.668	-3.573	-3.8(3)	-3.503	-3.444
X_{14}	675	0.608	0.712	928	-1.087	$.109^{c}$	+0.64(7)	.056	.013
X_{22}	339^{a}	-0.284	-0.254	339	339	168	0.0(5)	168	168
X_{23}	-2.372	-0.544	-0.543	-2.315	-2.264	989	-1.5(10)	979	968
X_{24}	004	-0.283	-0.273	.003	.007	.506	0.0(5)	.503	.500
X_{33}	-4.157	-5.642	-5.536	-3.917	-3.713	-2.845	-3.0058(7)	-2.779	-2.717
X_{34}	-3.960^{b}	-1.407	-1.423	-3.756	-3.594	-1.386	-0.5(4)	-1.350	-1.318
X_{44}	129^{b}	-0.606	-0.604	122	114	.165 ^c	-0.22(10)	.163	.161
G_{22}	$.542^{a}$	0.286	0.257	.542	.542	.391	[0]	.391	.391
G_{33}	3.833	4.317	4.241	3.618	3.432	1.799	1.7828(6)	1.746	1.698
G_{34}	518	-1.270	-1.320	442	401	414	≈ -0.5	377	345
G_{44}	$.025^{b}$	0.088	0.084	.025	.025	$.000^{c}$	-0.05(15)	.000	.001
T_{23}	039	0.045	0.044	037	035	101	[0]	098	096
T_{24}	037	-0.006	-0.005	038	039	.005	[0]	.004	.003
T_{33}	.221	0.683	0.670	.221	.222	.196	0.20292(10)	.195	.195
T_{34}	$.080^{b}$	0.391	0.394	.065	.051	133	≈ -0.5	131	129
T_{44}	$.020^{b}$	0.091	0.090	.021	.021	.014	[0]	.014	.013
S_{34}	$.157^{b}$.211	.243	.019	[0]	.027	.033
$K_{1,44}$	0			0	0	2.493^{c}		2.465	2.438
$K_{1,22}$	-3.418^{a}			-3.418	-3.418	0		0	0
$K_{3,44}$	-5.223^{b}			-4.524	-3.941	0		0	0
$K_{1,222}$	0			0	0	0.063		0.063	0.063

TABLE VII. Computed (second order transformed CVPT Hamiltonian) and experimentally derived vibrational spectroscopic constants of SiF_4 and CF_4 (cm⁻¹).

Spectroscopic constants are identical between curvilinear and rectilinear formalisms except for those affected by resonances: ^a Due to $\omega_1 \approx 2\omega_2$ resonance, in rectilinear formalism: $X_{12} = -0.606$; $X_{22} = -0.416$; $G_{22} = 0.619$; and $K_{1,22} = -2.908$ cm⁻¹. ^b Due to $\omega_3 \approx 2\omega_4$ resonance, in rectilinear formalism: $X_{34} = -3.916$; $X_{44} = -0.142$; $G_{34} = -0.550$; $G_{44} = 0.018$; $T_{34} = 0.067$; $T_{44} = 0.024$; $S_{34} = 0.161$; and $K_{3,44} = 5.051$ cm⁻¹. ^c Due to resonance $\omega_1 \approx 2\omega_4$, in rectilinear formalism: $X_{14} = -0.231$; $X_{44} = 0.250$; $G_{44} = -0.085$; and $K_{1,44} = 2.699$ cm⁻¹.

N^a	Sym	Obs^b	Obs-E(2)	Obs-E(4)	Obs-E(6)	$Obs - U(2)^c$	$\nu_1 \nu_2 \nu_3 \nu_4 (c_i^2)^d$	
2	E	264.2(10)	.00	03	03	21	0100(100%)	
3	F_2	388.4448(2)	01	03	03	41	0001(100%)	
6	A_1	800.6(3)	24	30	30	1.03	1000(93%)	0002(6%)
6	F_2	776.3(5)	81	84	83	.97	0002(100%)	
8	F_2	1031.3968(3)	64	76	75	1.72	0010(100%)	
8	E	1064.2(4)	.00	06	06	.01	1100(92%)	0102(7%)
9	F_2	1164.2(2)	.48	.48	.51	.03	0003(90%)	1001(10%)
9	F_2	1189.7(3)	70	83	82	1.07	1001(90%)	0003(10%)
10	F_2	1294.05(10)	.25	.11	.11	.15	0110(100%)	
11	F_2	1418.75(10)	.16	.08	.08	.22	0011(100%)	
14	F_2	$1804.5(1)^{e}$	1.05	.70	.72	21	0012(97%)	1010(2%)
14	F_2		1805.34	1805.49	1805.48		0012(96%)	1010(4%)
14	F_2		1807.20	1807.38	1807.37		0012(100%)	
14	F_2	1828.17(2)	56	65	65	57	1010(93%)	0012(7%)
16	F_2	2059.1(3)	04	17	18	1.09	0020(100%)	
20	F_2	$2602.55(10)^{f}$	14	71	67	-1.24	1012(70%)	2010(12%)
20	F_2		2604.97	2605.28	2605.09		1012(81%)	0014(9%)
20	F_2		2606.96	2607.20	2607.04		1012(86%)	0014(14%)
20	F_2	2623.8(1)	-1.30	-1.47	-1.43	.12	2010(83%)	1012(15%)
24	F_2	3068.5(1)	74	-1.71	-1.68	63	0030(100%)	
RMSD			0.59	0.74	0.73	0.79		

TABLE VIII. Comparison of experimental, CVPT^a and U(2) algebraic model band origins (cm^{-1}) for SiF₄. The *ab initio* force contants of Table III are used for the CVPT calculation.

 $\overline{a} N = 6v_1 + 2v_2 + 8v_3 + 3v_4$. This polyad number results from three independent resonances: $\omega_1 + \omega_2 \approx \omega_3$, $\omega_1 \approx 2\omega_4$, and $\omega_1 \approx 3\omega_2$.

 b The observations are from Ref. 35. Standard deviations given in parentheses.

 c The U(2) algebraic model calculations are from Ref. 36.

^d The largest two components in terms of $v_1v_2v_3v_4$ and their percentage (c_i^2) in the basis, based on the sixth order calculation. Only the components with the percentage larger than 1% are listed.

^e There are three (0012) F_2 states.

 f There are three (1012) F_{2} states.