Stopping of ion beam in a temperature anisotropic magnetized plasma

H. B. Nersisyan,[†] M. Walter, and G. Zwicknagel

Institut für Theoretische Physik II, Universität Erlangen, D - 91058 Erlangen, Germany

Abstract

Using the dielectric theory for a weakly coupled plasma we investigate the stopping power of the ion in a temperature anisotropic magnetized electron plasma. The analysis is based on the assumption that the energy variation of the ion is much less than its kinetic energy. The obtained general expression for stopping power is analyzed for weak and strong magnetic fields (i.e., for the electron cyclotron frequency less than and greater than the plasma frequency), and for low and high ion velocities. It is found that the friction coefficient contains, in addition to the usual velocity independent friction coefficient, an anomulous term which diverges logarithmically as the projectile velocity approaches zero. The physical origin of this anomulous term is the coupling between the cyclotron motion of the electrons and the long-wave length, low-frequency fluctuations produced by the projectile ion.

[†]Permanent address: Division of Theoretical Physics, Institute of Radiophysics and Electronics, 1 Alikhanian Brothers Str., Ashtarak -2, 378410, Armenia E-mail: hrachya@irphe.am

I INTRODUCTION

Energy loss of the ions in a plasma has been a topic of great interest due to its considerable importance for the study of basic interactions of the charged particles in real media. Recent applications are electron cooling of heavy ion beams [1, 2, 3] and energy transfer for inertial confinement fusion (ICF) (see [4] for an overview). Electron cooling is realized by mixing the ion beam periodically with a cold electron beam of the same average velocity. The interaction length is normally about a few meters and the electron beam is guided by a magnetic field parallel to its direction of motion. The cooling of the ion beam may then be viewed as an energy loss in the common rest frame of both beams. Similar questions arise in heavy-ion-induced ICF. There a frozen hydrogen pellet is heated and compressed by stopping of ion beams in the surrounding converter. In this case the electrons of the solid state converter are acting like a plasma and absorb the incoming energy.

In the electron cooling process the velocity distribution of the electron beam is highly anisotropic because of the acceleration from the cathode to the cooling section. It can be described by a Maxwell distribution with two different temperatures, a longitudinal T_{\parallel} and a transversal T_{\perp} [1, 2, 3]. Furthermore, an external, longitudinal magnetic field is needed to guide the electrons from the cathode to and through the electron cooler and to stabilize the anisotropic velocity distribution by suppressing the transverse-longitudinal relaxation.

In the present paper we are interested in the influences of the magnetic field and the temperature anisotropy on the ion beam stopping power.

Since the early 1960's several theoretical calculations of the stopping power in a magnetized plasma have been presented [5, 6, 7, 8, 9, 10, 11, 12, 13, 14]. Stopping of a fast test particle moving with velocity V much higher than the electron thermal velocity $v_{\rm th}$ was studied in Refs. [5, 6, 8]. Energy loss of a charged particle moving with arbitrary velocity was studied in Ref. [7]. The expression obtained there for the Coulomb logarithm, $\Lambda = \ln(\lambda_D/\rho_{\perp})$ (where λ_D is the Debye length and ρ_{\perp} is the impact parameter for scattering for an angle $\vartheta = \pi/2$), corresponds to the classical description of collisions. In the quantum-mechanical case, the Coulomb logarithm is $\Lambda = \ln(\lambda_D/\lambda_B)$, where λ_B is the de Broglie wavelength of plasma electrons [15].

In Ref. [10], the expressions were derived describing the stopping power of a charged particle in Maxwellian plasma placed in a classically strong (but not quantizing) magnetic field ($\lambda_B \ll a_c \ll \lambda_D$, where a_c is the electron Larmor radius), under the conditions when scattering must be described quantum mechanically. Calculations were carried out for slow test particles whose velocities satisfy the conditions $(m/m_i)^{1/3}v_{\rm th} < V \ll v_{\rm th}$, where m_i is the mass of the plasma ions and m is the electron mass.

In the recent paper [11] the stopping power in the magnetized plasma has been investigated for high-velocity light particles taking into account the Larmor rotation of a test projectile in a magnetic field. It has been shown that the stopping power can exhibit an oscillatory dependence on the magnetic field and that it is much greater than in the case without magnetic field.

More attention has been paid on the stopping power in a strongly magnetized plasma for ions which move along the magnetic field [11, 12, 13]. Both uncorrelated [11, 13] and correlated [12] situations have been discussed.

These investigations have concentrated on the stopping power in temperature isotropic plasma. Extensions to nonlinear effects of ion stopping and temperature anisotropy have been done recently by particle-in-cell (PIC) computer simulation [14], where the case $T_{\parallel} \ll$ T_{\perp} has been investigated which is interesting for electron cooling process. Here, in the framework of dielectric theory, we will focus on the stopping power at arbitrary temperature anisotropy T_{\perp}/T_{\parallel} .

The paper is organized as follows. We start in Sec. II, with solving the linearized Vlasov-Poisson equations by means of Fourier transformation. This provides the general form of the linearized potential generated in a temperature anisotropic magnetized Maxwellian plasma by a projectile ion from which the stopping power is deduced.

In the next Sec. III, is dedicated to apply our results to nonmagnetized plasma. Calculations are carried out for small projectile velocities at arbitrary temperature anisotropy and arbitrary direction of ion motion with respect to the anisotropy axis.

Then we turn to the effect of a weak magnetic field on the stopping power in Sec. IV, while we concentrate on the influence of a strong magnetic field in Sec. V. In contrast with the papers [11, 13] we consider an ion motion in arbitrary direction.

As the last issue we investigate in Sec. VI the stopping power for small projectile velocities at arbitrary magnetic field and temperature anisotropy. The friction coefficient there contains an anomalous term which increases logarithmically when the projectile velocity approaches to zero.

The achieved results are finally summarized and discussed in Sec. VII.

II DIELECTRIC THEORY

For the temperature anisotropic plasma with two different temperatures T_{\parallel} , T_{\perp} of the electrons we define an average temperature $\overline{T} = \frac{1}{3}T_{\parallel} + \frac{2}{3}T_{\perp}$. Within the dielectric theory the electron plasma is described as a continuous, polarizable fluid (medium), which is represented by the phase-space density of the electrons $f(\mathbf{r}, \mathbf{v}, t)$. Here, only a mean-field interaction between the electrons is considered and hard collisions are neglected and the evolution of the distribution function $f(\mathbf{r}, \mathbf{v}, t)$ is determined by the Vlasov-Poisson equation is valid for weakly coupled plasmas where the number of electrons in the Debye sphere $N_{\rm D} = 4\pi n_0 \overline{\lambda}_{\rm D}^3 \gg 1$ is very large. Here n_0 is the electron density, $\overline{\lambda}_{\rm D} = (k_{\rm B}\overline{T}/4\pi n_0 e^2)^{1/2}$ is an averaged Debye length.

In the following, we consider a nonrelativistic projectile ion with charge Ze and with a velocity \mathbf{V} that moves in a magnetized temperature anisotropic plasma at an angle ϑ with respect to the magnetic field \mathbf{B}_0 . The axis defined by \mathbf{B}_0 also coincides with the degree of freedom with temperature T_{\parallel} . We assume that the energy variation of the ion is much smaller than its kinetic energy. The strength of the coupling betweeen an ion moving with velocity V and the electron plasma is given by the coupling parameter

$$\mathcal{Z} = \frac{|Z|}{N_{\rm D} \left[1 + V^2 / \overline{v}_{\rm th}^2\right]^{3/2}}.$$
 (1)

Here $\overline{v}_{\rm th} = (k_{\rm B}\overline{T}/m)^{1/2}$ is the average thermal velocity of an electron. The derivation of Eq. (1) is discussed in detail in Ref. [16]. The parameter \mathcal{Z} characterizes the ion-target coupling, where $\mathcal{Z} \ll 1$ corresponds to weak, almost linear coupling and $\mathcal{Z} \gtrsim 1$ to strong, nonlinear coupling.

For a sufficiently small perturbation ($\mathcal{Z} \ll 1$) the linearized Vlasov equation of the plasma may be written as

$$\frac{\partial f_1}{\partial t} + \mathbf{v} \,\frac{\partial f_1}{\partial \mathbf{r}} - \omega_c \left[\mathbf{v} \times \mathbf{b}\right] \frac{\partial f_1}{\partial \mathbf{v}} = -\frac{e}{m} \,\frac{\partial \varphi}{\partial \mathbf{r}} \,\frac{\partial f_0}{\partial \mathbf{v}},\tag{2}$$

where $f = f_0 + f_1$ and the self-consistent electrostatic potential φ is determined by the Poisson equation

$$\nabla^2 \varphi = -4\pi Z e \delta(\mathbf{r} - \mathbf{V}t) + 4\pi e \int d\mathbf{v} f_1(\mathbf{r}, \mathbf{v}, t).$$
(3)

The **b** is the unit vector parallel to \mathbf{B}_0 , -e and $\omega_c = eB_0/mc$ are the charge and Larmor frequency of plasma electrons respectively, f_0 is the unperturbed distribution function of plasma electrons, which in the case of temperature anisotropic, homogeneous electron plasma is given by two Maxwellians for the longitudinal and transversal degrees of freedom

$$f_0(v_{\parallel}, v_{\perp}) = \frac{n_0}{(2\pi)^{3/2} v_{\rm th\perp}^2 v_{\rm th\parallel}} \exp\left(-\frac{v_{\perp}^2}{2v_{\rm th\perp}^2}\right) \exp\left(-\frac{v_{\parallel}^2}{2v_{\rm th\parallel}^2}\right) , \qquad (4)$$

where $\langle v_{\parallel}^2 \rangle = v_{\rm th\parallel}^2 = k_{\rm B} T_{\parallel}/m, \quad \langle v_{\perp}^2 \rangle = 2v_{\rm th\perp}^2 = 2k_{\rm B} T_{\perp}/m.$

By solving Eqs. (2) and (3) in space-time Fourier components, we obtain the electrostatic potential

$$\varphi(\mathbf{r},t) = \frac{Ze}{2\pi^2} \int d\mathbf{k} \, \frac{\exp\left[i\mathbf{k}(\mathbf{r}-\mathbf{V}t)\right]}{k^2 \varepsilon(\mathbf{k},\mathbf{kV})} \,, \tag{5}$$

which provides the dynamical response of the temperature anisotropic plasma to the motion of the projectile ion in the presence of the external magnetic field. Here $\epsilon(\mathbf{k}, \omega)$ is the dielectric function of a temperature anisotropic, magnetized plasma which is given by

$$\varepsilon(\mathbf{k},\omega) = 1 + \frac{1}{k^2 \lambda_{\mathrm{D}\parallel}^2} \left[G(s) + iF(s) \right]$$

$$= 1 + \frac{1}{k^2 \lambda_{\mathrm{D}\parallel}^2} \left\{ 1 + is\sqrt{2} \int_0^\infty dt \exp\left[ist\sqrt{2} - X(t)\right] \right.$$

$$+ \frac{k v_{\mathrm{th}\parallel} \sqrt{2}}{\omega_c} \sin^2 \alpha (1-\tau) \int_0^\infty dt \sin\left(\frac{\omega_c t \sqrt{2}}{k v_{\mathrm{th}\parallel}}\right) \exp\left[ist\sqrt{2} - X(t)\right] \right\}$$

$$(6)$$

with

$$X(t) = t^2 \cos^2 \alpha + k^2 a_{c\perp}^2 \sin^2 \alpha \left[1 - \cos\left(\frac{\omega_c t \sqrt{2}}{k v_{\text{th}\parallel}}\right) \right]$$
(7)

where $\lambda_{\mathrm{D}\parallel} = v_{\mathrm{th}\parallel}/\omega_p$, ω_p is the plasma frequency, $s = \omega/kv_{\mathrm{th}\parallel}$, $\tau = T_{\perp}/T_{\parallel}$, $a_c = v_{\mathrm{th}\perp}/\omega_c$ and α is the angle between the wave vector **k** and the magnetic field.

As shown in Appendix A, Eqs. (6) and (7) are identical with the Bessel function representation of $\varepsilon(\mathbf{k}, \omega)$ derived e.g. by Ichimaru [17]. Eqs. (6) and (7) are, however, more convenient when studying the weak and strong magnetic field limits in Secs. IV and V.

The stopping power S of an ion is defined as the energy loss of the ion in a unit length due to interactions with the plasma electrons. From Eq. (5) it is straightforward to calculate the electric field $\mathbf{E} = -\nabla \varphi$, and the stopping force acting on the ion. Then, the stopping power of the projectile ion becomes

$$S = -\frac{dE}{dl} = Ze \frac{\partial}{\partial \mathbf{r}} \varphi(\mathbf{r}, t) \Big|_{\mathbf{r} = \mathbf{V}t}$$

$$= \frac{2Z^2 e^2 \lambda_{\mathrm{D}\parallel}^2}{\pi^2} \int_0^{k_{\mathrm{max}}} k^3 dk \int_0^1 d\mu \int_0^{\pi} d\varphi \frac{\cos \Theta F(s)}{[k^2 \lambda_{\mathrm{D}\parallel}^2 + G(s)]^2 + F^2(s)} ,$$
(8)

where $\mu = \cos \alpha$ was the angle between **k** and **B**₀, Θ is the angle between **k** and **V**, $s = \mathbf{k} \cdot \mathbf{V} / k v_{\text{th}\parallel} = (V / v_{\text{th}\parallel}) \cos \Theta$, $\cos \Theta = \mu \cos \vartheta - \sqrt{1 - \mu^2} \sin \vartheta \cos \varphi$, and ϑ is the angle between

V and \mathbf{B}_0 . In Eq. (8) we introduced a cutoff parameter $k_{\text{max}} = 1/r_{\text{min}}$ (where r_{min} is the effective minimum impact parameter) in order to avoid the logarithmic divergence at large k. This divergence corresponds to the incapability of the linearized Vlasov theory to treat close encounters between the projectile ion and the plasma electrons properly. For r_{min} we thus use the effective minimum impact parameter of classical binary Coulomb collisions $r_{\text{min}} = Ze^2/mv_r^2$ for relative velocities $v_r \simeq (V^2 + \overline{v}_{\text{th}}^2)^{1/2}$, which is often called the "distance of closest approach." Hence

$$k_{\max} = \frac{1}{r_{\min}} = \frac{m(V^2 + \overline{v}_{\rm th}^2)}{Ze^2}.$$
(9)

A two temperature description of an electron plasma is valid only when the ion beamplasma interaction time is less than the relaxation time between the two temperatures, T_{\parallel} and T_{\perp} . For an estimate we will briefly consider the field-free case, because the external magnetic field suppresses the relaxation between the transversal and longitudinal temperatures during the time of flight of the ion beam through plasma.

The problem of a temperature relaxation in a temperature anisotropic plasma with and without of an external magnetic field was considered by Ichimaru [17]. Within the dominantterm approximation the relaxation time $\Delta \tau_{\rm rel}$ for the plasma without magnetic field is given by

$$\frac{1}{\Delta\tau_{\rm rel}} = \frac{8}{15} \sqrt{\frac{\pi}{m}} \frac{n_0 e^4}{(k_{\rm B} T_{\rm eff})^{3/2}} \ln \Lambda_c , \qquad (10)$$

where $\ln \Lambda_c = \ln(N_D)$ is the Coulomb logarithm and the effective electron temperature T_{eff} is defined through

$$\frac{1}{T_{\text{eff}}^{3/2}} = \frac{15}{2} \int_{0}^{1} \frac{\mu^2 (1 - \mu^2) d\mu}{[\mu^2 T_{\parallel} + (1 - \mu^2) T_{\perp}]^{3/2}}$$

$$= \frac{5\sqrt{3}}{12\overline{T}^{3/2}} \frac{(1 + 2\tau)^{3/2}}{(\tau - 1)^2} \left[\frac{\tau + 2}{\sqrt{|\tau - 1|}} p_0(\tau) - 3 \right] ,$$
(11)

$$p_0(\tau) = \begin{cases} \ln \frac{1+\sqrt{1-\tau}}{\sqrt{\tau}}, & \tau < 1\\ \arctan \sqrt{\tau-1}, & \tau > 1 \end{cases}$$
(12)

The relaxation time calculated from Eq. (11) are of the order of 10^{-6} s, 0.5×10^{-5} s and 10^{-3} s for averaged temperatures $\overline{T} = 10^{-2}$ eV, $\overline{T} = 0.1$ eV and $\overline{T} = 1$ eV, respectively, for anisotropies $\tau \simeq 0.01 - 100$. The interaction time (for instance, for ICF or for electron cooling) is about $10^{-7} - 10^{-8}$ s. Therefore, ion beam-plasma interaction time can be very small compared to the plasma relaxation time.

III STOPPING POWER IN PLASMA WITHOUT MAGNETIC FIELD

Let us analyse expression (8) in the case when a projectile ion moves in a temperature anisotropic plasma without magnetic field. The plasma dielectric function from Eqs. (6) and (7) now takes the form

$$\varepsilon(\mathbf{k},\omega) = 1 + \frac{1}{k^2 \lambda_{\text{D}\parallel}^2} \frac{1}{A^2} W\left(\frac{s}{A}\right) . \tag{13}$$

Here $A = (\mu^2 + \tau(1 - \mu^2))^{1/2}$ and $W(s) = g_0(s) + if_0(s)$ is the plasma dispersion function [18],

$$g_0(s) = 1 - s\sqrt{2} Di\left(\frac{s}{\sqrt{2}}\right); \quad f_0(s) = \sqrt{\frac{\pi}{2}}s \exp\left(-\frac{s^2}{2}\right), \tag{14}$$

where

$$Di(s) = \exp(-s^2) \int_{0}^{s} dt \exp(t^2)$$
(15)

is the Dawson integral [18] which has for large arguments s the asymptotic $Di(s) \simeq 1/2s + 1/4s^3$.

Substituting Eq. (13) into Eq. (8) and performing the k-integration we obtain

$$S_0 = \frac{Z^2 e^2}{2\pi^2 \lambda_{\text{D}\parallel}^2} \int_0^1 d\mu \int_0^\pi d\varphi \frac{\cos\Theta}{A^2} Q_0 \left(\frac{v}{v_{\text{th}\parallel}} \frac{\cos\Theta}{A} , \xi_\parallel A \right) , \qquad (16)$$

where $\xi_{\parallel} = k_{\max} \lambda_{D\parallel}$ and

$$Q_0(x,\xi) = f_0(x) \ln \frac{f_0^2(x) + [\xi^2 + g_0(x)]^2}{f_0^2(x) + g_0^2(x)}$$

$$+2g_0(x) \left[\arctan \frac{g_0(x)}{f_0(x)} - \arctan \frac{\xi^2 + g_0(x)}{f_0(x)} \right] .$$
(17)

In the case of temperature isotropic plasma $(T_{\perp} = T_{\parallel} \equiv T, \text{ and } \tau = 1) A = 1$ and Eq. (16) coincides with the result of e.g. Ref. [19]

$$S_0 = \frac{Z^2 e^2}{2\pi \lambda_D^2} \frac{v_{\rm th}^2}{V^2} \int_0^{V/v_{\rm th}} d\mu \mu Q_0(\mu, \xi) , \qquad (18)$$

where $v_{\rm th} = v_{\rm th\parallel} = v_{\rm th\perp}$, $\lambda_D = v_{\rm th}/\omega_p$, and $\xi = k_{\rm max}\lambda_D$.

When a projectile ion moves slowly through a plasma, the electrons have much time to experience the ion attractive potential. They are accelerated towards the ion, but when they reach its trajectory the ion has already moved forward a little bit. Hence, we expect an increased density of electrons at some place in the trail of the ion. This negative charge density pulls back the positive ion and gives rise to the stopping. This drag force is of particular interest for the electron cooling process. In the limit of small velocities $S \simeq R \cdot V$. This looks like the friction law of a viscous fluid, and accordingly R is called the friction coefficient. However, in the case of an ideal plasma it should be noted that this law does not depend on the plasma viscosity and is not a consequence of electron-electron collisions which are neglected in the Vlasov equation.

The Taylor expansion of Eq. (16) for small $V(V \ll \overline{v}_{\rm th})$ yields the friction law

$$S_0 = \frac{Z^2 \left(e^2 / \overline{\lambda}_D^2\right)}{3\sqrt{2\pi}} \frac{V}{\overline{v}_{\rm th}} \psi(\overline{\xi}) \left[I_1(\tau) + I_2(\tau) \sin^2 \vartheta\right],\tag{19}$$

where $\overline{\xi} = k_{\max} \overline{\lambda}_D = (1 + V^2 / \overline{v}_{\text{th}}^2) / \mathcal{Z} \simeq 1 / \mathcal{Z},$

$$I_1(\tau) = \frac{3}{\psi(\overline{\xi})} \left(\frac{2\tau+1}{3}\right)^{3/2} \int_0^1 d\mu \; \frac{\mu^2 \psi(\xi_{\parallel} A(\mu))}{A^3(\mu)},\tag{20}$$

$$I_2(\tau) = \frac{3}{2\psi(\overline{\xi})} \left(\frac{2\tau+1}{3}\right)^{3/2} \int_0^1 d\mu \; \frac{(1-3\mu^2)\psi(\xi_{\parallel}A(\mu))}{A^3(\mu)},\tag{21}$$

and the function ψ is

$$\psi(\xi) = \ln(1+\xi^2) - \frac{\xi^2}{1+\xi^2}.$$
(22)

In the case of temperature isotropic plasma ($\tau = 1$) we have $I_1 = 1$ and $I_2 = 0$. Then the Eq. (19) becomes the usual friction law in an isotropic plasma [19]. For the strongly temperature anisotropic case, when $\tau \ll 1$ ($T_{\perp} \ll T_{\parallel}$) we have $\xi_{\parallel} \simeq \sqrt{3}/\mathcal{Z}$ and

$$I_1 \simeq -\frac{\sqrt{3}}{6\psi(\overline{\xi})} \left[Li_2(1+\xi_{\parallel}^2) + \ln(1+\xi_{\parallel}^2) \right] , \qquad (23)$$

$$I_2 \simeq \frac{\sqrt{3}}{12\psi(\overline{\xi})} \left[\xi_{\parallel}^2 + 2\ln(1+\xi_{\parallel}^2) + 3Li_2(1+\xi_{\parallel}^2) \right].$$
(24)

Here the functions I_1 and I_2 do not depend on τ , and $Li_2(x)$ is the dilogarithm function [20]. Note that $\mathcal{Z} \ll 1$ and therefore $\overline{\xi} \gg 1$, $\xi_{\parallel} \gg 1$ in Eqs. (23) and (24). The Coulomb logarithms in Eqs. (23) and (24) are then the leading terms and

$$I_1 \simeq \frac{\sqrt{3}}{6} \ln \frac{1}{\mathcal{Z}} \ll I_2 \simeq \frac{\sqrt{3}}{8\mathcal{Z}^2} \frac{1}{\ln(1/\mathcal{Z})} .$$

$$(25)$$

The normalized friction coefficient (Eq. (19)) is thus dominated by the second term and increases with increasing ϑ .

In the opposite case, $\tau \gg 1$ $(T_{\perp} \gg T_{\parallel})$, the evoluation of Eqs. (20) and (21) yields

$$I_1 \simeq \frac{\pi\sqrt{6}}{3\psi(\overline{\xi})} \left(\sqrt{1 + \frac{3}{2}\overline{\xi}^2} - 1 - 2\ln\frac{1 + \sqrt{1 + \frac{3}{2}\overline{\xi}^2}}{2} \right) , \qquad (26)$$

$$I_2 \simeq \frac{\pi\sqrt{6}}{6\psi(\overline{\xi})} \left(1 + \frac{1}{\sqrt{1 + \frac{3}{2}\overline{\xi}^2}} - 2\sqrt{1 + \frac{3}{2}\overline{\xi}^2} + 6\ln\frac{1 + \sqrt{1 + \frac{3}{2}\overline{\xi}^2}}{2} \right) , \qquad (27)$$

and

$$I_1 \simeq -I_2 \simeq \frac{\pi}{2\mathcal{Z}\ln(1/\mathcal{Z})}.$$
(28)

Then $I_1 + I_2 \sin^2 \vartheta \simeq I_1 \cos^2 \vartheta$ and the normalized friction coefficient decreases with increasing of ϑ in this case.

In Fig. 1 the normalized friction coefficient $I_1 + I_2 \sin^2 \vartheta$ is plotted as a function of temperature anisotropy τ for $\vartheta = 0$ (solid line), $\vartheta = \pi/6$ (dotted line), $\vartheta = \pi/3$ (dashed line), $\vartheta = \pi/2$ (dot-dashed line) and for fixed plasma density and average temperature $(\mathcal{Z} = 0.2)$. Fig. 1 shows an enhancement of the friction coefficient when the ion moves along the direction with low temperature. This effect can be easily explained in a binary collision picture. Let us consider the particular case of strongly anisotropic plasma $T_{\perp} \gg T_{\parallel}$. In this case the plasma electrons move mostly in the direction across to the anisotropy axis. For $\vartheta \simeq \pi/2$ the projectile ion moves along the plasma electrons thermal fluctuation direction and effective impact parameter for electron-ion collision is very small. Then the friction coefficient decreases. For $\vartheta \simeq 0$ the projectile ion moves across to the direction of plasma electrons thermal fluctuation. Therefore, the impact parameter for electron-ion collisions increases which rises the friction coefficient.

For arbitrary projectile velocities we evaluated Eq. (16) numerically. In Figs. 2 and 3 the stopping power is plotted for strongly temperature anisotropic plasmas ($\tau = 10^{-2}$ and $\tau = 10^2$ in Figs. (2) and (3) respectively) with $n_0 = 10^8 \text{cm}^{-3}$, $\overline{T} = 0.1 \text{eV}$ and for four values of ϑ ; $\vartheta = 0$ (dotted line), $\vartheta = \pi/6$ (dashed line), $\vartheta = \pi/3$ (long-dashed line), $\vartheta = \pi/2$ (dot-dashed line). The solid lines are plotted for temperature isotropic plasma with $T = \overline{T} = 0.1 \text{eV}$. The general behaviour of the stopping power for two anisotropy parameters τ is characterized by an increase by comparision with the isotropic case. At $\vartheta \simeq \pi/2$ and $\tau = 10^{-2}$ (Fig. (2)) the ion moves in direction accross to the longitudinal electron motion with the lower temperature T_{\perp} and the maximum of the stopping power is around $V \simeq v_{\text{th}\perp}$, whereas the maximum for an ion motion in longitudinal direction is at $V \simeq v_{\text{th}\parallel} \gg v_{\text{th}\perp}$.

IV STOPPING IN PLASMAS WITH WEAK MAGNETIC FIELD

For the case when the magnetic field is weak, in the sense that the dimensionless parameter $\eta = \omega_c/\omega_p$ is much less than unity, the functions G and F, Eqs. (6) and (7), which define the dielectric function, can be expanded about its field free values $g_0(s/A)/A^2$, $f_0(s/A)/A^2$ Eqs. (14) and (15)

$$G(s) + iF(s) = \frac{1}{A^2} \left[g_0\left(\frac{s}{A}\right) + if_0\left(\frac{s}{A}\right) \right] + \eta^2 \frac{\sin^2 \alpha}{(k\lambda_{D\parallel})^2} \left[g_1(s) + if_1(s) \right],$$
(29)

where

$$g_{1}(s) + if_{1}(s) = \frac{2}{3}(1-\tau) \int_{0}^{\infty} t^{3} dt \left(\frac{t^{2}}{2}\tau \sin^{2}\alpha - 1\right) \exp(ist\sqrt{2} - A^{2}t^{2})$$

$$+ \frac{is\sqrt{2}}{6}\tau \int_{0}^{\infty} t^{4} dt \exp(ist\sqrt{2} - A^{2}t^{2}),$$
(30)

 $s = \omega / k v_{\text{th}\parallel}$. Substituting this expression (29) into Eq. (8) leads to

$$S = S_0 + \eta^2 S_1, (31)$$

where S_0 is the stopping power in plasma without magnetic field Eq. (16) and $\eta^2 S_1$ represents the change due to a weak magnetic field. After some simplifications it becomes

$$S_{1} = \sqrt{\frac{\pi}{2}} \frac{Z^{2} e^{2}}{24\pi^{2} \lambda_{D\parallel}^{2}} \frac{V}{v_{\text{th}\parallel}} \int_{0}^{1} d\mu \int_{0}^{\pi} d\varphi \frac{(1-\mu^{2})\cos^{2}\Theta}{A^{5}}$$

$$\times \exp\left(-\frac{V^{2}}{v_{\text{th}\parallel}^{2}} \frac{\cos^{2}\Theta}{2A^{2}}\right) \frac{\tau\left(7 - \frac{V^{2}}{v_{\text{th}\parallel}^{2}} \frac{\cos^{2}\Theta}{A^{2}}\right) - 4A^{2}}{f_{0}^{2}\left(\frac{V}{v_{\text{th}\parallel}} \frac{\cos\Theta}{A}\right) + g_{0}^{2}\left(\frac{V}{v_{\text{th}\parallel}} \frac{\cos\Theta}{A}\right)}.$$

$$(32)$$

In the temperature isotropic plasma ($\tau = 1$) Eq. (32) coincides with the results by May and Cramer [7] after integration over φ . Note that the additional term S_1 does not depend on the cutoff parameter k_{max} .

In the next subsections we evaluate Eq. (32) for small and large projectile velocities.

A Small projectile velocities

When the projectile ion moves slowly $(V < \overline{v}_{th})$ in plasma Eq. (32) leads to the simplified expression

$$S_1 = \frac{Z^2 e^2}{60\pi \overline{\lambda}_D^2} \sqrt{\frac{\pi}{2}} \frac{V}{\overline{v}_{\rm th}} P(\vartheta, \tau), \qquad (33)$$

with

$$P(\vartheta,\tau) = \left(\frac{1+2\tau}{3}\right)^{3/2} \left[P_1(\tau) + P_2(\tau)\sin^2\vartheta\right],\tag{34}$$

$$P_1(\tau) = \frac{5}{6(1-\tau)^2} \left\{ 14\tau + 25 - \frac{3(9\tau+4)}{\sqrt{|1-\tau|}} p_0(\tau) \right\},\tag{35}$$

$$P_2(\tau) = \frac{5}{12\tau(1-\tau)^2} \left\{ \frac{3\tau(23\tau+16)}{\sqrt{|1-\tau|}} p_0(\tau) - 28\tau^2 - 91\tau + 2 \right\} .$$
(36)

Here, the function $p_0(\tau)$ is given by Eq. (12). In temperature isotropic plasma with $\tau = 1$ we have $P_1(1) = P_2(1) = 1$.

In Fig. 4 the normalized friction coefficient $P(\vartheta, \tau)$ for the additional stopping power S_1 is plotted as a function of τ for $\vartheta = 0$ (solid line), $\vartheta = \pi/6$ (dotted line), $\vartheta = \pi/3$ (dashed line), $\vartheta = \pi/2$ (dot-dashed line). The general behaviour of $P(\vartheta, \tau)$ is similar to the friction coefficient of the plasma without magnetic field (see Fig. 1). Here, the correction $P(\vartheta, \tau)$ can be also negative at small τ and ϑ , which then corresponds to a slight decrease of the stopping power, Eq. (31).

B High projectile velocities

When the projectile ion moves with large velocity $(V \gg \overline{v}_{\rm th})$, Eq. (32), yields

$$S_1 = -\frac{Z^2 e^2 \omega_p^2}{8V^2} \left\{ 2C_1 (1 + \cos^2 \vartheta) - C_2 B(\vartheta, \tau) \left[\cos^2 \vartheta + \frac{\sin^2 \vartheta}{B(\vartheta, \tau) + 1} \right] \right\},\tag{37}$$

where

$$B(\vartheta,\tau) = \left(\frac{\tau}{\tau\cos^2\vartheta + \sin^2\vartheta}\right)^{1/2},\tag{38}$$

$$C_1 = \frac{1}{3\sqrt{2\pi}} \int_0^\infty \frac{x^2 \exp(-x^2/2) dx}{f_0^2(x) + g_0^2(x)} , \ C_2 = \frac{1}{3\sqrt{2\pi}} \int_0^\infty \frac{x^2(7-x^2) \exp(-x^2/2) dx}{f_0^2(x) + g_0^2(x)}.$$
 (39)

For numbers C_1 and C_2 we get the accurate values $C_1 = 1$ and $C_2 = 0$ ($C_2 \simeq 10^{-12}$) respectively. Therefore from Eq. (37) we have finally

$$S_1 = -\frac{Z^2 e^2 \omega_p^2}{4V^2} (1 + \cos^2 \vartheta).$$
(40)

This result is in accord with the results of Honda et al. [6] and May and Cramer [7], who, however, kept the terms $O(V^{-4})$ in their work as well. Although the function S_1 in Eq. (40) is proportional to the plasma density, the full correction term $\eta^2 S_1$ does not depend on the plasma density.

In Figs. (5) and (6) we show the velocity dependence of the function S_1 for $\tau = 10^{-2}$ and $\tau = 10^2$ respectively. The different curves are $\vartheta = 0$ (solid line), $\vartheta = \pi/6$ (dotted line), $\vartheta = \pi/3$ (dashed line), $\vartheta = \pi/2$ (dot-dashed line). For small and medium projectile velocities the weak magnetic field decreases the total stopping power for small τ and increases it in the high τ limit. For high projectile velocities the magnetic field always reduces the stopping power independent of the temperature anisotropy, see Eq. (40).

V STOPPING IN PLASMAS WITH STRONG MAGNETIC FIELD

We now turn to the case when a projectile ion moves in a temperature anisotropic plasma with a strong magnetic field, which is on one hand, sufficiently weak to allow a classical description ($\hbar\omega_c < k_{\rm B}T_{\perp}$ or $\hbar/mv_{\rm th\perp} < a_c$, and, on the other hand, comparatively strong so that the cyclotron frequency of the plasma electrons exceeds the plasma frequency $\omega_c \gg \omega_p$. This limits the values of the magnetic field itself and values of perpendicular temperature and plasma density. From these conditions we can obtain

$$3 \times 10^{-6} n_0^{1/2} < B_0 < 10^5 T_\perp, \tag{41}$$

where n_0 is measured in cm⁻³, T_{\perp} is measured in eV, and B_0 in kG. Conditions (41) are always true in the range of parameters $n_0 < 10^{15}$ cm⁻³, $B_0 < 100$ kG, $T_{\perp} > 10^{-3}$ eV. Then the perpendicular motion of the electrons is completely quenched and the stopping power depends only on the longitudinal electron temperature T_{\parallel} . The dependence on the transversal temperature will be only introduced by the cutoff parameter Eq. (9).

In the limit of sufficiently strong magnetic field, Eq. (8) becomes

$$S_{\rm inf} = \frac{2Z^2 e^2}{\pi^2 \lambda_{D\parallel}^2} \int_0^{\xi_{\parallel}} k^3 dk \int_0^1 d\mu \int_0^{\pi} d\varphi \frac{\cos \Theta f_0(s)}{\left[k^2 + g_0(s)\right]^2 + f_0^2(s)},\tag{42}$$

with $s = (V/v_{\text{th}\parallel})(\cos \Theta/\mu)$ and g_0 , f_0 from Eqs. (14), which gives after integration over k

$$S_{\rm inf} = \frac{Z^2 e^2}{2\pi^2 \lambda_{D\parallel}^2} \int_0^1 d\mu \int_0^{\pi} d\varphi \cos \Theta Q_0 \left(\frac{V}{v_{\rm th\parallel}} \frac{\cos \Theta}{\mu}, \xi_{\parallel} \right).$$
(43)

Here the function Q_0 is given by Eq. (17). For further simplification of Eq. (43) we introduce the new variable of integration $x = \cos \Theta/\mu$. After φ integration in Eq. (43) we finally find the stopping power in the presence of a strong magnetic field as

$$S_{\rm inf}(V,\vartheta) = \frac{Z^2 e^2}{8\pi \lambda_{D\parallel}^2} Q\left(\frac{V}{v_{\rm th\parallel}},\vartheta\right),\tag{44}$$

where

$$Q\left(\frac{V}{v_{\rm th\parallel}},\vartheta\right) = \sin^2\vartheta \int_{-\infty}^{\infty} \frac{Q_0\left(\frac{V}{v_{\rm th\parallel}}x,\xi_{\parallel}\right)xdx}{(x^2+1-2x\cos\vartheta)^{3/2}}.$$
(45)

In the previous works [11, 12, 13] only the case of $\vartheta = 0$ the motion of the projectile along the magnetic field direction has been investigated. In this case the integral in Eq. (45) diverges, while prefactor $\sin^2 \vartheta$ tends to zero. Introducing the new variable of the integration in Eq. (45) $y = (x - \cos \vartheta) / \sin \vartheta$ we obtain for vanishing angle ϑ

$$Q\left(\frac{V}{v_{\rm th\parallel}}, \vartheta \to 0\right) = 2Q_0\left(\frac{V}{v_{\rm th\parallel}}, \xi_{\parallel}\right).$$
(46)

Thus expression (44) reproduces the known results for the stopping power on an ion which moves along the direction of the magnetic field [11, 12, 13].

In the following paragraphs we will discuss its low and high velocity limits.

A Small projectile velocities

In the low velocity limit $(V \ll v_{\rm th})$ Eq. (45) becomes

$$Q\left(\frac{V}{v_{\rm th\parallel}},\vartheta\right) \simeq \frac{2V}{v_{\rm th\parallel}} \left\{\sqrt{2\pi}\psi(\xi_{\parallel}) \left[\sin^2\vartheta\ln\left(\frac{2v_{\rm th\parallel}}{V\sin\vartheta}\right) + 1 - 2\sin^2\vartheta\right] + C_1(\xi_{\parallel})\sin^2\vartheta\right\}, \quad (47)$$

where

$$C_{1}(\xi_{\parallel}) = \int_{0}^{1} \frac{dx}{x^{2}} \left[Q_{0}(x,\xi_{\parallel}) - \sqrt{2\pi}\psi(\xi_{\parallel})x \right] + \int_{1}^{\infty} \frac{dx}{x^{2}} Q_{0}(x,\xi_{\parallel}).$$
(48)

Here, the function ψ is defined by Eq. (22). Since we deal with small ion beam-plasma coupling $\mathcal{Z} \ll 1$ we have, $\xi_{\parallel} \gg 1$ in Eqs. (47) and (48) and the function $C_1(\xi)$ simplifies

$$C_1(\xi_{\parallel}) \simeq \sqrt{2\pi} \ln \frac{2}{\gamma} \ln \xi_{\parallel} + 0.6, \qquad (49)$$

where $\gamma = 0.5772$ is Euler's constant.

We note that the friction coefficient S_{inf}/V from Eqs. (44) and (47) contains a logarithmically large term which vanishes for $\vartheta \to 0$. It will be shown in the next section that this behaviour is a characteristic feature of the stopping power at low velocities and the friction coefficient for arbitrary strength of the magnetic field.

B High projectile velocities

In the case of high projectile velocities $(V \gg v_{\rm th\parallel})$ the general expression (45) becomes

$$Q\left(\frac{V}{v_{\rm th\parallel}},\vartheta\right) \simeq \frac{4\pi v_{\rm th\parallel}^2}{V^2} \left\{\sin^2\vartheta \left[\ln\left(\frac{2V}{v_{\rm th\parallel}\sin\vartheta}\right) + C_2(\xi_{\parallel}) - 2\right] + 1\right\},\tag{50}$$

where

$$C_{2}(\xi_{\parallel}) = \frac{1}{2\pi} \int_{0}^{1} Q_{0}(x,\xi_{\parallel}) x dx + \int_{1}^{\infty} \frac{dx}{x} \left[\frac{x^{2}}{2\pi} Q_{0}(x,\xi_{\parallel}) - 1 \right]$$
(51)

which gives for $\xi_{\parallel} \gg 1 \ C_2(\xi_{\parallel}) \simeq \ln \xi_{\parallel}$. The stopping power for strong magnetic fields shows in the low and high velocity limits (Eqs. (47) and (50)) an enhancement for ions moving transversal to the magnetic field compared to the case of the longitudinal motion ($\vartheta = 0$). This effect is in agreement with PIC simulation results [14]. In contrast to the field-free case, at strong magnetic field and for $\vartheta = 0$, $V \gg v_{\text{th}\parallel}$ (Eqs. (44) and (50)) we have $S_{\text{inf}} \simeq Z^2 e^2 \omega_p^2 / 2V^2$ independent of k_{max} . The cutoff k_{max} necessary at low ion velocities is, however, less well defined here than for the field-free case, where the cutoff (9) was deduced from the binary collision picture. Now, the electrons are forced to move parallel to \mathbf{B}_0 . Since we assumed the motion of the ion in this direction as well the ion and an electron just pass each other along a straight line. For symmetry reasons the total momentum transfer and the stopping power is zero. Purely binary interactions contribute nothing and the stopping of the ion is only due to the collective response of the plasma, that is, due to modes with long wavelengths $k < 1/\lambda_{D\parallel}$. This suggests taking k_{max} of the order of $1/\lambda_{D\parallel}$, but further investigations are clearly needed here for a more precise description in this particular case.

In Figs. (7) and (8), the stopping power S_{inf} is plotted as a function of projectile velocity (in units of $v_{\text{th}\parallel}$) for $n_0 = 10^6 \text{cm}^{-3}$, $T_{\parallel} = 10^{-4} \text{eV}$, $T_{\perp} = 10^{-5} \text{eV}$ (Fig. (7)), $T_{\perp} = 0.1 \text{eV}$ (Fig. (8)), and for four different values of angle ϑ : $\vartheta = 0$ (solid line), $\vartheta = \pi/6$ (dotted line), $\vartheta = \pi/3$ (dashed line) and $\vartheta = \pi/2$ (dash-dotted line). The enhancement of $S_{\text{inf}}(V, \vartheta)$ with respect to $S_{\text{inf}}(V, 0)$ in the low and in high velocity limit by increasing of the angle ϑ is documented in Fig. (9), for $T_{\parallel} = 10^{-4} \text{eV}$, $T_{\perp} = 0.1 \text{eV}$, $n_0 = 10^6 \text{cm}^{-3}$, $\vartheta = \pi/6$ (solid line), $\vartheta = \pi/4$ (dotted line), $\vartheta = \pi/3$ (dashed line) and $\vartheta = \pi/2$ (dash-dotted line). The physical origin of this angular behaviour in the low and high velocity limits is the enhancement of the effective impact parameter for an individual electron-ion collision with increasing ϑ . For medium projectile velocities $V \simeq v_{\text{th}\parallel}$ the collective excitations in plasma become important and then stopping power is higher for small ϑ .

VI STOPPING AT ARBITRARY MAGNETIC FIELD AND IN LOW-VELOCITY LIMIT. ANOMALOUS FRICTION COEFFICIENT

We now proceed with a projectile ion at low velocities and at arbitrary magnetic field. This regime is of particular importance for the electron cooling process [1, 2, 3]. In the presence of a magnetic field the friction coefficient here contains a term which diverges like $\ln(v_{\text{th}\parallel}/V)$ in addition to the usual (see e.g. Sec. III) constant one.

For this consideration it is convenient to use the Bessel function representation of the dielectric function which has been given e.g. by Ichimaru [17], see Appendix A Eq. (A7), and to write the real and imaginary parts of Eq. (A7) separately

$$G = 1 - \frac{\sqrt{2\omega}}{|k_{\parallel}|v_{\rm th\parallel}} \Lambda_0(z) Di \left(\frac{\omega}{|k_{\parallel}|v_{\rm th\parallel}\sqrt{2}}\right)$$

$$- \frac{\sqrt{2}}{|k_{\parallel}|v_{\rm th\parallel}} \sum_{n=1}^{\infty} \Lambda_n(z) \left\{ \omega \left[Di \left(\frac{\omega + n\omega_c}{|k_{\parallel}|v_{\rm th\parallel}\sqrt{2}}\right) + Di \left(\frac{\omega - n\omega_c}{|k_{\parallel}|v_{\rm th\parallel}\sqrt{2}}\right) \right]$$

$$+ n\omega_c \left(\frac{1}{\tau} - 1\right) \left[Di \left(\frac{\omega - n\omega_c}{|k_{\parallel}|v_{\rm th\parallel}\sqrt{2}}\right) - Di \left(\frac{\omega + n\omega_c}{|k_{\parallel}|v_{\rm th\parallel}\sqrt{2}}\right) \right] \right\},$$
(52)

$$F = \sqrt{\frac{\pi}{2}} \left\{ \frac{\omega}{|k_{\parallel}|v_{\text{th}\parallel}} \Lambda_0(z) \exp\left(-\frac{\omega^2}{2k_{\parallel}^2 v_{\text{th}\parallel}^2}\right) + \frac{2}{|k_{\parallel}|v_{\text{th}\parallel}} \sum_{n=1}^{\infty} \Lambda_n(z) \exp\left(-\frac{\omega^2 + n^2 \omega_c^2}{2k_{\parallel}^2 v_{\text{th}\parallel}^2}\right) \right\}$$
(53)

$$\times \left[\omega \operatorname{ch} \left(\frac{n \omega_c \omega}{k_{\parallel}^2 v_{\operatorname{th} \parallel}^2} \right) + n \omega_c \left(\frac{1}{\tau} - 1 \right) \operatorname{sh} \left(\frac{n \omega_c \omega}{k_{\parallel}^2 v_{\operatorname{th} \parallel}^2} \right) \right] \right\} \; .$$

The notations in Eqs. (52) and (53) are explained in Appendix A.

For the friction coefficient we have to consider S, given by Eq. (8) in the low-velocity limit and thus the functions G and F given by Eqs. (52) and (53), when $\omega = \mathbf{kV}$. Now we have to write the Taylor expansion of Eqs. (52) and (53) for small $\omega = \mathbf{kV}$. However, the first term of Eq. (53) exhibits a singular behaviour in the limit of $\omega = \mathbf{kV} \rightarrow 0$ where the k_{\parallel} integration diverges logarithmically for small k_{\parallel} . We must therefore keep $\omega = \mathbf{kV}$ finite in that integration to avoid such a divergence. This anomalous contribution which arises from the first term of Eq. (53) in low-velocity limit is

$$S_{\rm an} \simeq \left(\frac{2}{\pi^3}\right)^{1/2} \frac{Z^2 e^2}{\lambda_{D\parallel}^2} \frac{V}{v_{\rm th\parallel}} \int_0^{\xi_{\parallel}} k^3 dk \int_0^1 \frac{d\mu}{\mu} \int_0^{\pi} d\varphi \cos^2 \Theta \frac{\Lambda_0(z) \exp\left(-\frac{V^2}{2v_{\rm th\parallel}^2} \frac{\cos^2 \Theta}{\mu^2}\right)}{[k^2 + E_2(k,\mu)]^2} , \qquad (54)$$

where $\Lambda_0(z) = \exp(-z)I_0(z)$ and $E_2(k,\mu) = G(\omega = 0)$ is

$$E_2(k,\mu) = 1 + \frac{2\sqrt{2\eta}}{k\mu} \left(\frac{1}{\tau} - 1\right) \sum_{n=1}^{\infty} n\Lambda_n(z) Di\left(\frac{n\eta}{k\mu\sqrt{2}}\right) .$$
(55)

Here $z = (k^2 \tau / \eta^2)(1 - \mu^2)$, $\mu = \cos \alpha = k_{\parallel}/k$, and Θ is the angle between **k** and **V**. After μ and φ integration, see Appendix B, Eq. (54) reads

$$S_{\rm an} \simeq \left(\frac{2}{\pi}\right)^{1/2} \frac{Z^2 e^2}{4\lambda_{D\parallel}^2} \frac{V}{v_{\rm th\parallel}} \sin^2 \vartheta \ln \left(\frac{v_{\rm th\parallel}}{V} \frac{2.26}{\sin \vartheta}\right) \mathcal{F}(\tau, \eta), \tag{56}$$

with

$$\mathcal{F}(\tau,\eta) = \int_{0}^{\tau\xi_{\parallel}^{2}} \frac{\Lambda_{0}(x/\eta^{2})xdx}{[x+1+(\tau-1)\Lambda_{0}(x/\eta^{2})]^{2}}.$$
(57)

The function \mathcal{F} and thus S_{an} (56) vanishes in the limit $B_0 \to 0$ (or $\eta \to 0$) like

$$\mathcal{F}(\tau,\eta) \simeq \frac{\eta}{(2\pi)^{1/2}} \left[\arctan(k_{\max}\lambda_{D\perp}) - \frac{k_{\max}\lambda_{D\perp}}{1 + (k_{\max}\lambda_{D\perp})^2} \right] \,. \tag{58}$$

The anomalous term Eqs. (56) and (57) therefore represents a new effect arising from the presence of the magnetic field, which is not restricted to anisotropic plasmas.

For temperature isotropic plasma ($\tau = 1$) and for a sufficiently weak magnetic field $\eta < \xi_{\parallel}$ (or $\omega_c < k_{\max} v_{\text{th}\parallel}$), Eq. (57) takes the form

$$\mathcal{F}(\tau,\eta) \simeq \exp\left(\frac{1}{\eta^2}\right) \left[\left(1 + \frac{1}{\eta^2}\right) K_0\left(\frac{1}{\eta^2}\right) - \frac{1}{\eta^2} K_1\left(\frac{1}{\eta^2}\right) \right] , \qquad (59)$$

where K_0 and K_1 are the modified Bessel functions of the second kind. In the case of very strong magnetic field $\eta > \xi_{\parallel}\sqrt{\tau}$ (or $\omega_c > k_{\max}\lambda_{D\perp}$) the function $\mathcal{F}(\tau,\eta)$ reads

$$\mathcal{F}(\tau,\eta) \simeq \Psi(\xi_{\parallel}) = \ln(1+\xi_{\parallel}^2) - \frac{\xi_{\parallel}^2}{1+\xi_{\parallel}^2} \,. \tag{60}$$

The physical origin of such an anomalous friction coefficient may be traced to the spiral motion of the electrons along the magnetic field lines. These electrons naturally tend to couple strongly with long-wavelength fluctuations (i.e., small k_{\parallel}) along the magnetic field. In addition, when such fluctuations are characterized by slow variation in time (i.e., small $\omega = \mathbf{kV}$), the contact time or the rate of energy exchange between the electrons and the fluctuations will be further enhanced. In a plasma such low-frequency fluctuations are provided by the slow projectile ion. The above coupling can therefore be an efficient mechanism of energy exchange between the electrons and the projectile ion. In the limit of $V \to 0$, the frequency $\omega = \mathbf{kV} \to 0$ tends to zero as well. The contact time thus becomes infinite and the friction coefficient diverges.

The anomalous friction coefficient (see Eq. (56)) vanishes, however, when the ion moves along the magnetic field ($\vartheta = 0$). Then the friction coefficient is solely given by the second term of Eq. (53). The contribution of this term to the stopping power leads to the usual friction law in plasma and reads for arbitrary angles ϑ

$$S \simeq \left(\frac{2}{\pi}\right)^{1/2} \frac{2Z^2 e^2}{\lambda_{D\parallel}^2} \frac{V}{v_{\text{th}\parallel}} \int_0^{\xi_\parallel} k^3 dk \int_0^1 \frac{d\mu}{\mu} \frac{E_1(k,\mu)}{[k^2 + E_2(k,\mu)]^2}$$
(61)

$$\times \left[\mu^2 \cos^2 \vartheta + \frac{1}{2} \left(1 - \mu^2\right) \sin^2 \vartheta\right]$$

with

$$E_1(k,\mu) = \sum_{n=1}^{\infty} \Lambda_n(z) \exp\left(-\frac{n^2 \eta^2}{2k^2 \mu^2}\right) \left[1 + \left(\frac{1}{\tau} - 1\right) \frac{n^2 \eta^2}{k^2 \mu^2}\right]$$
(62)

and $E_2(k,\mu)$ as defined by Eq. (55).

In Figs. (10) and (11) we compare the anomalous term $S_{\rm an}$ with the low velocity stopping without magnetic field S_0 see Eq. (19), where $S_{\rm an}/S_0$ is plotted as a function of ω_c/ω_p for $\vartheta = \pi/6$ (solid line), $\vartheta = \pi/3$ (dotted line), $\vartheta = \pi/2$ (dashed line), $\mathcal{Z} = 0.1$, $V/\overline{v}_{\rm th} = 0.2$, and for two values of the anisotropy parameter τ : $\tau = 0.1$ (Fig. (10)), $\tau = 10$ (Fig. (11)).

We conclude that the anomalous term $S_{\rm an}$ gives espessially for strong magnetic fields $(\omega_c > \omega_p)$ and for strongly temperature anisotropic plasma $(T_{\perp} \gg T_{\parallel})$ an important contribution to the stopping. It should be noted that the observed enhancement of stopping due to $S_{\rm an}$ for $T_{\perp} \gg T_{\parallel}$ can be potentially interesting for future electron cooling experiments.

VII SUMMARY

The purpose of this work was to analyze the stopping power of an ion in temperature anisotropic magnetized classical plasma. A general expression obtained for stopping power was analyzed in four particular cases: in a plasma without magnetic field; in a plasma with weak and very strong magnetic fields; and in a plasma with arbitrary magnetic field and for low-velocity projectile.

From the results obtained in Secs. III-V, we found that the stopping power essentially depends on the plasma temperature anisotropy. In field-free case and for small ion velocities the effect of the anisotropy results in an enhancement of the stopping power when the ion moves in the direction with low temperature.

For small projectile velocities a weak magnetic field slightly decreases the field-free stopping power for small τ , in the opposite case (large τ) the field-free stopping power slightly increases. In the high-velocity limit correction to the field-free stopping power for weak magnetic fields is always negative and the stopping power is reduced by the magnetic field.

In the case of strong magnetic fields we demonstrated an enhancement of the stopping power with increasing of ϑ for low and high-velocity regions compared to the case of an ion which moves along **B**₀.

In low-velocity limit but for arbitrary magnetic field, we find an enhanced stopping power compared to the field-free value mainly due to the strong coupling between the spiral motion of the electrons and the long-wavelength, low-frequency fluctuations excited by the projectile ion. This anomalous stopping power increases with the angle ϑ (the angle between ion velocity **V** and magnetic field **B**₀) and depends strongly on the temperature anisotropy $\tau = T_{\perp}/T_{\parallel}$, as seen in Figs. (10) and (11). Although the nature of the anomalous stopping power is conditioned by the external magnetic field the temperature anisotropy of the plasma can intensify this effect when $T_{\perp} \gg T_{\parallel}$ (see Fig. (11)).

This emphasizes the importance of the special role of fluctuations with small k_{\parallel} and small ω (small projectile velocity V) and as another significant contribution to the energy exchange processes arising from the collective modes of plasma. Potentially, the electron plasma waves and the ion acoustic waves in a magnetized plasma might provide a significant energy-exchange mechanism between projectile ion and plasma particles. This fact then makes it necessary to consider the influence of plasma collective modes to anomalous stopping process. This problem will be treated in a subsequent work.

ACKNOWLEDGMENTS

Finally, it is pleasure to thank Prof. Christian Toepffer for helpful discussions. One of the authors (H.B.N.) is grateful to Prof. Christian Toepffer for hospitality at the Institut für Theoretische Physik II, Universität Erlangen-Nürnberg, where this work was concluded and would like to thank the Deutscher Akademischer Austauschdienst for financial support. We are indebted to Claudia Schlechte for her help in preparing the manuscript. Here we describe the evaluation of the dielectric function in the temperature anisotropic case where the velocity distribution of the unperturbated distribution function was given by Eq. (4). We next introduce the Fourier transformations of $f_1(\mathbf{r}, \mathbf{v}, t)$ with respect to variables \mathbf{r} and t, $f_1(\mathbf{k}, \omega, \mathbf{v})$. Because of the cylindrical symmetry (around the magnetic field direction $\mathbf{b} = \mathbf{B}_0/B_0 = \hat{\mathbf{z}}$) of the problem, we choose

$$\mathbf{v} = v_{\perp} \cos \sigma \hat{\mathbf{x}} + v_{\perp} \sin \sigma \hat{\mathbf{y}} + v_{\parallel} \hat{\mathbf{z}}.$$
 (1)

Then the Vlasov Eq. (2) for the distribution function becomes

$$\frac{\partial}{\partial\sigma}f_1(\mathbf{k},\omega,\mathbf{v}) + \frac{i}{\omega_c}(\mathbf{k}\mathbf{v} - \omega - i0)f_1(\mathbf{k},\omega,\mathbf{v}) = -\frac{ie}{m\omega_c}\varphi(\mathbf{k},\omega)\left(\mathbf{k}\frac{\partial f_0}{\partial\mathbf{v}}\right),\tag{2}$$

where $\varphi(\mathbf{k}, \omega)$ is the Fourier transformation of $\varphi(\mathbf{r}, t)$. The positive infinitesimal +i0 in Eq. (A2) serves to assure the adiabatic turning on of the disturbance and thereby to guarantee the causality of the response. The solution of the Eq. (A2) has the form

$$f_1(\mathbf{k},\omega,\mathbf{v}) = -\frac{ie}{m\omega_c}\varphi(\mathbf{k},\omega)\int\limits_{\infty}^{\sigma} d\sigma_2 \left(\mathbf{k}\frac{\partial f_0}{\partial \mathbf{v}}\right)_{\sigma=\sigma_2} \exp\left[\frac{i}{\omega_c}\int\limits_{\sigma}^{\sigma_2} d\sigma_1 \left[-\omega - i0 + (\mathbf{k}\mathbf{v})_{\sigma=\sigma_1}\right]\right].$$
 (3)

Combining Eq. (A3) with the Poisson equation (3) we find for the dielectric function

$$\varepsilon(\mathbf{k},\omega) = 1 - \frac{4\pi i e^2}{m\omega_c k^2} \int_0^\infty v_\perp dv_\perp \int_0^{2\pi} d\sigma \int_{-\infty}^{+\infty} dv_\parallel \int_\infty^\sigma d\sigma_2 \left[k_\parallel \frac{\partial f_0}{\partial v_\parallel} + k_\perp \cos(\varphi - \sigma_2) \frac{\partial f_0}{\partial v_\perp} \right] \\ \times \exp\left[\frac{i}{\omega_c} \int_\sigma^{\sigma_2} d\sigma_1 \left[k_\parallel v_\parallel - \omega - i0 + k_\perp v_\perp \cos(\varphi - \sigma_1) \right] \right], \tag{4}$$

where $k_x = k_{\perp} \cos \varphi$, $k_y = k_{\perp} \sin \varphi$. After integration by the variables σ_1 , σ_2 and σ , and using the expression [20]

$$\exp(-iz\sin\theta) = \sum_{n=-\infty}^{+\infty} J_n(z)\exp(-in\theta),\tag{5}$$

where J_n is the Bessel function of the *n*th order, we obtain the expression [17]

$$\varepsilon(\mathbf{k},\omega) = 1 - \frac{8\pi^2 e^2}{mk^2} \sum_{n=-\infty}^{+\infty} \int_0^\infty v_\perp dv_\perp \int_{-\infty}^{+\infty} dv_\parallel \left(\frac{n\omega_c}{v_\perp} \frac{\partial f_0}{\partial v_\perp} + k_\parallel \frac{\partial f_0}{\partial v_\parallel}\right) \frac{J_n^2(k_\perp v_\perp/\omega_c)}{n\omega_c + k_\parallel v_\parallel - \omega - i0}.$$
 (6)

Substituting Eq. (4) for the unperturbed distribution function f_0 into Eq. (A6) we finally results in

$$\varepsilon(\mathbf{k},\omega) = 1 + \frac{1}{k^2 \lambda_{D\parallel}^2} \left\{ 1 + \sum_{n=-\infty}^{+\infty} \left(1 + \frac{T_{\parallel}}{T_{\perp}} \frac{n\omega_c}{\omega - n\omega_c} \right) \left[W\left(\frac{\omega - n\omega_c}{|k_{\parallel}|v_{\rm th\parallel}}\right) - 1 \right] \Lambda_n(\beta) \right\}, \quad (7)$$

where $\beta = k_{\perp}^2 v_{\text{th}\perp}^2 / \omega_c^2 = k_{\perp}^2 a_c^2$, $\Lambda_n(z) = \exp(-z) I_n(z)$, $I_n(z)$ is the modified Bessel function of the *n*th order, and W(z) is the plasma dispersion function [18].

To show the identity of the two forms (Eqs. (6) and (A7)) of the dielectric function we will use the expansion in modified Bessel functions [20]

$$\exp(z\cos\theta) = \sum_{n=-\infty}^{\infty} I_n(z)\exp(in\theta).$$
(8)

This allows to rewrite $\exp[-X(t)]$ with X(t) from Eq. (7) as

$$\exp[-X(t)] = \exp(-t^2 \cos^2 \alpha) \sum_{n=-\infty}^{+\infty} \Lambda_n(\beta) \exp\left(\frac{in\omega_c t\sqrt{2}}{kv_{\text{th}\parallel}}\right).$$
(9)

Substituting Eq. (A9) into expression (6) and integration over the variable t leads to Eq. (A7).

В

We now give a more detail derivation of the anomalous term S_{an} (Eq. (56)). We start with the expression (see Eq. (54))

$$Q(k,\varphi,\lambda) = \int_{0}^{1} \frac{d\mu}{\mu} \Phi(\mu,k,\varphi) \exp\left(-\frac{\lambda^2 \phi^2(\mu,\varphi)}{2\mu^2}\right),\tag{1}$$

where $\phi(\mu, \varphi) = \cos \Theta$, $\lambda = V/v_{\text{th}\parallel}$,

$$\Phi(\mu, k, \varphi) = \frac{\Lambda_0(z) \cos^2 \Theta}{\left[k^2 + E_2(k, \mu)\right]^2}.$$
(2)

For $\lambda \to 0$ a leading-term approximation of (B1) leads to

$$Q(k,\varphi,\lambda) \simeq \Phi(0,k,\varphi) \ln \frac{\sqrt{2}}{\lambda |\phi(0,\varphi)| \sqrt{\gamma}} + \mathcal{O}(1),$$
(3)

where γ is the Euler's constant, $|\phi(0,\varphi)| = \sin \vartheta |\cos \varphi|$,

$$\Phi(0,k,\varphi) = \frac{\Lambda_0(k^2\tau/\eta^2)\sin^2\vartheta\cos^2\varphi}{\left[k^2 + E_2(k,0)\right]^2}$$
(4)

and

$$E_2(k,0) = 1 + 2\left(\frac{1}{\tau} - 1\right) \sum_{n=1}^{\infty} \Lambda_n(k^2 \tau / \eta^2).$$
 (5)

Using the relation [17, 20]

$$\sum_{n=-\infty}^{+\infty} \Lambda_n(z) = 1, \tag{6}$$

the function $E_2(k,0)$ we finally takes the form

$$E_2(k,0) = \frac{1}{\tau} + \left(1 - \frac{1}{\tau}\right) \Lambda_0(k^2 \tau / \eta^2).$$
(7)

Substituting Eqs. (B3), (B4) and (B7) into Eq. (54) and integration over φ we finally come to expression (56).

REFERENCES

- [1] A. H. Sørensen and E. Bonderup, Nucl. Instrum. Methods **215**, 27 (1983).
- [2] H. Poth, Phys. Reports **196**, 135 (1990).
- [3] I. N. Meshkov, Phys. Part. Nucl. **25**, 631 (1994).

- [4] Proceedings of the 12th International Symposium on Heavy Ion Inertial Fusion, (Heidelberg, Germany, Sept. 1997), Nucl. Instrum. Methods A 415, (1998).
- [5] I. A. Akhiezer, Zh. Eksp. Teor. Fiz. 40, 954 (1961) [Sov. Phys. JETP 13, 667 (1961)].
- [6] N. Honda, O.Aona, and T.Kihara, J. Phys. Soc. Jpn. 18, 256 (1963).
- [7] R. M. May and N. F. Cramer, Phys. Fluids **13**, 1766 (1970).
- [8] G. G. Pavlov and D. G. Yakovlev, Zh. Eksp. Teor. Fiz. 70, 753 (1976) [Sov. Phys. JETP 43, 389 (1976)].
- [9] J. G. Kirk and D. I. Galloway, Plasma Phys. 24, 339 (1982).
- [10] S. V. Bozhokin and É. A. Choban, Fiz. Plazmy 10, 779 (1984) [Sov. J. Plasma Phys. 10, 452 (1984)].
- [11] H. B. Nersisyan, Phys. Rev. E 58, 3686 (1998).
- [12] H. B. Nersisyan and C. Deutsch, Phys. Lett. A **246**, 325 (1998).
- [13] C. Seele, G. Zwicknagel, C. Toepffer, and P.-G. Reinhard, Phys. Rev. E 57, 3368 (1998).
- [14] M. Walter, C. Toepffer, and G. Zwicknagel, Nucl. Instrum. Meth. B, (1999) (to be published).
- [15] E. Lifshitz and L. P. Pitaevskij, *Physical Kinetics* (Pergamon Press, Oxford, 1981).
- [16] G. Zwicknagel, C. Toepffer, and P.-G. Reinhard, Phys. Reports 309, 117 (1999).
- [17] S. Ichimaru, Basic Principles of Plasma Physics (Benjamin, Reading, MA 1973), Sec. 7.4.
- [18] D. B. Fried and S. D. Conte, *The Plasma Dispersion Function* (Academic Press, New York, 1961).
- [19] Th. Peter and J. Meyer-ter-Vehn, Phys. Rev. A 43, 1998 (1991).
- [20] I. S. Gradshteyn and I. M. Ryzhik, Table of Integrals, Series and Products (Academic, New York, 1980).