

Determination of meteor showers on other planets using comet ephemerides

Shane L. Larson

Department of Physics, Montana State University, Bozeman, Montana 59717

ABSTRACT

Meteor showers on the Earth occur at well known times, and are associated with the decay of comets or other minor bodies whose orbital paths pass close to the Earth's trajectory. On the surface, determining the closest proximity of two orbital paths appears to be a computationally intensive procedure. This paper describes a simple geometric method for determining the proximity between the orbital paths of two bodies (*i.e.*, a comet and a planet) in the solar system from the known ephemerides of the objects. The method is used to determine whether or not meteor showers on other planets in the solar system could be associated with any of 250 known comets.

Subject headings: comets: general — meteors — solar system: general

1. INTRODUCTION

As they traverse their orbits about the Sun, comets slowly evaporate and fragment, leaving small bits of cometary debris along their orbital tracks. Some comet orbits intersect the Earth's path, and the planet sweeps up a portion of these particulates each year. Generally, these particles are drawn into the atmosphere, where they burn up at high altitudes, producing the yearly meteor showers. A sample of the meteor showers expected on a regular basis for Earth-bound observers is given in Table 1. A very detailed list of meteor streams encountered by the Earth has been composed based on ground-based observations of amateur astronomers around the world (Jenniskens 1994).

Given the large number of meteor showers seen on the Earth, it seems natural to ask about the possibility of meteor showers on other planets. It may be impractical for a sky-observer of the future to view meteor showers from some worlds: Mercury has no atmosphere, the clouds of Venus are so thick most meteors will likely burn up before a planetbound observer could see them, Jupiter has no solid surface to sit on while viewing the shower, and so forth. Never-the-less, predict-

ing regular meteor showers on other worlds may be important for protecting explorers and spacecraft from incoming particles, and could be useful for planning expeditions and experiments to collect cometary material. It has been suggested (Adolfsson *et al.* 1996) that future Mars landers may be able to detect meteors from the surface of Mars.

A great deal of modern research has been devoted to analysis of the evolution of meteor streams in the solar system, particularly those that intersect the Earth's orbit (for example, detailed analyses of the evolution of the Quadrantid stream can be found in (Hughes *et al.* 1979), (Williams *et al.* 1979) and (Murray *et al.* 1980); the Geminid stream is analyzed in (Fox *et al.* 1982) and (Jones 1985)). These analyses take into account perturbations to the orbits of the parent bodies, as well as the subsequent evolution of the debris trail after the comet or minor body has continued on in its orbit. Over time, streams may wander into a planet's path causing new meteor storms, or may wander out the planet's path quenching a shower which has been periodic for decades or centuries (*e.g.*, (Murray *et al.* 1980) estimates that the Quadrantid shower will vanish by the year 2100).

To a first approximation, however, meteor showers will occur if the orbit of a planet and the orbit of a minor body intersect (or pass close to one another). One way to determine if this occurs is to evolve the two orbits on a computer and watch for an intersection. A generalized method for finding the minimum separation between two orbits has been described by Murray, Hughes and Williams (1980), but the method reduces to a coupled set of equations for the orbital anomalies which requires numerical solution. Alternatively, the methods described in this paper approach the problem of determining the intersection of orbital planes in a completely analytical fashion, requiring only geometrical methods and matrix algebra.

Section 2 describes the basic parameters and coordinate systems used to characterize orbits in this paper. Section 3 describes the rotations used to correctly orient two orbits with respect to each other, and applies the rotations to essential vectors needed for the analysis. Section 4 uses the rotated vectors to determine the intersection between two orbital planes, and computes the distance between the orbital paths when the planes intersect. Section 5 proposes a criteria for the existence of a meteor shower based on the distance between the orbits at intersection. The “time” of showers meeting the criterion is determined. Section 6 applies the condition of Section 5 to 250 known comets, summarizes the results, and discusses the limitations of determining meteor showers using this method.

Throughout this paper, SI (*Système Internationale*) units are employed, except where the size of the units makes it convenient to work in standard units employed in astronomy (*e.g.*, on large scales, astronomical units (AU) will be used, rather than meters).

2. Describing Orbits

As is well known, one of the great discoveries of Johannes Kepler was that the planets travel on elliptical paths, with the Sun at one focus of the ellipse (Kepler’s First Law of Planetary Motion, published in 1609). Since then, an enormous body of knowledge has been developed regarding the analysis of orbital motion (see, for example (Marion & Thornton 1988)), allowing the determination of the position of virtually any object

in the solar system at any moment in time.

For the work presented here, a time dependent analysis of the orbital motion is not necessary ¹. The only information which is required is a knowledge of the trajectory of the orbit through space. The distance of the orbital path from the Sun may be written for elliptical orbits as

$$r = \frac{a(1 - e^2)}{1 + e \cos \theta} , \quad (1)$$

where a is the semi-major axis of the orbit, e is the eccentricity, and θ is the angle (the anomaly) between the body and the axis defined by perihelion, as measured in the orbital plane. The perihelion distance for the object can be found from Eq. (1) by taking $\theta = 0$, yielding

$$r_p = a(1 - e) . \quad (2)$$

The distance expressed in Eq. (1) describes the correct size and shape of an elliptical orbit for any object around the Sun, but more information is needed to correctly orient the orbit in three-dimensional space. This information is typically collected in the orbital ephemeris, which expresses orbital parameters with respect to the plane which is coincident with the orbital plane of the Earth (the ecliptic). This paper will use a (cartesian) reference coordinate system defined in the ecliptic plane as shown in Figure 1. The $+z$ axis is defined perpendicular to the ecliptic and in the right handed sense with respect to the Earth’s orbital motion (*i.e.*, when viewed looking down the $+z$ axis, the Earth’s motion is counter-clockwise in the xy -plane). The $+x$ axis is defined along the direction of the Earth’s perihelion.

The orbital ephemeris of any body describes its orbit relative to the ecliptic plane, and locates the object along its orbital path as a function of time. For the problem of determining the possible intersection of two orbital paths, only three elements of the full ephemeris for a body will be needed: Ω_o (a *modified* longitude of the ascending node), i (inclination), and ω (argument of perihelion). Each of these parameters is described below, and shown in Figure 2.

¹We are interested in knowing the proximity of two orbits when they cross. An interesting (but ultimately more difficult) question to address is whether two bodies might actually *collide* because their orbits intersect.

The longitude of the ascending node, Ω , is the angle in the ecliptic plane between the vernal equinox (the first point of Aries) and the point at which the orbit crosses the ecliptic towards the $+z$ direction (“northward” across the ecliptic). The parameter, Ω_o , used in this paper, is an offset longitude measured from the perihelion of Earth, rather than the first point of Aries (see Figure 3). The inclination, ι , is the angle between the normal vector of the orbit and the normal vector of the ecliptic. Lastly, the argument of perihelion, ω , is the angle between the position of the body as it crosses the ascending node and the position at perihelion, as measured *in the orbital plane of the body*.

In addition to these three angles, it will be useful to define two vectors for each orbit of interest: \hat{n} , the unit normal vector to the plane of the orbit, and \vec{r}_p , the vector pointing to perihelion in the plane of the orbit.

3. Rotations for Orbital Orientation

In order to correctly orient an orbit with respect to the ecliptic, assume (initially) that the orbit of interest is in the plane of the ecliptic, with the perihelion of the orbit aligned along the $+x$ axis (*i.e.*, the orbit is co-aligned with the Earth’s orbit). A series of three rotations, based on the angles $\{\omega, \iota, \Omega\}$ from the orbital ephemeris will produce the correct orientation. The first rotation will set the value of the ascending node with respect to perihelion, the second rotation will set the inclination to the ecliptic, and the third rotation will move the ascending node to the correct location in the ecliptic plane.

A useful method for describing rotations is in terms of matrices. While it is possible to construct a rotation matrix for rotations about a general axis, it is more convenient to conduct rotations about the coordinate axes shown in Figure 1. The matrices describing rotations about the x -, y -, and z -axes will be denoted $\tilde{M}_x(\phi)$, $\tilde{M}_y(\xi)$, and $\tilde{M}_z(\psi)$, respectively.

To demonstrate the rotations needed to orient the orbit, consider a general vector, \vec{A} , which is rigidly attached to the orbital plane, maintaining its orientation as the plane is rotated.

The first rotation locates the ascending node with respect to perihelion; the rotation depends on

the value of the argument of perihelion, ω . This is done by rotating around the z -axis by $\psi = \omega$. In terms of rotating a general vector \vec{A} , this can be written

$$\vec{A}_1 = \tilde{M}_z(\omega)\vec{A}. \quad (3)$$

When this operation is applied to the orbit, the ascending node will be located on the $+x$ axis.

The orbit is inclined around an axis which passes through the ascending node and through the Sun (at one focus of the orbit). Since the first rotation placed the ascending node on the $+x$ axis, and the Sun lies at the origin of coordinates, a rotation around the x -axis by the inclination angle, $\phi = \iota$, will correctly incline the orbit. In terms of the vector \vec{A}_1 (resulting from Eq. (3)), this yields

$$\vec{A}_2 = \tilde{M}_x(\iota)\vec{A}_1. \quad (4)$$

Before the final rotation, it will be convenient to offset the longitude of the ascending node such that it is measured from the perihelion of the Earth, rather than the vernal equinox (this makes the x -axis the origin for measuring the longitude of the ascending node). The angle between the vernal equinox and perihelion of Earth is simply the argument of perihelion for Earth, ω_\oplus , giving (see Figure 3)

$$\Omega_o = 2\pi - \omega_\oplus + \Omega. \quad (5)$$

After the second rotation, the ascending node is still located on the $+x$ axis. Rotation about the z -axis by the offset longitude, $\psi = \Omega_o$, will rotate the longitude of the ascending node to its correct location in the ecliptic plane. In terms of the vector \vec{A}_2 (resulting from Eq. (4)), this yields

$$\vec{A}_3 = \tilde{M}_z(\Omega_o)\vec{A}_2. \quad (6)$$

The vector \vec{A}_3 (which is rigidly attached to the orbit) is correctly oriented with respect to the ecliptic.

The two vectors which will be of use later are the unit normal vector to the orbit, \hat{n} , and the perihelion vector, \vec{r}_p . When the orbital plane is co-aligned with the Earth’s (before any rotations have been performed), these vectors have the form

$$\hat{n} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \quad \vec{r}_p = \begin{bmatrix} r_p \\ 0 \\ 0 \end{bmatrix}. \quad (7)$$

The rotation operations described by Eqs. (3), (4), and (6) must be applied to these vectors so they correctly describe the orbit with respect to the ecliptic. Conducting the rotation procedure yields

$$\hat{n}' = \begin{bmatrix} \sin \iota \sin \Omega_o \\ -\sin \iota \cos \Omega_o \\ \cos \iota \end{bmatrix}, \quad (8)$$

and

$$\vec{r}_p' = r_p \begin{bmatrix} \cos \omega \cos \Omega_o - \sin \omega \cos \iota \sin \Omega_o \\ \cos \omega \sin \Omega_o + \sin \omega \cos \iota \cos \Omega_o \\ \sin \omega \sin \iota \end{bmatrix}. \quad (9)$$

4. Intersection of Orbits

The procedure described in Section 3 will correctly orient any orbit with respect to the ecliptic. One could take any planet's ephemeris (*e.g.*, from the ephemerides given in Table 2) and construct the normal vector \hat{n} and perihelion vector \vec{r}_p in accordance with Eqs. (8) and (9)². Similar vectors could be generated for cometary ephemerides.

The real question of interest is not how the orbital planes of planets and comets are related to the ecliptic, but rather how they are oriented with respect to each other, and in particular where they intersect. The line defining the intersection of the orbital planes can be used to determine whether or not the orbits actually intersect.

Hereafter, assume that vectors related to a comet's orbit will bear the subscript 'c' and vectors related to a planet's orbit will bear the subscript '+'. Further, suppose the components of the normal vector for a comet's orbit are $\hat{n}_c = (a, b, c)$, and the components of the normal vector of a planet's orbit are $\hat{n}_+ = (e, f, g)$. Both orbital planes automatically share one point in common: the origin, which lies at the focus of each orbital ellipse. Given this point and the two vectors \hat{n}_c and \hat{n}_+ , the equations describing the two orbital planes are

$$\begin{array}{ll} \text{CometPlane} & ax + by + cz = 0 \\ \text{PlanetPlane} & ex + fy + gz = 0 \end{array} \quad (10)$$

²To ease the notation, we will drop the primed notation for rotated vectors from here on. It will be understood that the normal vectors and perihelion vectors have been correctly oriented with respect to the ecliptic.

The intersection of the two planes is a line which is the common solution of the two expressions in Eq. (10). Using determinants, the common solution to these equations is found to be

$$\frac{x}{\begin{vmatrix} b & c \\ f & g \end{vmatrix}} = \frac{-y}{\begin{vmatrix} a & c \\ e & g \end{vmatrix}} = \frac{z}{\begin{vmatrix} a & b \\ e & f \end{vmatrix}} = k, \quad (11)$$

where k is an arbitrary constant. The solutions $\{x, y, z\}$ of Eq. (11) will be points along the line of intersection. It is useful to use these values to define a new vector, $\vec{\lambda}$, called the 'node vector.' It points along the line of nodes (the intersection of the two planes), and has components

$$\vec{\lambda} = k \begin{bmatrix} bg - cf \\ ce - ag \\ af - be \end{bmatrix}. \quad (12)$$

To determine if the orbital paths intersect, one must know the radii of the orbits along the line of nodes. An orbital radius may be determined from Eq. (1) if the value of the anomaly, θ , is known. In terms of two orbits inclined with respect to each other, the angles of interest will be the angle between the perihelion vector for each orbit, \vec{r}_p , and the node vector, $\vec{\lambda}$. For each orbit, the angle is defined in terms of the dot product of the two vectors, yielding

$$\cos \theta = \frac{\vec{r}_p \cdot \vec{\lambda}}{|\vec{r}_p| \cdot |\vec{\lambda}|}. \quad (13)$$

The orbits have two opportunities to intersect: at the ascending node, and at the descending node. Eq. (13) gives the angle at a single node. To obtain the value of the anomaly at the other node, dot the perihelion vector, \vec{r}_p , into the negative of the node vector, $-\vec{\lambda}$.

Once the anomaly is known, the distance between the orbital paths when the planes intersect is simply

$$\Delta = |r_+ - r_c|, \quad (14)$$

where r_+ and r_c are computed using Eq. (1) with the anomaly defined by Eq. (13) and the appropriate orbital parameters derived from tabulated ephemerides.

5. Is there a meteor shower?

The occurrence of a meteor shower associated with a particular comet will depend on the value of the separation between the orbital paths, Δ . A variety of proximity criteria could be developed, depending on specific considerations one would like to make regarding the likelihood that particles from a given stream might reach a planetary body (such as the proximity of the mean orbits of a comet and planet, or the spread of cometary particles around the orbit of the parent comet). The basic criteria may be expressed such that

$$\Delta \leq \hat{r}, \quad (15)$$

where \hat{r} is the minimum separation one considers likely for a shower to occur.

In this paper, the criteria for an orbit intersection causing a meteor shower will be scaled to a region where the gravitational potential of the planet dominates over the gravitational potential of the Sun. The criteria will be

$$\Delta \leq \kappa R_l, \quad (16)$$

where R_l is the ‘‘Roche-lobe radius’’ (defined as the radius of a sphere which has the same volume as the planet’s Roche lobe) and κ is an arbitrary adjustable scaling factor. The Roche-lobe radius can be approximated by

$$R_l \sim 0.52 \cdot a \left[\frac{m_+}{M_\odot + m_+} \right]^{0.44}, \quad (17)$$

where m_+ and M_\odot are the mass of the planet and the Sun, and a is the semi-major axis of the planet’s orbit (Iben & Tutukov 1984).

Once an intersection (in the sense of Eq. (16)) has been found, one would like to know *when* the associated meteor shower might occur, such that it is possible to mount an observational effort to detect the shower. A good marker for a planetary encounter with the meteor stream is the value of the planetary anomaly along the line of nodes, given by Eq. (13).

Given a particular value of the anomaly, θ , it is possible to write a closed-form expression $t(\theta)$ for the time at which the planet will arrive at that point in its orbit. The areal velocity may be written

$$\frac{dA}{dt} = \frac{\pi ab}{\tau}, \quad (18)$$

where a and b are the semi-major and -minor axes respectively, and τ is the period of the orbit. Separating Eq. (18) and writing the area element as $dA = (1/2)r^2 d\theta'$ allows one to write an integral for the time t as

$$t(\theta) = \left(\frac{\tau}{\pi ab} \right) \frac{1}{2} \int_0^\theta r^2 d\theta'. \quad (19)$$

Using the shape equation (Eq. (1)) to write $r = r(\theta)$, and using $b = a(1 - e^2)^{1/2}$ to express the semi-minor axis, Eq. (19) can be integrated to give

$$t(\theta) = \frac{\tau}{2\pi} \left[2 \tan^{-1} \left(\sqrt{\frac{1-e}{1+e}} \tan \frac{\theta}{2} \right) - \frac{e\sqrt{1-e^2} \sin \theta}{1+e \cos \theta} \right], \quad (20)$$

which is the time it takes the planet to pass from perihelion ($\theta = 0$) to the anomaly value θ .

6. Results & Discussion

As an example of the methods presented here, a search for comet-planet orbital intersections in the solar system was carried out using the planetary ephemerides shown in Table 2 (Standish *et al.* 1992), and the comet ephemerides provided in the Jet Propulsion Laboratory’s DASTCOM (Database of ASTeroids and COMets) (JPL DASTCOM 1999). The DASTCOM is a collection of orbital parameters and physical characteristics for the numbered asteroids, unnumbered asteroids, and periodic comets, used for analyses of solar system dynamics. The DASTCOM includes all known comets with periods less than 200 years (periodic comets), and long period comets which have made a return after 1995 (there are currently 73 long period comets in the DASTCOM, and 208 comets with periods less than 200 years). The orbital elements given in the DASTCOM are osculating elements, computed from observations during the comet’s most recent apparition.

The results of the search for comet-planet orbital intersections are listed in Table 3, which specified an encounter distance of

$$\Delta \leq 5R_l. \quad (21)$$

In all, 128 possible showers were detected: 3 at Earth, 1 at Mars, 106 at Jupiter³, 17 at Saturn,

³In fact, the results of Table 3 show that Jupiter’s orbit

and 1 at Uranus. If one reduces the encounter distance to $\Delta \leq 1R_l$, only 32 possible showers are detected (shown at the top of Table 3): 1 at Earth, 28 at Jupiter, 2 at Saturn, and 1 at Uranus. If one allows the encounter distance to expand to $\Delta \leq 10R_l$, 188 possible showers are detected (data not shown in Table 3): 4 at Earth, 5 at Mars, 148 at Jupiter, 24 at Saturn, 6 at Uranus, and 1 at Neptune.

Comets with orbital scales smaller than the solar system ('short period comets') have evolved largely under the influence of perturbations due to Jupiter (the mass of Jupiter is greater than the mass of the other planets combined), giving a large population of comets which cross Jupiter's orbit. The search for the origin of these "Jovian family comets" has been a matter of much numerical simulation and debate (see, for example, (Quinn *et al.* 1990)). The disproportionately large number of showers detected for Jupiter can be attributed to this feature of the comet population.

A good check of the procedure described in this paper is to consider the predicted showers at Earth. In particular, the method outlined in this work predicts two meteor streams which can be identified with known showers. The first is the stream from Comet Tempel-Tuttle, occurring at $t \sim 318$ d. This stream can be identified with the Leonid meteor shower (known to be a stream from Tempel-Tuttle), which occurs in mid-November each year. The second is a stream from Comet Swift-Tuttle, occurring at $t \sim 221$ d. This stream can be identified with the Perseid meteor shower (known to be a stream from Swift-Tuttle), which occurs in mid-August each year.

The possible showers computed here have all assumed that the orbits of the comets are static and do not precess. Further, it is assumed that the meteor streams remain attached to those static orbits without wandering under the influence of gravitational perturbations in the solar system. In addition, the influences of 'local' bodies around each planet (*e.g.*, Earth's moon, or the Galilean satellites around Jupiter) have been ignored. Never-

intersects the path of comet P/Spahr (1998 U4) *twice*: one intersection at a separation of $\Delta \simeq 1.6R_l$, and a second intersection (at the other node) with a separation of $\Delta \simeq 3.3R_l$.

theless, the method provides a useful way for determining the possibility that a given planet will encounter a meteor stream from minor bodies in the solar system.

I would like to thank M. B. Larson for comments and suggestions, and E. M. Standish who provided helpful discussions regarding planetary ephemerides. D. K. Yeomans and P. Chodas provided helpful details about the DASTCOM database. I would also like to acknowledge the hospitality of the Solar System Dynamics Group at the Jet Propulsion Laboratory during the time this work was completed. This work was supported in part by NASA Cooperative Agreement No. NCC5-410.

REFERENCES

- Adolfsson, L. G., Gustafson, B. A. S. and Murray, C. D., *Icarus*, 1996, **119**, 144
- DASTCOM (Database of Asteroids and Comets), Jet Propulsion Laboratory, 1999, <http://ssd.jpl.nasa.gov/dastcom.html>
- Fox, K. A., Hughes, D. W. and Williams, I. P., *MNRAS*, 1982, **200**, 313
- Hughes, D. W., Williams, I. P. and Murray, C. D., *MNRAS*, 1979, **189**, 493
- Iben, Jr. I. and Tutukov, A. V., *ApJS*, 1984, **54**, 335
- Jenniskens, P., *A&A*, 1994, **287**, 990
- Jones, J., *MNRAS*, 1985, **217**, 523
- Marion, J. B. and Thornton, S. T., *Classical dynamics of particles and systems* (Harcourt Brace Jovanovich, New York, 1988).
- Murray, C. D., Hughes, D. W. and Williams, I. P., *MNRAS*, 1980, **190**, 733
- Quinn, T., Tremaine, S. and Duncan, M., *ApJ*, 1990, **355**, 667
- Schlosser, W., Schmidt-Kaler, T. and Milone, E. F., *Challenges of astronomy: hands-on experiments for the sky and laboratory* (Springer-Verlag, New York, 1991).

Standish, E. M., Newhall, XX, Williams, J. G. and Yeomans, D. K. in *Explanatory supplement to the Astronomical Almanac*, P. K. Seidelmann, Ed. (University Science Books, Mill Valley, CA, 1992).

Williams, I. P., Murray, C. D. and Hughes, D. W., MNRAS, 1979, **189**, 483

This 2-column preprint was prepared with the AAS L^AT_EX macros v5.0.

Fig. 1.— The reference coordinate system in the ecliptic plane. The z -axis is defined in the right-handed sense with respect to the Earth's motion, and the x -axis points towards the perihelion of the Earth.

Fig. 2.— The three essential angles for correctly orienting orbits in three dimensional space are (a) Ω_o , the (*modified*) longitude of the ascending node; (b) ι , the inclination; and (c) ω , the argument of perihelion.

Fig. 3.— The *modified* longitude of the ascending node, Ω_o , defined in terms of the Earth's argument of perihelion, ω_{\oplus} , and the conventional value of Ω for the orbit. v indicates the first point of Aries.

Fig. 4.— The intersection of two orbits, showing the essential quantities for defining the occurrence of a meteor shower: the separation of the orbits at crossing, Δ and the node vector, $\vec{\lambda}$, which defines the intersection of the two orbits and is used to determine the time of the meteor shower.

TABLE 1
SOME YEARLY METEOR SHOWERS SEEN FROM EARTH.

Shower Name	Date
Quadrantids	early January
Lyrids	mid April
η Aquarids	early May
δ Aquarids	late July
Perseids	mid August
Orionids	mid October
Leonids	mid November
Geminids	mid December

TABLE 2
THE MEAN EPHEMERIDES FOR THE PLANETS OF THE SOLAR SYSTEM (EPOCH J2000).

Planet	a (AU)	e	ι ($^{\circ}$)	Ω ($^{\circ}$)	ω ($^{\circ}$)
Mercury	0.38709893	0.20563069	7.00487	48.33167	29.12478
Venus	0.72333199	0.00677323	3.39471	76.68069	54.85229
Earth	1.00000011	0.01671022	0.00005	-11.26064	114.20783
Mars	1.52366231	0.09341233	1.85061	49.57854	286.4623
Jupiter	5.20336301	0.04839266	1.3053	100.55615	-85.8023
Saturn	9.53707032	0.0541506	2.48446	113.71504	-21.2831
Uranus	19.19126393	0.04716771	0.76986	74.22988	96.73436
Neptune	30.06896348	0.00858587	1.76917	131.72169	-86.75034
Pluto	39.348168677	0.24880766	17.14175	110.30347	113.76329

TABLE 3

THE RESULTS OF A METEOR SHOWER SEARCH USING THE COMETS IN THE JPL DASTCOM DATABASE.

Planet	Comet	$(\Delta/R_t)^a$	θ , degrees	$t(\theta)$, d
Earth	109P/Swift-Tuttle	0.5043	216.49717	220.81696
Jupiter	P/LONEOS-Tucker (1998 QP54)	0.02759	143.72224	1688.24555
Jupiter	117P/Helin-Roman-Alu 1	0.05761	54.79245	605.7507
Jupiter	43P/Wolf-Harrington	0.10689	241.59541	2965.94337
Jupiter	P/Hergenrother (1998 W2)	0.13008	158.6528	1883.37507
Jupiter	C/Hale-Bopp (1995 O1)	0.15907	267.71691	3287.20798
Jupiter	78P/Gehrels 2	0.2102	206.19238	2510.80676
Jupiter	75P/Kohoutek	0.21718	256.88227	3155.62992
Jupiter	P/Spahr (1998 W1)	0.24054	267.30471	3282.24601
Jupiter	124P/Mrkos	0.32574	344.80757	4164.72657
Jupiter	14P/Wolf	0.32874	191.77241	2321.02269
Jupiter	53P/Van Biesbroeck	0.37063	143.86166	1690.05544
Jupiter	59P/Kearns-Kwee	0.39771	294.48157	3602.23635
Jupiter	91P/Russell 3	0.41994	56.45964	624.68681
Jupiter	76P/West-Kohoutek-Ikemura	0.44039	248.71583	3054.87529
Jupiter	26P/Grigg-Skjellerup	0.45251	21.59907	236.07478
Jupiter	132P/Helin-Roman-Alu 2	0.50932	176.74286	2122.18424
Jupiter	P/Kushida (1994 A1)	0.54051	239.05142	2933.93263
Jupiter	16P/Brooks 2	0.57961	175.77613	2109.39006
Jupiter	83P/Russell 1	0.59462	34.84717	382.18644
Jupiter	D/Kowal-Mrkos (1984 H1)	0.61747	57.35711	634.89823
Jupiter	135P/Shoemaker-Levy 8	0.68357	28.92759	316.71554
Jupiter	139P/Vaisala-Oterma	0.72789	241.18336	2960.76693
Jupiter	86P/Wild 3	0.73742	55.46464	613.3804
Jupiter	104P/Kowal 2	0.81618	233.8782	2868.46755
Jupiter	P/LINEAR-Mueller (1998 S1)	0.87419	157.76011	1871.64355
Jupiter	P/Shoemaker-Levy 6 (1991 V1)	0.87826	199.12624	2418.00496
Jupiter	18P/Perrine-Mrkos	0.94922	228.33725	2797.82077
Jupiter	85P/Boethin	0.97138	164.48654	1960.19405
Saturn	P/Jager (1998 U3)	0.06673	210.03052	6366.0197
Saturn	126P/IRAS	0.53161	83.17079	2299.91062
Uranus	C/Li (1999 E1)	0.84435	137.45475	11360.58497
Earth	55P/Tempel-Tuttle	4.15373	312.31161	318.28395
Earth	26P/Grigg-Skjellerup	4.518	110.36191	110.13972
Mars	C/LINEAR (1998 U5)	1.98779	90.14579	151.61158
Jupiter	47P/Ashbrook-Jackson	1.00145	162.12665	1929.08903
Jupiter	15P/Finlay	1.00753	186.90971	2256.74179
Jupiter	97P/Metcalf-Brewington	1.09545	175.71325	2108.55792
Jupiter	81P/Wild 2	1.10168	320.59158	3897.71867
Jupiter	121P/Shoemaker-Holt 2	1.10447	264.90153	3253.24816
Jupiter	54P/de Vico-Swift	1.1133	152.93248	1808.3284
Jupiter	56P/Slaughter-Burnham	1.1211	143.93726	1691.03684
Jupiter	P/Korlevic-Juric (1999 DN3)	1.15484	347.38897	4192.9561
Jupiter	P/Mueller 4 (1992 G3)	1.16212	312.27537	3804.66421
Jupiter	52P/Harrington-Abell	1.20057	316.89669	3856.48162
Jupiter	69P/Taylor	1.2392	274.64237	3370.06049
Jupiter	P/Larsen (1997 V1)	1.269	224.13556	2743.90598
Jupiter	46P/Wirtanen	1.31801	245.23688	3011.54806
Jupiter	100P/Hartley 1	1.32008	21.70502	237.23779
Jupiter	87P/Bus	1.34382	15.87435	173.3314
Jupiter	C/Ferris (1999 K2)	1.3836	285.50894	3498.14812
Jupiter	77P/Longmore	1.43731	358.02608	4309.06758
Jupiter	C/Mueller (1997 J1)	1.44651	262.26797	3221.3349
Jupiter	P/Hartley-IRAS (1983 V1)	1.51103	166.77415	1990.37975

TABLE 3—*Continued*

Planet	Comet	$(\Delta/R_1)^a$	θ , degrees	$t(\theta)$, d
Jupiter	119P/Parker-Hartley	1.53129	236.51823	2901.93741
Jupiter	114P/Wiseman-Skiff	1.54424	256.92325	3156.13216
Jupiter	102P/Shoemaker 1	1.58558	143.00259	1678.90784
Jupiter	P/Spahr (1998 U4)	1.61195	169.07697	2020.79437
Jupiter	33P/Daniel	1.64668	229.96736	2818.65939
Jupiter	62P/Tsuchinshan 1	1.6969	261.53064	3212.37462
Jupiter	70P/Kojima	1.78536	288.99826	3538.80021
Jupiter	60P/Tsuchinshan 2	1.84257	272.21092	3341.08206
Jupiter	4P/Faye	1.91557	192.45922	2330.09358
Jupiter	67P/Churyumov-Gerasimenko	1.95263	207.28864	2525.16175
Jupiter	6P/d'Arrest	2.01111	126.65145	1468.85806
Jupiter	36P/Whipple	2.02544	175.2223	2102.06112
Jupiter	C/LINEAR (1998 U1)	2.08822	12.83484	140.08603
Jupiter	C/Spacewatch (1997 BA6)	2.15772	302.67334	3696.04414
Jupiter	P/Levy (1991 L3)	2.21811	131.97724	1536.80051
Jupiter	116P/Wild 4	2.27558	347.04647	4189.21226
Jupiter	P/Shoemaker-Levy 1 (1990 V1)	2.39831	215.04783	2626.37534
Jupiter	9P/Tempel 1	2.41105	50.10947	552.78419
Jupiter	C/LINEAR (1999 H3)	2.59549	138.47987	1620.39097
Jupiter	31P/Schwassmann-Wachmann 2	2.63332	303.86438	3709.59249
Jupiter	P/LINEAR (1999 J5)	2.69947	98.43266	1117.77483
Jupiter	C/LINEAR (1998 W3)	2.71291	108.75604	1244.38557
Jupiter	21P/Giacobini-Zinner	2.73222	182.72852	2201.41502
Jupiter	40P/Vaisala 1	2.84407	304.29294	3714.46212
Jupiter	108P/Cifre	2.88171	214.58058	2620.30119
Jupiter	136P/Mueller 3	2.92721	128.57415	1493.32993
Jupiter	42P/Neujmin 3	2.93144	153.24921	1812.47528
Jupiter	7P/Pons-Winnecke	3.13747	78.25359	876.539
Jupiter	103P/Hartley 2	3.19169	209.67222	2556.32919
Jupiter	128P/Shoemaker-Holt 1-B	3.22265	213.44774	2605.56299
Jupiter	D/van Houten (1960 S1)	3.25993	336.58556	4074.58242
Jupiter	P/Kushida-Muramatsu (1993 X1)	3.26773	250.65642	3078.93931
Jupiter	C/Zhu-Balam (1997 L1)	3.26781	218.82993	2675.43314
Jupiter	P/LINEAR (1998 VS24)	3.26871	158.78542	1885.11843
Jupiter	P/Spahr (1998 U4)	3.29681	349.07697	4211.40146
Jupiter	61P/Shajn-Schaldach	3.3833	164.21702	1956.63978
Jupiter	65P/Gunn	3.43747	49.5381	546.34347
Jupiter	120P/Mueller 1	3.59615	161.60426	1922.20873
Jupiter	129P/Shoemaker-Levy 3	3.6607	284.2452	3483.36888
Jupiter	30P/Reinmuth 1	3.66617	288.52425	3533.29093
Jupiter	22P/Kopff	3.67008	113.53434	1303.71706
Jupiter	112P/Urata-Nijima	3.69814	194.43253	2356.14237
Jupiter	110P/Hartley 3	3.70194	272.28798	3342.00224
Jupiter	49P/Arend-Rigaux	3.78256	288.50492	3533.06618
Jupiter	P/Lagerkvist (1996 R2)	3.9786	175.43706	2104.90302
Jupiter	19P/Borrelly	4.00335	239.67933	2941.845
Jupiter	17P/Holmes	4.00357	130.61884	1519.42495
Jupiter	137P/Shoemaker-Levy 2	4.01952	229.51888	2812.93064
Jupiter	131P/Mueller 2	4.05027	208.10376	2535.82718
Jupiter	P/Jedicke (1995 A1)	4.07511	102.29178	1164.85082
Jupiter	D/Tritton (1978 C2)	4.234	282.865	3467.19321
Jupiter	48P/Johnson	4.31823	104.30891	1189.57815
Jupiter	106P/Schuster	4.41102	212.99144	2599.62197
Jupiter	C/LINEAR (1998 M5)	4.67076	318.4884	3874.26616

TABLE 3—*Continued*

Planet	Comet	$(\Delta/R_l)^a$	θ , degrees	$t(\theta)$, d
Jupiter	P/LONEOS (1999 RO28)	4.84285	141.12541	1654.58458
Jupiter	98P/Takamizawa	4.88896	113.76163	1306.55052
Jupiter	C/Spacewatch (1997 P2)	4.89278	286.39551	3508.49848
Jupiter	73P/Schwassmann-Wachmann 3	4.9158	51.54895	569.03117
Jupiter	P/Helin-Lawrence (1993 K2)	4.99431	76.00361	850.14018
Saturn	C/Catalina (1999 F1)	1.01329	107.67064	3035.62842
Saturn	P/Gehrels (1997 C1)	1.27361	164.60138	4862.59895
Saturn	P/Hermann (1999 D1)	1.35745	252.08077	7703.6729
Saturn	C/LINEAR (1998 Q1)	1.36923	70.34971	1928.09742
Saturn	P/Shoemaker 4 (1994 J3)	2.45512	358.40089	10704.16966
Saturn	P/Montani (1997 G1)	2.89611	185.28975	5549.17747
Saturn	P/Helin (1987 Q3)	3.16833	76.81205	2114.42428
Saturn	63P/Wild 1	3.2706	260.23561	7952.44828
Saturn	140P/Bowell-Skiff	3.30576	231.84877	7070.65503
Saturn	D/Bradfield 1 (1984 A1)	3.57748	262.73062	8027.80737
Saturn	134P/Kowal-Vavrova	3.64885	319.9063	9665.71078
Saturn	C/Spacewatch (1997 BA6)	3.95068	44.94144	1214.46702
Saturn	C/LINEAR (1999 N4)	3.95944	78.38428	2160.08505
Saturn	C/LINEAR (1999 H3)	4.19679	241.0887	7362.46033
Saturn	P/Lagerkvist-Carsenty (1997 T3)	4.90066	120.32592	3428.86878

^aThe first 32 entries (above the line) are for intersections having $\Delta < R_l$. All other encounters are for $\Delta < 5R_l$.







