

A Star Product for Complex Grassmann Manifolds

Joachim Schirmer*

September 12, 1997

FR-THEP 21

Abstract

We explicitly construct a star product for the complex Grassmann manifolds using the method of phase space reduction. Functions on $\mathbb{C}^{(p+q) \times p}$, the space of $(p+q) \times p$ matrices of rank p , invariant under the right action of $Gl(p, \mathbb{C})$ can be regarded as functions on the Grassmann manifold $G_{p,q}(\mathbb{C})$, but do not form a subalgebra whereas functions only invariant under the unitary subgroup $U(p) \subset Gl(p, \mathbb{C})$ do. The idea is to construct a projection from $U(p)$ - onto $Gl(p, \mathbb{C})$ -invariant functions, whose kernel is an ideal. This projection can be used to define a star-algebra on $G_{p,q}(\mathbb{C})$ onto which this projection acts as an algebra-epimorphism.

*e-mail: Joachim.Schirmer@physik.uni-freiburg.de

1 Introduction

The idea of deformation quantization is to modify the multiplication on the algebra of smooth complex valued functions in a noncommutative way, such that in zeroth order the normal multiplication is preserved, the commutator equals in first order the Poisson bracket, the constant function with value one acts again as one in the new algebra and the pointwise complex conjugation remains an antilinear involution. This concept of a quantization, first introduced by Berezin [1], later as a program formulated by F. Bayen, M. Flato, c. Fronsdal, A. Lichnerowicz D. Sternheimer[2], is obviously founded on the existence of this kind of a deformed product. This nontrivial question was settled for symplectic manifolds by M. DeWilde and P.B.A. Lecomte [3], independently by B. Fedosov [4, 5] and by H. Omori, Y. Maeda and A. Yoshioka [6].

Explicit examples of star products for nontrivial phase spaces of physical interest are still rare. Although Grassmann manifolds $G_{p,q}(\mathbb{C})$ may be seen as the reduced phase space of a system of $(p+q) \cdot p$ harmonic oscillators where certain energy and angular momentum sums are fixed, it is not really a physical example. So the aim of this paper is principally to show that phase space reduction may be used to construct star products on more difficult symplectic manifolds. The methods used might be relevant as well to the possible deformation quantization of constrained systems.

C. Moreno [7] has already stated a recursion formula for a star product on nonexceptional Kähler symmetric spaces. M. Cahen, S. Gutt and J. Rawnsley [8] defined a star product on every compact Hermitian symmetric space and proved the convergence for the algebra of representative functions. Regarding $G_{p,q}(\mathbb{C})$ as a coadjoint orbit of $U(p+q)$ Karabegov [9] has given a method of deformation quantization in terms of representative functions which probably could be specialized explicitly in this case. The existence of star products on coadjoint orbits was treated as well by D. Arnal, J. Ludwig and M. Masmoudi [10]. For the special case of the projective space a closed formula for a star product of Wick type was given by M. Bordemann *et al.* [11]. It was constructed as well by phase space reduction and will be generalized in the present paper. Whereas there an equivalence transformation of the Wick product on $\mathbb{C}^{q+1} \setminus \{0\}$ onto a new star product was constructed which could be projected to $\mathbb{C}P^q$ trivially, in the case of Grassmann manifolds $G_{p,q}(\mathbb{C})$ we will directly head for the projection, because a similar equivalence transformation does not exist since the group $Gl(p, \mathbb{C})$ is noncommutative for $p > 1$.

The paper will be organized as follows: In the next section we will show how to construct a projection of functions on $\mathbb{C}^{(p+q) \cdot p}$ onto functions on $G_{p,q}(\mathbb{C})$ whose kernel is an ideal with respect to the Wick star product on $C^\infty(\mathbb{C}^{(p+q) \cdot p})$. Then we will calculate this projection for the well known case of the projective space $\mathbb{C}P^q = G_{1,q}(\mathbb{C})$, thereby verifying the result of Bordemann *et al.* [11] in a very simple way. We use this as a guideline in the fourth section where the formula for the star product on Grassmann manifolds is proved. The arising coefficients are treated in the last section, where we will state some open questions as well. Throughout this paper we use Einstein's summation convention.

2 The decomposition of the space of invariant functions

Our aim is to construct a star product on Grassmann manifolds $G_{p,q}(\mathbb{C})$ by making use of the Wick product on $\mathbb{C}^{(p+q)\cdot p^*}$, the space of complex $(p+q) \times p$ matrices of rank p , from which the Grassmann manifolds can be obtained by projection π . Unfortunately the homogeneous functions on $\mathbb{C}^{(p+q)\cdot p^*}$ – functions invariant under the $Gl(p, \mathbb{C})$ -action, which are the bijective image of functions on $G_{p,q}(\mathbb{C})$ by the pullback π^* – do not form a subalgebra. Yet the set of invariant functions – we reserve this term for functions invariant under the unitary subgroup $U(p)$ of $Gl(p, \mathbb{C})$ – is a subalgebra which we will denote by $(C_0^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]], \tilde{*})$. So the Wick star product of two homogeneous functions will only be an invariant function and we are forced to search for a certain "projection" $P := C_0^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]] \rightarrow C_{hom}^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]]$, which satisfies $P(P(f \tilde{*} g) \tilde{*} h) = P(f \tilde{*} P(g \tilde{*} h))$ for any three homogeneous functions f, g, h . Then we can obviously equip $C_{hom}^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]]$ with an algebra structure by defining $f * g := P(f \tilde{*} g)$. For the unity function we would like the equation $f = f * 1 = P(f \tilde{*} 1) = Pf$ to hold, thus $P|_{C_{hom}^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]]} = id|_{C_{hom}^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]]}$ and $P^2 = P$, so the name projection for P is justified. We can split the algebra $C_0^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]] = PC_0^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]] \oplus \ker P$ and notice that if P satisfies the above rules and is thus an algebra epimorphism of $(C_0^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]], \tilde{*})$ onto $(C_{hom}^\infty(\mathbb{C}^{(p+q)\cdot p^*})[[\lambda]], *)$, obviously $\ker P$ must be a $\tilde{*}$ -ideal $\mathcal{J}_{\tilde{*}}$. This condition is in fact an equivalent formulation, the projection P is an algebra epimorphism if and only if $\ker P$ forms a $\tilde{*}$ -ideal $\mathcal{J}_{\tilde{*}}$. The decomposition of the space of invariant functions shall now be constructed.

The complex Grassmann manifold $G_{p,q}(\mathbb{C})$ is the space of equivalence classes of the complex $(p+q) \times q$ matrices of rank p where the equivalence class is defined as follows:

$$\begin{aligned} z_1 \sim z_2 & \iff \exists g \in Gl(p, \mathbb{C}) \quad z_2 = z_1 g \quad z_1, z_2 \in \mathbb{C}^{(p+q)\cdot p^*} \\ \pi : \mathbb{C}^{(p+q)\cdot p^*} & \longrightarrow G_{p,q}(\mathbb{C}), \quad z \longmapsto z/\sim \end{aligned} \quad (1)$$

The Grassmann manifold $G_{1,q}$ is the projective space $\mathbb{C}P^q$. Grassmann manifolds can also be regarded as the reduced phase space associated to the right action Φ of the unitary group $U(p)$:

$$\Phi : \mathbb{C}^{(p+q)\cdot p^*} \times U(p) \longrightarrow \mathbb{C}^{(p+q)\cdot p^*}, \quad (z, U) \longmapsto \Phi_U z = zU$$

This action is symplectic with respect to the Kähler symplectic form

$$\omega = \frac{i}{2} \text{tr} dz \wedge dz^\dagger = \frac{i}{2} \partial \bar{\partial} \text{tr} z^\dagger z = \frac{i}{2} \partial \bar{\partial} \text{tr} x \quad x := z^\dagger z \quad (2)$$

and there is an Ad^* -equivariant momentum mapping

$$J : \mathbb{C}^{(p+q)\cdot p^*} \longrightarrow u(p)^*, \quad J = \frac{i}{2} z^\dagger z = \frac{i}{2} x \quad (3)$$

where $u(p)^*$ denotes the dual of the space of antihermitian matrices. Since for any positive definite hermitian $\mu \in H^+(p)$ there exists a representative of z/\sim on $J^{-1}(\frac{i}{2}\mu)$, namely $z(z^\dagger z)^{-\frac{1}{2}} \mu^{\frac{1}{2}}$, the Grassmann manifold $G_{p,q}$ is the quotient $J^{-1}(\frac{i}{2}\mu)/\mu^{-\frac{1}{2}} U \mu^{\frac{1}{2}}$, and this is the reduced phase space if and only if μ is a positive scalar multiple of the unit matrix, which is

always the case for $p = 1$. So we choose in the following $\mu \in \mathbb{R}^+$ and elucidate the situation by the usual phase space reduction diagramm [12]:

$$\begin{array}{ccc}
& \mathbb{C}^{(p+q) \cdot p * } & \\
& \nearrow i & \downarrow \pi \\
J^{-1}(\frac{i}{2}\mu) & & G_{p,q}(\mathbb{C}) \\
& \searrow \bar{\rho} & \\
& \searrow \pi_\mu &
\end{array}$$

Additionally to the general case there exist the direct projections π and $\bar{\rho}$. Identifying $\mathbb{C}^{(p+q) \cdot p * }$ with $J^{-1}(\frac{i}{2}\mu) \times H^+$ by the global diffeomorphism

$$z \longmapsto (\zeta, x) = (z(z^\dagger z)^{-\frac{1}{2}}\mu^{\frac{1}{2}}, z^\dagger z) \quad (4)$$

the mappings i and $\bar{\rho}$ adopt the explicit form

$$i(\zeta) = (\zeta, \mu) \quad \bar{\rho}(\zeta, x) = \bar{\rho}(z) = \zeta = z(z^\dagger z)^{-\frac{1}{2}}\mu^{\frac{1}{2}}, \quad (5)$$

and an invariant function $f \in C_0^\infty(\mathbb{C}^{(p+q) \cdot p * })$ satisfies

$$f(z) = f(\zeta, x) = f(zU) = f(\zeta U, U^\dagger x U). \quad (6)$$

Looking for an appropriate choice of the star ideal and the associated projection we consider the trivial case of the pointwise product as a guideline. There is obviously a 1:1 mapping between invariant functions on the orbit $J^{-1}(\frac{i}{2}\mu)$ and functions on $G_{p,q}(\mathbb{C})$, i.e. π_μ^* is invertible for invariant functions. All invariant functions on the orbit may be obtained by restriction of invariant functions on $\mathbb{C}^{(p+q) \cdot p * }$ and thus there is a simple projection P .

$$P : C_0^\infty(\mathbb{C}^{(p+q) \cdot p * }) \longrightarrow C_{hom}^\infty(\mathbb{C}^{(p+q) \cdot p * }), \quad P := \pi^* \circ \pi_\mu^{*-1} \circ i^* = \bar{\rho}^* \circ i^* \quad (7)$$

We can easily decompose an invariant function into its homogeneous and its ideal part, thereby describing the ideal:

$$[(1 - P)F](z) = F(\zeta, x) - F(\zeta, \mu) = \left\langle \int_0^1 dt \frac{\partial F}{\partial x}(\zeta, tx + (1 - t)\mu), x - \mu \right\rangle$$

So $\ker P = (1 - P)C_0^\infty(\mathbb{C}^{(p+q) \cdot p * })$ is generated by $x - \mu$:

$$\ker P = \mathcal{J} = \langle x - \mu \rangle = \{F \mid F = \langle x - \mu, G \rangle, \quad G \in C_{eq}^\infty(\mathbb{C}^{(p+q) \cdot p * }, u(p))\}$$

Note that in our case of invariant functions we can restrict the domain of functions paired with $x - \mu$ onto Ad-equivariant functions since for

$$H : C_0^\infty(\mathbb{C}^{(p+q) \cdot p * }) \longrightarrow C_{eq}^\infty(\mathbb{C}^{(p+q) \cdot p * }), \quad H(F)(\zeta, x) := \int_0^1 dt \frac{\partial F}{\partial x}(\zeta, tx + (1 - t)\mu) \quad (8)$$

holds

$$H(F \circ \Phi_U) = \text{Ad}_{U^{-1}} \circ H(F) \circ \Phi_U = H(F). \quad (9)$$

The decomposition

$$F = P.F + (1 - P).F = \bar{\rho}^* i^* F + \langle H(F), x - \mu \rangle \quad (10)$$

is unique whereas the function H is not, unless $p = 1$, because we can add to $H(F)$ any function orthogonal to $x - \mu$. We have chosen H here to be the integral along the straight line between (ζ, x) and (ζ, μ) , but any other curve connecting these points will do as well. For the solution a convenient choice of $H(F)$ will be important as well as the proof that the decomposition is unique, such that the result is independent of the choice of $H(F)$.

Analogously we would like to define a star ideal \mathcal{J}_* generated by $x - \mu$. First of all we have of course to extend our considerations to power series in a formal parameter λ with coefficients in the respective function spaces. The ideal

$$\mathcal{J}_* := \langle x - \mu \rangle_{\tilde{*}} = \{F \mid \langle G \tilde{*} x - \mu \rangle = G^i_j \tilde{*} (x - \mu)^j_i, \quad G \in C_{eq}^\infty(\mathbb{C}^{(p+q) \cdot p}, u(p))[[\lambda]]\}$$

may be defined independently of a global basis for $\mathbb{C}^{(p+q) \cdot p}$ and is indeed a twosided ideal because the Wick star product is strongly $U(p)$ -invariant

$$f \tilde{*} J - J \tilde{*} f = \frac{i\lambda}{2} \{f, J\} \quad \forall f \in C^\infty(\mathbb{C}^{(p+q) \cdot p}), \quad (11)$$

while $U(p)$ is compact – note that G is only equivariant. Now we can formulate the important

Lemma 2.1 *There exists a unique decomposition of invariant functions*

$$C_0^\infty(\mathbb{C}^{(p+q) \cdot p})[[\lambda]] = C_{hom}^\infty(\mathbb{C}^{(p+q) \cdot p})[[\lambda]] \oplus \mathcal{J}_*$$

into a homogeneous and a $\tilde{*}$ -ideal part, where the ideal is spanned by $x - \mu$.

As an immediate consequence of the proposition and the remarks in the introduction we obtain the following

Corollary 2.1 *There exists a star product on $G_{p,q}(\mathbb{C})$ and an algebra epimorphism*

$$\pi^{*-1} \circ P : (C_0^\infty(\mathbb{C}^{(p+q) \cdot p})[[\lambda]], \tilde{*}) \longrightarrow (C^\infty(G_{p,q}(\mathbb{C}))[[\lambda]], *) . \quad \square$$

Proof: We start with the point multiplicative decomposition and start deforming the usual product in the following way:

$$F = \bar{\rho}^* i^* F + \langle H(F), x - \mu \rangle = \bar{\rho}^* i^* F + \langle H(F) \tilde{*} x - \mu \rangle + \lambda \Delta(F) \quad (12)$$

$$\Delta(F) := \frac{1}{\lambda} (\langle H(F) \tilde{*} x - \mu \rangle - \langle H(F), x - \mu \rangle) \quad (13)$$

$\Delta(F)$ is a power series in λ since the coefficient of λ^{-1} vanishes and we can proceed decomposing $\Delta(F)$. Using induction, keeping in mind the linearity of $\bar{\rho}^*$, i^* , H and Δ yields:

$$F = \bar{\rho}^* i^* \left(\frac{1}{1 + \lambda \Delta} F \right) + \left\langle H \left(\frac{1}{1 + \lambda \Delta} F \right) \tilde{*} x - \mu \right\rangle \quad (14)$$

So there is a decomposition, but since H is not uniquely defined, Δ is not either, so uniqueness might be lost during the deformation process. To insure uniqueness assume a function F in the $\tilde{*}$ -ideal \mathcal{J}_* is simultaneously homogeneous, i.e.

$$\bar{\rho}^* \varphi = F = \langle G \tilde{*}; x - \mu \rangle, \quad G \in C_{eq}^\infty(\mathbb{C}^{(p+q) \cdot p}, u(p)) [[\lambda]] \quad \varphi \in C_0^\infty\left(J^{-1}\left(\frac{i}{2}\mu\right)\right) [[\lambda]]$$

It has to be shown that F vanishes identically, thus both sides of the above equation. We can make use of the fact, that the series for the Wick product with x breaks off after the first term:

$$\langle G \tilde{*}; x \rangle = \langle G, x \rangle + \lambda \tilde{M}_1 \langle G, x \rangle, \quad \tilde{M}_1 \langle G, x \rangle := z_i^A \frac{\partial}{\partial z_j^A} G^i_j \quad G \in C_{eq}^\infty(\mathbb{C}^{(p+q) \cdot p}, u(p)) \quad (15)$$

Let $\varphi_\nu \in C_0^\infty(J^{-1}(\frac{i}{2}\mu))$ and $G_{(\nu)} \in C_{eq}^\infty(\mathbb{C}^{(p+q) \cdot p}, u(p))$ be the coefficients of φ and G respectively:

$$\bar{\rho}^* \varphi = \sum_{\nu=0}^{\infty} \lambda^\nu \bar{\rho}^* \varphi_\nu = \sum_{\nu=0}^{\infty} \lambda^\nu \left(\langle G_{(\nu)}, x - \mu \rangle + \tilde{M}_1 \langle G_{(\nu-1)}, x \rangle \right) \quad G_{(-1)} \equiv 0$$

Considering all orders separately we obtain after restriction to $x = \mu$

$$\varphi_\nu = i^* \tilde{M}_1 \langle G_{(\nu-1)}, x - \mu \rangle \quad \varphi_0 = 0. \quad (16)$$

The strategy is the following: Assume $\varphi_\nu = 0$ for $\nu < n$. Consequently it holds

$$\tilde{M}_1 \langle G_{(\nu-1)}, x - \mu \rangle = -\langle G_{(\nu)}, x - \mu \rangle \quad \text{for } \nu < n. \quad (17)$$

Express $\varphi_n = i^* \tilde{M}_1 \langle G_{(n-1)}, x \rangle$ by derivatives of $\langle G_{(n-1)}, x - \mu \rangle$ and use the $n - 1$ equations (17) successively in order to write φ_n as derivatives of $\langle G_{(0)}, x - \mu \rangle$ which vanishes identically. Then $\varphi_n = 0$ and by induction $\varphi_\nu = 0$ for all ν what completes the proof.

Obviously we have to consider derivatives at $J^{-1}(\frac{i}{2}\mu)$ transversally to this submanifold. Thus in what follows we set $z = \zeta y$ and we investigate derivatives with respect to y at the point $y = 1$. In a tedious, but simple calculation one verifies the following formulas.

$$\left. \frac{\partial^r}{\partial y_{j_1}^{i_1} \dots \partial y_{j_r}^{i_r}} \right|_{y=1} \left(\langle G_{(\nu)}, x - \mu \rangle (\zeta y) \right) = \mu \sum_{s=1}^r \frac{\partial^{r-1} G_{(\nu)}^{j_s}_{i_s}}{\partial z_{j_1}^{A_1} \dots \partial z_{j_r}^{A_r}} (\zeta) \zeta_{i_1}^{A_1} \dots \zeta_{i_r}^{A_r} \quad (18)$$

$$\begin{aligned} \left. \frac{\partial^r}{\partial y_{j_1}^{i_1} \dots \partial y_{j_r}^{i_r}} \right|_{y=1} \left(\tilde{M}_1 \langle G_{(\nu)}, x - \mu \rangle (\zeta y) \right) &= \delta_{j_{r+2}}^{i_{r+1}} \delta_{j_{r+1}}^{i_{r+2}} \frac{\partial^{r+1} G_{(\nu)}^{j_{r+2}}_{i_{r+2}}}{\partial z_{j_1}^{A_1} \dots \partial z_{j_{r+1}}^{A_{r+1}}} (\zeta) \zeta_{i_1}^{A_1} \dots \zeta_{i_{r+1}}^{A_{r+1}} \\ &+ \sum_{s=1}^r \delta_{j_{r+1}}^{i_{r+1}} \frac{\partial^r G_{(\nu)}^{j_s}_{i_{r+1}}}{\partial z_{j_1}^{A_1} \dots \partial z_{j_{r+1}}^{A_{r+1}} \dots \partial z_{j_s}^{A_r}} (\zeta) \zeta_{i_1}^{A_1} \dots \zeta_{i_r}^{A_r} \quad (19) \end{aligned}$$

Before proceeding with the proof let us summarize some notation for conjugacy classes of the symmetric group. By the r -tuple $\alpha = (\alpha_1, \dots, \alpha_r)$ we denote the conjugacy class in S_r having α_i i -cycles in the cycle representation, by $Con(S_r)$ the set of all conjugacy classes

Obviously it holds $\sum i\alpha_i = r$ and we define $|\alpha| = \alpha_1 + \dots + \alpha_r$ as usual. The number of elements in the conjugacy class α is

$$h_\alpha = \frac{r!}{1^{\alpha_1}\alpha_1! \cdot \dots \cdot r^{\alpha_r}\alpha_r!}. \quad (20)$$

Since $i - 1$ transpositions are necessary to compose an i -cycle from 1-cycles (resp. to decompose it into 1-cycles) the lowest number of transpositions, denoted by $\#$, that is needed to compose a certain permutation $\sigma \in \alpha$ from the identity (or vice versa to decompose it into the identity) is only dependent on the conjugacy class α and it holds

$$\#\alpha = \#[\sigma] = \sum_{i=1}^r (i-1)\alpha_i = r - |\alpha| \quad (21)$$

We abbreviate by $D := \left(\frac{\partial}{\partial y^i_j}\right)_{i,j=1\dots p}$ and set $D_k = \text{tr}D^k := \frac{\partial^r}{\partial y^{i_1}_{i_2} \partial y^{i_2}_{i_3} \dots \partial y^{i_k}_{i_1}}$. Now assume $\varphi_\nu \equiv 0$ for $\nu < n$ thus also equations (17). Then we find successively

$$\varphi_n = -\frac{1}{(-\mu)^k} \sum_{r=k+1}^{2k} \frac{1}{r!} \sum_{\substack{\alpha \in \text{Con}(S_r) \\ \alpha_1=0 \quad \#\alpha=k}} h_\alpha D_2^{\alpha_2} \dots D_r^{\alpha_r} \langle G_{(n-k)}, x - \mu \rangle. \quad (22)$$

This formula can be proved by induction using the above formulas. Using it for $k = n$ shows that φ_n vanishes identically and the proof is completed. \square

A final lemma will be useful in the next section:

Lemma 2.2 *Let $F = PF + \langle G \tilde{*}; x - \mu \rangle$ be the unique decomposition of the invariant function F and let f be homogeneous $Pf = f$. Then $fF = fPF + \langle fG \tilde{*}; x - \mu \rangle$ is the decomposition of the product fF .*

Proof: For a homogeneous function f holds $z^A_i \frac{\partial}{\partial z^A_j} f = 0 \quad i, j = 1 \dots p$. Hence we have $\langle fG \tilde{*}; x - \mu \rangle = f \langle G \tilde{*}; x - \mu \rangle$. \square

3 The solution for the projective space $\mathbb{C}P^q$

The first step towards the construction of a star product on the quotient manifold for the projective space as well as for the general Grassmann manifolds consists in "homogenizing" the expression for the Wick product of two homogeneous functions. The summands in the series are written as products of homogeneous differential operators – differential operators that assign two homogeneous functions another homogeneous function and that are thus well defined on the quotient manifold – and powers of a certain nonhomogeneous function. According to the lemma in the last paragraph it suffices to calculate the projection for this certain function and its powers. In case of the projective space $\mathbb{C}P^q$ the homogeneization is simple and was also used in [11] to find an explicit expression for the star product on $\mathbb{C}P^q$.

$$\begin{aligned} F \tilde{*} G &= \sum_{r=0}^{\infty} \frac{\lambda^r}{r!} x^{-r} \underbrace{\left(x^r \frac{\partial^r F}{\partial z^{A_1} \dots \partial z^{A_r}} \cdot \frac{\partial^r G}{\partial z^{\dagger A_1} \dots z^{\dagger A_r}} \right)} \\ &= \sum_{r=0}^{\infty} \frac{\lambda^r}{r!} x^{-r} M_r(F, G) \end{aligned} \quad (23)$$

Obviously $x \frac{\partial}{\partial z^A} \otimes \frac{\partial}{\partial z^\dagger_A}$ is a homogeneous differential operator, assigning two homogeneous functions the tensor product of homogeneous functions, so its power (followed by multiplication) M_r are as well homogeneous. Now a star product for homogenous functions $F, G \in C_{hom}^\infty(\mathbb{C}^{q+1} \setminus \{0\})$ is easily constructed

$$F * G = P(F \tilde{*} G) = \sum_{r=0}^{\infty} P(x^{-r} M_r(F, G)) = \sum_{r=0}^{\infty} P(x^{-r}) M_r(F, G) \quad , \quad (24)$$

where the last equation sign holds because of lemma 2.2. Following the program described in section 2 we obtain

$$\langle H(x^{-r}), x - \mu \rangle = x^{-r} - \bar{\rho}^* i^* x^{-r} = x^{-r} - \mu^{-r} = -(x - \mu) \sum_{\nu=1}^r x^{-\nu} \mu^{\nu-r-1}$$

In this case $H(F)$ is a uniquely determined invariant function.

$$\Delta(x^{-r}) = z^A \frac{\partial}{\partial z^A} H(x^{-r}) = x \frac{\partial}{\partial x} H(x^{-r}) = \sum_{\nu=1}^r \nu x^{-\nu} \mu^{\nu-r-1}$$

Applying the general recursion formula

$$P(F) = \bar{\rho}^* i^* F - \lambda P(\Delta F) \quad (25)$$

to the cases $F = x^{-r}$ and $F = \mu^{-1} x^{-r+1}$ as well as to $F = x^{-1}$ yields a recursion formula and its beginning:

$$P(x^{-r}) = \frac{1}{\mu + \lambda r} P(x^{-(r-1)}) \quad P(x^{-1}) = \frac{1}{\mu + \lambda}$$

This proves the simple

Lemma 3.1
$$P(x^{-r}) = \prod_{s=1}^r \frac{1}{\mu + \lambda s} = \lambda^{-r} \frac{1}{[c]_r} \quad ,$$

where $[y]_r := y(y+1) \dots (y+r-1)$, $[y]_0 := 1$ and $c := \mu \lambda^{-1} + 1$. □

In the same way as described here one could calculate $P(x^r) = \prod_{s=0}^{r-1} (\mu - \lambda s)$.

Corollary 3.1 *There exists a star product on $G_{1,q}(\mathbb{C}) = \mathbb{C}P^q$ defined by $(f, g \in C^\infty(\mathbb{C}P^q))$*

$$\pi^*(f * g) := P(\pi^* f \tilde{*} \pi^* g) = \sum_{r=0}^{\infty} \frac{1}{r! [c]_r} M_r(\pi^* f, \pi^* g) \quad , \quad (26)$$

where the homogeneous operators M_r are defined by

$$M_r(F, G) := x^r \frac{\partial^r F}{\partial z^{A_1} \dots \partial z^{A_r}} \cdot \frac{\partial^r G}{\partial z^\dagger_{A_1} \dots \partial z^\dagger_{A_r}} = \mu^r \bar{\rho}^* i^* \left(\frac{\partial^r F}{\partial z^{A_1} \dots \partial z^{A_r}} \cdot \frac{\partial^r G}{\partial z^\dagger_{A_1} \dots \partial z^\dagger_{A_r}} \right) \quad \square \quad (27)$$

At this point it seems to be only a matter of curiosity that for this star product not only the limit $\lambda \rightarrow 0$, but also the opposite one, $\lambda \rightarrow \infty$ is well defined as a formal power series. And at a glance the space of functions for which this product converges is nonempty. This strange possibility of a "strong quantum limit" will be preserved in the Grassmann case.

4 The projection for Grassmann manifolds

We have to find a transferred version of the homogeneization procedure (23), since x is no longer a scalar function. First note that according to the usual properties differentiation lowers the degree of homogeneity by one, so for a homogeneous function $F \in C_{hom}^\infty(\mathbb{C}^{(p+q) \cdot p} *)$ which is more precisely a homogeneous of degree 0 holds:

$$F(zg) = F(z) \quad \forall g \in Gl(p, \mathbb{C}) \quad \Longrightarrow \quad g^i_j \frac{\partial F}{\partial z^A_j}(zg) = \frac{\partial F}{\partial z^A_i}(z) \quad \begin{array}{l} i = 1 \dots p \\ A = 1 \dots p + q \end{array}$$

Thus the expression $z^B_i \frac{\partial F}{\partial z^A_i}$ is again homogeneous and this suggests to rewrite the expression for the Wick product for the case of homogeneous functions $F, G \in C_{hom}^\infty(\mathbb{C}^{(p+q) \cdot p} *)$ in the following way:

$$\begin{aligned} F \tilde{*} G &= m \circ \exp \left(\lambda \delta^A_B \delta^j_i \frac{\partial}{\partial z^A_i} \otimes \frac{\partial}{\partial z^{\dagger j}_B} \right) (F \otimes G) \\ &= m \circ \exp \left(\lambda S^D_C z^C_i \frac{\partial}{\partial z^A_i} \otimes \frac{\partial}{\partial z^{\dagger j}_A} z^{\dagger j}_D \right) (F \otimes G) = \sum_{r=0}^{\infty} \frac{\lambda^r}{r!} \left(S^{\otimes r}, M_r(F, G) \right) \end{aligned} \quad (28)$$

where

$$\begin{aligned} S &= z x^{-2} z^\dagger, \\ M_r(F, G)^{C_1 \dots C_r}_{D_1 \dots D_r} &= z^{C_1}_{i_1} \dots z^{C_r}_{i_r} \frac{\partial^r F}{\partial z^{A_1}_{i_1} \dots \partial z^{A_r}_{i_r}} \frac{\partial^r G}{\partial z^{\dagger j_1}_{A_1} \dots \partial z^{\dagger j_r}_{A_r}} z^{\dagger j_1}_{D_1} \dots z^{\dagger j_r}_{D_r}, \\ m(F \otimes G) &= F \cdot G, \end{aligned}$$

and (\cdot, \cdot) is the extension of the usual hermitian product $(A, B) = \text{tr} A^\dagger B$ to tensor products. The matrix S is obviously hermitian and in contrast to the momentum mapping J , which is only equivariant with respect to the $U(p)$ -action, even $U(p)$ -invariant. The homogeneous differential operators take values in the r -fold tensor product of the defining representation of $U(p+q)$.

Formally the construction of a star product for homogeneous functions on $\mathbb{C}^{(p+q) \cdot p} *$ is completely analogous to the $\mathbb{C}P^q$ -case of the last section

$$F * G := P(F \tilde{*} G) = \sum_{r=0}^{\infty} \frac{\lambda^r}{r!} P \left(S^{\otimes r}, M_r(F, G) \right) = \sum_{r=0}^{\infty} \frac{\lambda^r}{r!} \left(P(S^{\otimes r}), M_r(F, G) \right), \quad (29)$$

where again the last equation sign is justified by lemma (5). Yet the evaluation of the projection $P(S^{\otimes r})$ shows a new aspect for which we introduce the notion of the action of the symmetric group on the tensor product space $(\mathbb{C}^s)^{\otimes r}$:

Definition 4.1 For $\sigma \in S_r$ let $\rho(\sigma) : (\mathbb{C}^s)^{\otimes r} \longrightarrow (\mathbb{C}^s)^{\otimes r}$ be the representation defined by linear extension of the prescription $\rho(\sigma)(e_{A_1} \otimes \dots \otimes e_{A_r}) = e_{A_{\sigma(1)}} \otimes \dots \otimes e_{A_{\sigma(r)}} \quad 1 \leq A_i \leq s$, where $\{e_A\}_{A=1 \dots s}$ is a basis of \mathbb{C}^s . This representation can linearly be extended to the group algebra. As well let for $S \in Gl(s, \mathbb{C})$ be $D(S) : (\mathbb{C}^s)^{\otimes r} \longrightarrow (\mathbb{C}^s)^{\otimes r}$ the representation defined by $D(S) := S^{\otimes r}$, which in a basis reads $D(S)(e_{A_1} \otimes \dots \otimes e_{A_r}) := e_{B_1} \otimes \dots \otimes e_{B_r} S^{B_1}_{A_1} \dots S^{B_r}_{A_r}$.

Obviously the two representations commute, that means $\rho(\sigma) \circ D(S) = D(S) \circ \rho(\sigma)$ for all $\sigma \in S_r$ and $S \in Gl(s, \mathbb{C})$. Linear extension of D to a representation of the endomorphism algebra $End(\mathbb{C}^s)$ leads to the subalgebra of all mappings that commute with permutations. We will denote this subalgebra as $\mathbb{C}[D(Gl(s, \mathbb{C}))]$.

Using the notation introduced in lemma 2.1 concerning the conjugacy classes of the symmetric group we can formulate the essential

Lemma 4.1 *For the projection $P(S^{\otimes r})$ the following recursion relation holds:*

$$\lambda^r \rho \left(\sum_{\alpha \in Con(S_r)} c^{|\alpha|} k_\alpha \right) P(S^{\otimes r}) = T^{\otimes r} , \quad (30)$$

where $T := \mu \bar{\rho}^* i^* S = \mu^{-1} \zeta^\dagger \zeta$, $c := \lambda^{-1} \mu + p$ and $k_\alpha := \sum_{\sigma \in \alpha} \sigma$ is defined to be the sum over all permutations of the conjugation class α in the group algebra $\mathbb{C}[S_r]$.

Proof: The formula will be proved by induction. For any constant hermitian $p + q$ matrix Φ holds:

$$\left\langle H \left((S, \Phi) \right), x - \mu \right\rangle = -\mu^{-1} \langle zx^{-1}(x - \mu)x^{-1}z^\dagger, \Phi \rangle = -\mu^{-1} \langle x^{-1}z^\dagger \Phi zx^{-1}, x - \mu \rangle$$

and we can choose

$$H \left((S, \Phi) \right) = -\mu^{-1} z^\dagger \Phi zx^{-1} . \quad (31)$$

The calculation of $\Delta(S, \Phi)$ is simplified by the fact that $z^A_i \frac{\partial}{\partial z^A_j}$ is the fundamental vector field for the holomorphic $Gl(p, \mathbb{C})$ -action, i.e. $((z, z^\dagger), g) \rightarrow (zg, z^\dagger)$ under which the term zx^{-1} is invariant.

$$\Delta \left((S, \Phi) \right) = z^A_i \frac{\partial}{\partial z^A_j} H^i_j(S, \Phi) = \mu^{-1} p(S, \Phi) \quad \Longrightarrow \quad \Delta(S) = \mu^{-1} pS$$

Using the recursion formula (25) one finds $\lambda c P(S) = T$, which proves the start of the induction. Now assume the formula is proved for $i < r$. A possible choice for H is

$$H(S^{\otimes r}) = S^{\otimes r-1} \otimes H(S) + H(S^{\otimes r-1}) \otimes \mu^{-1} T$$

and it follows

$$\begin{aligned} \Delta(S^{\otimes r}) &= S^{\otimes r-1} \otimes \Delta(S) + \left(z^A_i \frac{\partial}{\partial z^A_j} S^{\otimes r-1} \right) \otimes H^i_j(S) + \Delta(S^{\otimes r-1}) \otimes \mu^{-1} T \\ &= \mu^{-1} p S^{\otimes r} + \sum_{s=1}^{r-1} \rho(\tau_{sr}) S^{\otimes r} + \mu^{-1} \Delta(S^{\otimes r-1}) \otimes T , \end{aligned}$$

where τ_{sr} is the transposition of s and r . Comparing the recursion formula (25) analogously to the proof of lemma (3.1) for $P(S^{\otimes r})$ and $\mu^{-1} P(S^{\otimes r-1}) \otimes T$ taking into consideration that P and ρ commute, yields:

$$\left(1 + \lambda \mu^{-1} p + \lambda \mu^{-1} \sum_{s=1}^{r-1} \rho(\tau_{ir}) \right) P(S^{\otimes r}) = \mu^{-1} P(S^{\otimes r-1}) \otimes T$$

Using the induction hypothesis and the relation (21) leads to

$$\lambda^r \left(c + \sum_{s=1}^{r-1} \rho(\tau_{sr}) \right) \left(\sum_{\substack{\sigma \in S_{r-1} \subset S_r \\ \sigma(r)=r}} c^{r-1-\#\sigma} \rho(\sigma) \right) P(S^{\otimes r}) = \lambda^r \sum_{\sigma \in S_r} c^{r-\#\sigma} \rho(\sigma) P(S^{\otimes r}) = T^{\otimes r},$$

which proves the lemma. \square

Remark: In the same way one can show that for the positive powers holds

$$P \left(z z^\dagger{}^{\otimes r} \right) = \rho \left(\sum_{\alpha \in \text{Con}(S_r)} (-\lambda \mu^{-1})^{\#\alpha} k_\alpha \right) \left(\zeta \zeta^\dagger{}^{\otimes r} \right). \quad (32)$$

Since $\{k_\alpha\}_{\alpha \in S_r}$ is a basis of the center of the group algebra $\mathbb{C}[S_r]$, $\sum_{\alpha \in \text{Con}(S_r)} c^{|\alpha|} k_\alpha$ is an element of the center which will be invertible for generic c . For the inverse element we can write

$$\left(\sum_{\alpha} c^{|\alpha|} k_\alpha \right)^{-1} =: \sum_{\alpha} s_\alpha(c) k_\alpha, \quad (33)$$

where s_α are rational functions which are to be determined. First we formulate a trivial corollary of lemma 4.1.

Corollary 4.1 *There exists a star product on $G_{p,q}(\mathbb{C})$ defined by $(f, g \in C^\infty(G_{p,q}(\mathbb{C}))[[\lambda]])$*

$$\pi^*(f * g) = P(\pi^* f \tilde{*} \pi^* g) = \sum_{r=0}^{\infty} \frac{\mu^r}{r!} \bar{\rho}^* i^* \left\langle \rho \left(\sum_{\alpha \in \text{Con}(S_r)} s_\alpha(c) k_\alpha \right) \frac{\partial^r \pi^* f}{\partial z^{A_1} \dots \partial z^{A_r}}, \frac{\partial^r \pi^* g}{\partial z^\dagger{}_{A_1} \dots \partial z^\dagger{}_{A_r}} \right\rangle, \quad (34)$$

where the rational functions s_α are defined by (33) and $c = \lambda^{-1} \mu + p$. Here we regard the derivatives $\frac{\partial^r F}{\partial z_{i_1}^{A_1} \dots \partial z_{i_r}^{A_r}}$ as (i_1, \dots, i_r) -component of a tensor in $(\mathbb{C}^p)^{\otimes r}$ and $\langle \cdot, \cdot \rangle$ denotes the hermitian inner product of $(\mathbb{C}^p)^{\otimes r}$. \square

A comparison with corollary 3.1 yields as a condition for s_α :

$$\sum_{\sigma \in S_r} s_{[\sigma]}(c) = \sum_{\alpha \in \text{Con}(S_r)} h_\alpha s_\alpha(c) = \frac{1}{[c]_r}. \quad (35)$$

5 The irreducible differential operators and their coefficients

This section is devoted to the determination of the rational functions s_α defined by (33) that arise in the star product on Grassmann manifolds (34). According to Wedderburn's theorem the group algebra $\mathbb{C}[G]$ of a finite group G decomposes directly into simple endomorphism rings $\text{End}(\mathbb{C}^{n_a})$, $a = 1 \dots f$ of dimension n_a^2 such that $\sum n_a^2$ equals the order $|G|$ of the group. Each simple ring $\text{End}(\mathbb{C}^{n_a})$ may be seen as an isotypical submodule of the regular representation of the group G on its group algebra $\mathbb{C}[G]$ obtained by left multiplication. These isotypical submodules $\text{End}(\mathbb{C}^{n_a})$ are associated to a certain irreducible representation

ρ^a , which is therein contained with multiplicity n_a . To sum up, all irreducible representations ρ^a of a finite group G are contained in the group algebra and their multiplicity equals their dimension n_a . The number f of inequivalent irreducible representations equals the number of conjugacy classes, which is in case of the symmetric group S_r the number of partitions of r . So the center of the group ring $\mathbb{C}[G]$ has two natural bases, the conjugation classes k_α and the units e_a of the simple rings $End(\mathbb{C}^{n_a})$ into which the group algebra decomposes. Between both there is a simple base transformation in terms of characters [13], which for the case of the symmetric group S_r reads

$$e_a = \frac{n_a}{r!} \sum_{\alpha=1}^f \chi_\alpha^a k_\alpha \quad k_\alpha = h_\alpha \sum_{a=1}^f \frac{\chi_\alpha^a}{n_a} e_a \quad (36)$$

Here χ_α^a is the character of the irreducible representation ρ^a evaluated at an element of the conjugacy class α . Note that for the symmetric group a group element and its inverse are in the same class, such that the character is real. Of course the basis $\{e_a\}_{a=1\dots f}$ is more appropriate for considering the inverse (33):

$$\left(\sum_{\alpha} c^{|\alpha|} k_\alpha \right)^{-1} = \sum_a \frac{n_a}{\sum_{\alpha} h_\alpha \chi_\alpha^a c^{|\alpha|}} e_a =: \sum_a \frac{1}{t_a(c)} e_a \quad t_a(c) := \frac{1}{n_a} \sum_{\alpha} h_\alpha \chi_\alpha^a c^{|\alpha|} \quad (37)$$

The relation between the rational functions s_α introduced in (33) and the polynomials t_a is

$$s_\alpha(c) = \sum_a \frac{n_a \chi_\alpha^a}{r!} \frac{1}{t_a(c)} \quad (38)$$

As noted above any isotypical submodule $End(\mathbb{C}^{n_a})$ of the group ring or equivalently any irreducible representation ρ^a (up to isomorphism) or any unit e_a is characterised by a partition, which is usually represented by a frame. We quote the dimension formula [13]:

Proposition 5.1 *The dimension $n_{[m]}$ of an irreducible representation $\rho^{[m]}$ of S_r characterised by a frame $[m] = [m_1, \dots, m_k]$ with k rows of lengths $m_1 \geq m_2 \geq \dots, m_k \geq 0$, $|m| := \sum_{i=1}^k m_i = r$, and thus the multiplicity in the group algebra is given by*

$$n_{[m]} = r! \frac{\prod_{i < j} (l_i - l_j)}{l_1! \dots l_k!} \quad l_i = m_i + k - i \quad (39)$$

Note that the dimension is of course invariant if one adds certain rows of length zero, so the right hand side is in fact unchanged under $[m] \mapsto [m'] = [m, 0]$.

Now we consider the decomposition of the r -fold product of the vector space \mathbb{C}^s into symmetry classes. $(\mathbb{C}^s)^{\otimes r}$ carries the representation ρ defined above of the symmetric group S_r and is thus decomposable into irreducible components characterised by a frame. The irreducible components may be contained with multiplicity d_a (possibly 0).

$$(\mathbb{C}^s)^{\otimes r} = d_1 \mathbb{C}^{n_1} \oplus \dots \oplus d_f \mathbb{C}^{n_f} \quad \text{w.r.t. } S_r \quad (40)$$

Since the action D of $Gl(s, \mathbb{C})$ does commute with the action of the symmetric group Schur's lemma states, that an element of $Gl(s, \mathbb{C})$ can only act as a multiple of unity between the

irreducible summands of (40) which has to be zero between \mathbb{C}^{n_i} and \mathbb{C}^{n_j} if $i \neq j$. This leads to another decomposition

$$(\mathbb{C}^s)^{\otimes r} = n_1 \mathbb{C}^{d_1} \oplus \dots \oplus n_f \mathbb{C}^{d_f} \quad \text{w.r.t. } Gl(s, \mathbb{C}), \quad (41)$$

where here \mathbb{C}^{d_a} are irreducible representation spaces of an irreducible representation D^a of $Gl(s, \mathbb{C})$, which arise with multiplicity n_a . As a trivial consequence the isotypical components are identical for both actions:

$$(\mathbb{C}^s)^{\otimes r} = \mathbb{C}^{d_1} \otimes \mathbb{C}^{n_1} \oplus \dots \oplus \mathbb{C}^{d_f} \otimes \mathbb{C}^{n_f} \quad (42)$$

The units e_a of the simple components of the group ring $\mathbb{C}[S_r]$ act as projectors into the isotypical components here as well. A unit e_a associated to a frame whose number of rows exceeds s , acts as an annihilator, $\rho(e_a) = 0$, since the degree of antisymmetrisation is too large, so n_a must be 0 in this case. So the representations D^a , $a = 1 \dots r$ of $Gl(s, \mathbb{C})$ are as well characterised by frames. We quote again the dimension formula [13]:

Proposition 5.2 *For a frame with more than s rows of nonzero length the dimension of the associated representation $D^{[m]}$ of $Gl(s, \mathbb{C})$ is zero. So let $[m]$ be a frame with s rows of length $m_1 \geq \dots \geq m_s \geq 0$, where the lowest rows may be of length zero. Then the dimension $d_{[m]}$ of the associated representation D^a of $Gl(s, \mathbb{C})$ is*

$$d_{[m]} = \frac{\prod_{i < j} (l_i - l_j)}{(s-1)!(s-2)! \dots 1!0!} \quad l_i = m_i + s - i. \quad (43)$$

Finally we recapitulate the considerations leading to Frobenius' formula relating the characters of the respective representations ρ^a and D^a of S_r and $Gl(s, \mathbb{C})$. Consider the composite actions of $A \in Gl(s, \mathbb{C})$ and $\sigma \in S_r$ on $(\mathbb{C}^s)^{\otimes r}$ and assume σ belongs to the conjugacy class α . Using the isotypical decomposition (42) the mapping decomposes as

$$D(A) \circ \rho(\sigma) = \rho(\sigma) \circ D(A) = D^1(A) \otimes \rho^1(\sigma) \oplus \dots \oplus D^f(A) \otimes \rho^f(\sigma)$$

and for its trace one finds on the one hand

$$\text{tr}\left(D(A) \circ \rho(\sigma)\right) = \sum_{a=1}^f \text{tr} D^a(A) \cdot \text{tr} \rho^a(\sigma) = \sum_{a=1}^f \varphi(A) \chi_\alpha^a,$$

where φ^a is the character of the representation D^a , and on the other hand according to definition 4.1

$$\text{tr}\left(D(A) \circ \rho(\sigma)\right) = A_{i_{\sigma(1)}}^{i_1} \cdot \dots \cdot A_{i_{\sigma(r)}}^{i_r} = a_1^{\alpha_1} \cdot \dots \cdot a_r^{\alpha_r},$$

where $a_k := \text{tr} A^k$ is the trace of the powers of A . Finally using the orthogonality relation, which may be read from equations (36) yields Frobenius'

Proposition 5.3 *The relation between the characters of the irreducible representations $D^{[m]}$ of $Gl(s, \mathbb{C})$ and $\rho^{[m]}$ of S_r associated to the same frame $[m] = [m_1, \dots, m_s]$, $|m| = r$ is given by*

$$\varphi^{[m]}(A) = \frac{1}{r!} \sum_{\alpha \in \text{Con}(S_r)} h_\alpha a_1^{\alpha_1} \cdot \dots \cdot a_r^{\alpha_r} \chi_\alpha^{[m]}, \quad (44)$$

where $a_i := \text{tr} A^i$ and h_α is the number of element in the conjugacy class α . \square

Summing up all informations the proof of the following corollary is trivial.

Corollary 5.1 *The polynomial $t_{[m]}$ introduced by (37) associated to the frame $[m]$ is given by*


$$t_{[m]}(c) = [c]_{m_1} [c-1]_{m_2} \cdot \dots \cdot [c-p+1]_{m_p} \quad (45)$$

Proof: It suffices to know the values of the polynomial for all natural values $s \geq p$. Then

$$t_{[m]}(s) = \frac{r!}{n_{[m]}} \varphi^{[m]}(\mathbf{1}_s) = [s]_{m_1} [s-1]_{m_2} \cdot \dots \cdot [1]_{m_s}$$

But the frames under consideration in our case have at most p rows, since otherwise $e_{[m]}$ acts as annihilator on tensors in $(\mathbb{C}^p)^{\otimes r}$. Thus in our case $m_{p+1} = \dots = m_s = 0$ and the formula is proved. \square

This formula has a simple graphical interpretation: Consider an arbitrary frame:



$$\longmapsto \quad t_{[m]}(c) = (c+6)(c+5)(c+4)(c+3)^2(c+2)^3 \times (c+1)^3 c^3 (c-1)^3 (c-2)^2 (c-3)^2 (c-4)$$

For any box lying on the main diagonal $t_{[m]}$ contains a factor c , for any box on the first upper resp. lower diagonal it contains a linear factor $c+1$ resp $c-1$ and so forth. The relation (35) is now recognized as a direct consequence of equation (38) and the orthogonality relation. The star product on the Grassmann manifolds $G_{p,q}(\mathbb{C})$ of corollary 4.1 then takes the form:

Proposition 5.4 *There exists a star product on $G_{p,q}(\mathbb{C})$ such that for $f, g \in C^\infty(G_{p,q}(\mathbb{C}))[[\lambda]]$:*

$$\begin{aligned} \pi^*(f * g) &= P(\pi^* f \tilde{*} \pi^* g) \\ &= \sum_{r=0}^{\infty} \frac{\mu^r}{r!} \sum_{\substack{|m|=r \\ m_{p+1}=0}} \frac{1}{t_{[m]}(c)} \bar{\rho}^* i^* \left\langle \rho(e_{[m]}) \frac{\partial^r \pi^* f}{\partial z^{A_1} \dots \partial z^{A_r}}, \rho(e_{[m]}) \frac{\partial^r \pi^* g}{\partial z^\dagger_{A_1} \dots \partial z^\dagger_{A_r}} \right\rangle \end{aligned} \quad (46)$$

Here $\rho(e_{[m]})$ denotes the projector of $(\mathbb{C}^p)^{\otimes r}$ onto the symmetry class characterised by the frame $[m]$ and $t_{[m]}(c) = [c]_{m_1} [c-1]_{m_2} \cdot \dots \cdot [c-p+1]_{m_p}$. In the sum over the frames only such frames yield a nonzero contribution whose number of rows is not greater than p . \square

Obviously this theorem contains Corollary (3.1) as a special case for $p = 1$. The product remains well defined in the limit $\lambda \rightarrow \infty$, which is here even more remarkable, since frames containing one factor $(c+p-1)$ do arise, but those containing the factor $c-p = \mu\lambda^{-1}$ do not. Karabegov already stated in Theorem 3 [9] that for representative functions f, g the dependence of $f * g$ on λ^{-1} is rational with no poles for $\lambda \rightarrow 0$. For arbitrary smooth functions formula (46) states that the dependence on λ of the product is given by an infinite sum of rational functions, obviously in accordance with Karabegov's result. By a description of the star product via representative functions as it is done for the projective space in [14]

one could verify Karabegov's proposition directly for Grassmann manifolds. The expected isomorphism of the star algebras $(C^\infty(G_{p,q}(\mathbb{C})), *)$ and $(C^\infty(G_{q,p}(\mathbb{C})), *)$ can probably as well be established most easily in terms of representative functions. This would extend the duality of $G_{p,q}(\mathbb{C})$ and $G_{q,p}(\mathbb{C})$ to their star algebras.

Acknowledgement: I would like to thank M. Bordemann, whose idea of the projection method was actually the starting point of this work.

References

- [1] F. BEREZIN *Quantization*, Izv. Mat. Nauk **38**, 1974, 1109-1165
- [2] F. BAYEN, M. FLATO, C. FRONSDAL, A. LICHNEROWICZ, D. STERNHEIMER *Deformation Theory and Quantization* Ann. of Phys. **111**, 1978, 61-151
- [3] M. DEWILDE, P.B.A. LECOMTE *Existence of star products and of formal deformations of the Poisson Lie algebra of arbitrary symplectic manifolds* Lett. Math. Phys. **7**, 1983, 487-496
- [4] B. FEDOSOV *Formal Quantization in Some Topics of Modern Mathematics and their Applications to Problems of Mathematical Physics*, Moscow, 1985, 129-136
- [5] B. FEDOSOV *A simple geometrical construction of deformation quantization* J. Diff. Geom. **40**, 213-238 (1994)
- [6] H. OMORI, Y. MAEDA, A. YOSHIOKA *Weyl manifolds and deformation quantization* Adv. Math **85**, 1991, 224-255
- [7] C. MORENO **-Products on Some Kähler Manifolds*, Lett. Math. Phys. **11** 1986, 361-372
- [8] M. CAHEN, S. GUTT, J. RAWNSLEY *Quantization of Kähler manifolds II*, Trans. Am. Math. Soc. **337**, 1993, 73-98
- [9] A. V. KARABEGOV *Berezin's Quantization on flag manifolds and spherical modules*, funct-an/9412001
- [10] D. ARNAL, J. LUDWIG, M. MASMOUDI *Déformations covariantes sur les orbites polarisées d'un groupe de Lie*, J. Geom. Phys. **14**, 1994, 309-331
- [11] M. BORDEMANN, M. BRISCHLE, C. EMMRICH, S. WALDMANN *Phase Space Reduction for Star-Products: An explicit Construction for $\mathbb{C}P^n$* , Lett. Math. Phys. **36**, 1996, 357-371
- [12] R. ABRAHAM, J. E. MARSDEN *Foundations of Mechanics*, 2nd edition, Addison Wesley, Reading Mass. 1985
- [13] H. BOERNER *Darstellungen von Gruppen*, 2. Auflage, Springer, Berlin 1967
- [14] M. BORDEMANN, M. BRISCHLE, C. EMMRICH, S. WALDMANN *Subalgebras with converging star products in deformation quantization: An algebraic construction for $\mathbb{C}P^n$* , J. Math. Phys. **37** (12), 1996, 6311-6323