

(Noncanonical) field quantization by means of a single harmonic oscillator

Marek Czachor

*Katedra Fizyki Teoretycznej i Metod Matematycznych
Politechnika Gdańska, ul. Narutowicza 11/12, 80-952 Gdańsk, Poland*

and

*Arnold Sommerferld Institut für Mathematische Physik
Technische Universität Clausthal, 38678 Clausthal-Zellerfeld, Germany*

A new scheme of field quantization is proposed. Instead of associating with different frequencies different oscillators we begin with a single oscillator that can exist in a (quantum) superposition of different frequencies. The idea is applied to the electromagnetic radiation field and nonrelativistic quantum optics. Employing a Dirac-type mode-quantization of the electromagnetic field and using a single oscillator we obtain several standard properties such as coherent states or spontaneous and stimulated emission. Extending the formalism to a greater number of oscillators we arrive at a structure analogous to the Fock space but without the standard cyclic “vacuum state”. In the modified formalism the notion of the vacuum state is replaced by a vacuum *subspace* spanned by ground states of the oscillators. As opposed to the standard approach the vacuum energy is finite and does not have to be removed by any ad hoc procedure. Atom-light interactions are described by an appropriately modified minimal-coupling Hamiltonian (no normal ordering of the free-field Hamiltonian is necessary). The Hamiltonian does not change the number of oscillators which leads to an additional conservation law. Using the “ $-(e/m)\vec{A}\cdot\vec{p}$ ” interaction we discuss in second-order perturbation theory a two-photon spontaneous emission. The result essentially agrees with the ordinary formulas but the nontrivial vacuum structure is explicitly seen in the two-photon amplitude. The probability of the 2-photon emission resulting from the new formalism consists of a product of several terms, a part of them resembling those arising in the standard formulation from detector inefficiency, and the remaining one being the well known quantum optics formula. The presence of the additional conservation law shows that the theory contains two kinds of bosons (oscillators, whose number is conserved, and their excitations, whose number is not conserved). Taking this distinction into account we calculate an analog of the blackbody radiation Planck law. For temperatures lower than some T_{critical} the result is indistinguishable from the Planck distribution. For $T > T_{\text{critical}}$ the distribution is Planck-like but with the maximum lowered and shifted towards higher frequencies.

I. INTRODUCTION

The standard quantization of a harmonic oscillator is based on quantization of p and q but ω is a parameter. To have, say, two different frequencies one has to consider two independent oscillators. On the other hand, it is evident that there can exist oscillators which are in a *quantum superposition* of different frequencies. The example is an oscillator wave packet associated with distribution of center-of-mass momenta. It is known that the superposition of momenta gets translated into a superposition of Doppler shifts and therefore also of frequencies. We stress here the word “quantum” since the superpositions we have in mind are not those we know from *classical* oscillations.

This trivial observation raises the question of the role of superpositions of frequencies for a description of a single harmonic oscillator. The motivation behind the problem is associated with the question of field quantization: Is it possible that a quantum field consists of oscillators whose frequencies are *indefinite*? If so, maybe to quantize the field it is sufficient to use only one oscillator which exists in a *quantum* superposition of all the possible frequencies allowed by the boundary conditions of a given problem?

The idea is very simple. It is known that a “one-particle” state vector can be regarded as a representation of an ensemble of particles in a given pure state. On the other hand, the classical electromagnetic field can be regarded as an ensemble of oscillators. The standard idea of quantization, going back to 1925 [1], is to treat the field as an ensemble of quantum oscillators. But the ensemble itself is, in a sense, a classical one since for each frequency we need a separate oscillator. This is analogous to a classical ensemble of particles forming a classical wave on a lake surface. For each point on the surface we need a separate particle because a classical particle can occupy only a single point in space. A quantum wave is of course different and we are all accustomed to the idea of a single-particle wave. In this case the properties of the entire ensemble are somehow encoded in properties of a single element of the ensemble.

For some reasons, probably partly historical and sociological, it seems that the idea of a single-particle state vector representation of the ensemble of oscillators has never been considered. The historical reason may be the fact that the very concept of field quantization occurred already in 1925. At that stage quantum mechanics existed still in a

matrix form and the Schrödinger paper “Quantisierung als Eigenwertproblem” [2], where the Schrödinger equation occurred for the first time and the role of eigenvalues was explained, was not yet published. Sociologically, the names and reputation of Heisenberg, Born, Jordan, Dirac, together with the unquestionable success of quantum optics, field theory, and statistical physics, made it almost impossible to question the very starting point of the theory. The ideas presented below are an accidental by-product of a work on a different problem.

It should be mentioned that several approaches towards an alternative description of the electromagnetic field at a fundamental level were proposed (e.g. Janes’ [3] neoclassical theory, stochastic electrodynamics [4]). But the main idea of all such alternatives was to treat the field in classical terms and to associate the observed discreteness of emission/absorption phenomena with the quantum nature of atoms and not with the field itself.

The approach we will discuss in this paper does not belong to this tradition, is much more radical and goes in the opposite direction. We will not try to make the field more classical. What we will try to do is to make it even *more quantum* by replacing classical parameters with eigenvalues.

The field will be quantized at a one-particle level, but then extended to multi-particle systems. Obviously, it is not possible to include in a single paper all the possible tests of the new formalism one should perform. We will therefore concentrate on these points where quantum electrodynamics produces results which are believed to be a consequence of the standard canonical quantization. Three areas should be checked first:

- (i) Vacuum effects in atomic physics.
- (ii) Emission of photons in entangled states.
- (iii) Boson statistics and the Planck law.

For the first two problems we shall choose the simplest approach, namely first and second order perturbation theory. The Planck law will be discussed in a more detailed way. As we shall see the new theory is not completely equivalent to the standard one, but the modifications one finds are surprisingly subtle and in principle subject to experimental tests.

In a separate paper we shall discuss perturbation theory to all orders, since then a kind of prescription for translating the old results into the new framework may appear.

II. HARMONIC OSCILLATOR IN SUPERPOSITION OF FREQUENCIES

We know that frequency is typically associated with an eigenvalue of some Hamiltonian or, which is basically the same, with boundary conditions. A natural way of incorporating different frequencies into a single harmonic oscillator is by means of the *frequency operator*

$$\Omega = \sum_{\omega_k, j_k} \omega_k |\omega_k, j_k\rangle \langle \omega_k, j_k| \quad (1)$$

where all $\omega_k \geq 0$. For simplicity we have limited the discussion to the discrete spectrum but it is useful to include from the outset the possibility of degeneracies, represented here by the additional discrete quantum numbers j_k . The corresponding Hamiltonian is defined by

$$H = \hbar \Omega \otimes \frac{1}{2} (a^\dagger a + a a^\dagger) \quad (2)$$

where $a = \sum_{n=0}^{\infty} \sqrt{n+1} |n\rangle \langle n+1|$. The eigenstates of H are $|\omega_k, j_k, n\rangle$ and satisfy the required formula

$$H |\omega_k, j_k, n\rangle = \hbar \omega_k \left(n + \frac{1}{2} \right) |\omega_k, j_k, n\rangle \quad (3)$$

justifying our choice of H . The standard case of the oscillator whose frequency is just ω corresponds either to $\Omega = \omega \mathbf{1}$ or to the subspace spanned by $|\omega_k, j_k, n\rangle$ with fixed $\omega_k = \omega$. Introducing the operators

$$a_{\omega_k, j_k} = |\omega_k, j_k\rangle \langle \omega_k, j_k| \otimes a \quad (4)$$

we find that

$$H = \frac{1}{2} \sum_{\omega_k, j_k} \hbar \omega_k \left(a_{\omega_k, j_k}^\dagger a_{\omega_k, j_k} + a_{\omega_k, j_k} a_{\omega_k, j_k}^\dagger \right). \quad (5)$$

The algebra of the oscillator is “noncanonical”:

$$[a_{\omega_k, j_k}, a_{\omega_l, j_l}^\dagger] = \delta_{\omega_k \omega_l} \delta_{j_k j_l} |\omega_k, j_k\rangle \langle \omega_k, j_k| \otimes \mathbf{1} \quad (6)$$

$$a_{\omega_k, j_k} a_{\omega_l, j_l} = \delta_{\omega_k \omega_l} \delta_{j_k j_l} (a_{\omega_k, j_k})^2 \quad (7)$$

$$a_{\omega_k, j_k}^\dagger a_{\omega_l, j_l}^\dagger = \delta_{\omega_k \omega_l} \delta_{j_k j_l} (a_{\omega_k, j_k}^\dagger)^2. \quad (8)$$

The dynamics in the Schrödinger picture is given by

$$i\hbar \partial_t |\Psi\rangle = H |\Psi\rangle = \hbar \Omega \otimes (a^\dagger a + \frac{1}{2} \mathbf{1}) |\Psi\rangle. \quad (9)$$

In the Heisenberg picture we obtain the important formula (see Appendix XI A)

$$a_{\omega_k, j_k}(t) = e^{iHt/\hbar} a_{\omega_k, j_k} e^{-iHt/\hbar} \quad (10)$$

$$= |\omega_k, j_k\rangle \langle \omega_k, j_k| \otimes e^{-i\omega_k t} a = e^{-i\omega_k t} a_{\omega_k, j_k}. \quad (11)$$

Taking a general state

$$|\psi\rangle = \sum_{\omega_k, j_k, n} \psi(\omega_k, j_k, n) |\omega_k, j_k\rangle |n\rangle \quad (12)$$

we find that the average energy of the oscillator is

$$\langle H \rangle = \langle \psi | H | \psi \rangle = \sum_{\omega_k, j_k, n} |\psi(\omega_k, j_k, n)|^2 \hbar \omega_k \left(n + \frac{1}{2} \right). \quad (13)$$

The average clearly looks as an average energy of an *ensemble of different and independent oscillators*. The ground state of the ensemble, i.e. the one with $\psi(\omega_k, j_k, n > 0) = 0$ has energy

$$\langle H \rangle = \frac{1}{2} \sum_{\omega_k, j_k} |\psi(\omega_k, j_k, 0)|^2 \hbar \omega_k \quad (14)$$

which is finite if

$$\sum_{\omega_k, j_k} \psi(\omega_k, j_k, 0) |\omega_k, j_k\rangle \quad (15)$$

belongs to the domain of Ω . The result is not surprising but still quite remarkable if one thinks of the problem of field quantization.

The very idea of quantizing the electromagnetic field, as put forward by Born, Heisenberg, Jordan [1] and Dirac [5], is based on the observation that the mode decomposition of the electromagnetic energy is analogous to the energy of an ensemble of independent harmonic oscillators. In 1925, after the work of Heisenberg, it was clear what to do: One had to replace each classical oscillator by a quantum one. But since each oscillator had a definite frequency, to have an infinite number of different frequencies one needed an infinite number of oscillators. The price one paid for this assumption was the infinite energy of the electromagnetic vacuum.

The infinity is regarded as an “easy” one since one can get rid of it by redefining the Hamiltonian and removing the infinite term. The result looks correct and many properties typical of a *quantum* harmonic oscillator are indeed observed in electromagnetic field. However, subtraction of infinite terms is in mathematics as forbidden as division by zero so to avoid evident absurdities one is forced to invent various ad hoc regularizations whose only justification is that otherwise the theory would not work. In larger perspective (say, in cosmology) it is not at all clear that an infinite (or arbitrarily cut off at the Planck scale) energy of the vacuum does not lead to contradictions with observational data [6]. Finally, Dirac himself had never been fully satisfied by the theory he created. As Weinberg put it, Dirac’s “demand for a completely finite theory is similar to a host of other aesthetic judgements that theoretical physicists always need to make” [7].

The oscillator that can exist in superpositions of different frequencies is a natural candidate as a starting point for Dirac-type field quantization. Symbolically, if the Heisenberg quantization is $p^2 + \omega^2 q^2 \mapsto \hat{p}^2 + \omega^2 \hat{q}^2$, where ω is a parameter, the new scheme is $p^2 + \omega^2 q^2 \mapsto \hat{p}^2 + \hat{\omega}^2 \hat{q}^2$, where $\hat{\omega}$ is an operator. Its spectrum can be related to boundary conditions imposed on the fields. The field now can exist in superposition of frequencies but the superposition is meant in the quantum sense i.e. the field may consist of (an indefinite number) of oscillators with indefinite frequency. In this meaning the approach we propose is even “more quantum” than the standard one since ω is not a (classical) parameter but an eigenvalue.

We do not need to remove the ground state energy since in the Hilbert space of physical states the correction is finite. The question we have to understand is whether one can obtain the well known quantum properties of the radiation field by this type of quantization.

III. PRELUDE: “FIRST QUANTIZATION” — FIELD OPERATORS FOR FREE MAXWELL FIELDS

The new quantization will be performed in two steps. In this section we describe the first step, a kind of first quantization. In next sections we shall perform an analogue of second quantization which will lead to the final framework. It is essential that the “second quantization” will not involve, in fact, any additional quantization but is simply a transition from one to many oscillators.

The energy and momentum operators of the field are defined in analogy to H from the previous section

$$H = \sum_{s, \kappa_\lambda} \hbar \omega_\lambda |s, \vec{\kappa}_\lambda\rangle \langle s, \vec{\kappa}_\lambda| \otimes \frac{1}{2} (a^\dagger a + a a^\dagger) \quad (16)$$

$$= \frac{1}{2} \sum_{s, \kappa_\lambda} \hbar \omega_\lambda (a_{s, \kappa_\lambda}^\dagger a_{s, \kappa_\lambda} + a_{s, \kappa_\lambda} a_{s, \kappa_\lambda}^\dagger) \quad (17)$$

$$\vec{P} = \sum_{s, \kappa_\lambda} \hbar \vec{\kappa}_\lambda |s, \vec{\kappa}_\lambda\rangle \langle s, \vec{\kappa}_\lambda| \otimes \frac{1}{2} (a^\dagger a + a a^\dagger) \quad (18)$$

$$= \frac{1}{2} \sum_{s, \kappa_\lambda} \hbar \vec{\kappa}_\lambda (a_{s, \kappa_\lambda}^\dagger a_{s, \kappa_\lambda} + a_{s, \kappa_\lambda} a_{s, \kappa_\lambda}^\dagger) \quad (19)$$

where $s = \pm 1$ corresponds to circular polarizations. Denote $P = (H/c, \vec{P})$ and $P \cdot x = Ht - \vec{P} \cdot \vec{x}$. We employ the standard Dirac-type definitions for mode quantization in volume V

$$\hat{A}(t, \vec{x}) = \sum_{s, \kappa_\lambda} \sqrt{\frac{\hbar}{2\omega_\lambda V}} (a_{s, \kappa_\lambda} e^{-i\omega_\lambda t} \vec{e}_{s, \kappa_\lambda} e^{i\vec{\kappa}_\lambda \cdot \vec{x}} + a_{s, \kappa_\lambda}^\dagger e^{i\omega_\lambda t} \vec{e}_{s, \kappa_\lambda}^* e^{-i\vec{\kappa}_\lambda \cdot \vec{x}}) \quad (20)$$

$$= e^{iP \cdot x / \hbar} \hat{A} e^{-iP \cdot x / \hbar} \quad (21)$$

$$\hat{E}(t, \vec{x}) = i \sum_{s, \kappa_\lambda} \sqrt{\frac{\hbar \omega_\lambda}{2V}} (a_{s, \kappa_\lambda} e^{-i\omega_\lambda t} e^{i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda} - a_{s, \kappa_\lambda}^\dagger e^{i\omega_\lambda t} e^{-i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda}^*) \quad (22)$$

$$= e^{iP \cdot x / \hbar} \hat{E} e^{-iP \cdot x / \hbar} \quad (23)$$

$$\quad (24)$$

$$\hat{B}(t, \vec{x}) = i \sum_{s, \kappa_\lambda} \sqrt{\frac{\hbar \omega_\lambda}{2V}} \vec{n}_{\kappa_\lambda} \times (a_{s, \kappa_\lambda} e^{-i\omega_\lambda t} e^{i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda} - a_{s, \kappa_\lambda}^\dagger e^{i\omega_\lambda t} e^{-i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda}^*) \quad (25)$$

$$= e^{iP \cdot x / \hbar} \hat{B} e^{-iP \cdot x / \hbar}, \quad (26)$$

where

$$a_{s, \vec{\kappa}_\lambda} = |s, \vec{\kappa}_\lambda\rangle \langle s, \vec{\kappa}_\lambda| \otimes a \quad (27)$$

$$a_{s, \vec{\kappa}_\lambda}^\dagger = |s, \vec{\kappa}_\lambda\rangle \langle s, \vec{\kappa}_\lambda| \otimes a^\dagger. \quad (28)$$

For later purposes we introduce the notation

$$[a_{s, \vec{\kappa}_\lambda}, a_{s, \vec{\kappa}_\lambda}^\dagger] = 1_{s, \vec{\kappa}_\lambda} = |s, \vec{\kappa}_\lambda\rangle \langle s, \vec{\kappa}_\lambda| \otimes \mathbf{1}. \quad (29)$$

Now take a state (say, in the Heisenberg picture)

$$|\Psi\rangle = \sum_{s, \vec{\kappa}_\lambda, n} \Psi_{s, \vec{\kappa}_\lambda, n} |s, \vec{\kappa}_\lambda, n\rangle \quad (30)$$

$$= \sum_{s, \vec{\kappa}_\lambda} \Phi_{s, \vec{\kappa}_\lambda} |s, \vec{\kappa}_\lambda\rangle |\alpha_{s, \vec{\kappa}_\lambda}\rangle \quad (31)$$

where $|\alpha_{s, \vec{\kappa}_\lambda}\rangle$ form a family of one-oscillator coherent states:

$$a |\alpha_{s, \vec{\kappa}_\lambda}\rangle = \alpha_{s, \vec{\kappa}_\lambda} |\alpha_{s, \vec{\kappa}_\lambda}\rangle \quad (32)$$

The averages of the field operators are

$$\langle \Psi | \hat{A}(t, \vec{x}) | \Psi \rangle = \sum_{s, \kappa_\lambda} |\Phi_{s, \vec{\kappa}_\lambda}|^2 \sqrt{\frac{\hbar}{2\omega_\lambda V}} \left(\alpha_{s, \kappa_\lambda} e^{-i\kappa_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda} + \alpha_{s, \kappa_\lambda}^* e^{i\kappa_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda}^* \right) \quad (33)$$

$$\langle \Psi | \hat{E}(t, \vec{x}) | \Psi \rangle = \sum_{s, \kappa_\lambda} |\Phi_{s, \vec{\kappa}_\lambda}|^2 \sqrt{\frac{\hbar\omega_\lambda}{2V}} \left(\alpha_{s, \kappa_\lambda} e^{-i\kappa_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda} - \alpha_{s, \kappa_\lambda}^* e^{i\kappa_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda}^* \right) \quad (34)$$

$$\langle \Psi | \hat{B}(t, \vec{x}) | \Psi \rangle = i \sum_{s, \kappa_\lambda} |\Phi_{s, \vec{\kappa}_\lambda}|^2 \sqrt{\frac{\hbar\omega_\lambda}{2V}} \left(\alpha_{s, \kappa_\lambda} e^{-i\kappa_\lambda \cdot \vec{x}} \vec{n}_{\kappa_\lambda} \times \vec{e}_{s, \kappa_\lambda} - \alpha_{s, \kappa_\lambda}^* e^{i\kappa_\lambda \cdot \vec{x}} \vec{n}_{\kappa_\lambda} \times \vec{e}_{s, \kappa_\lambda}^* \right) \quad (35)$$

These are just the classical fields. More precisely, the fields look like averages of monochromatic coherent states with probabilities $|\Phi_{s, \vec{\kappa}_\lambda}|^2$. The energy-momentum operators satisfy also the standard relations (see Appendix XI B)

$$H = \frac{1}{2} \int_V d^3x \left(\hat{E}(t, \vec{x}) \cdot \hat{E}(t, \vec{x}) + \hat{B}(t, \vec{x}) \cdot \hat{B}(t, \vec{x}) \right), \quad (36)$$

$$\vec{P} = \int_V d^3x \hat{E}(t, \vec{x}) \times \hat{B}(t, \vec{x}). \quad (37)$$

It should be stressed, however, that these relations have a completely different mathematical origin than in the usual formalism where the integrals are necessary in order to make plane waves into an orthonormal basis. Here orthogonality follows from the presence of the projectors in the definition of a_{s, κ_λ} and the integration in itself is *trivial* since

$$\hat{E}(t, \vec{x}) \cdot \hat{E}(t, \vec{x}) + \hat{B}(t, \vec{x}) \cdot \hat{B}(t, \vec{x}) = \hat{E} \cdot \hat{E} + \hat{B} \cdot \hat{B} \quad (38)$$

$$\hat{E}(t, \vec{x}) \times \hat{B}(t, \vec{x}) = \hat{E} \times \hat{B}. \quad (39)$$

Therefore the role of the integral is simply to produce the factor V which cancels with $1/V$ arising from the term $1/\sqrt{V}$ occuring in the mode decomposition of the fields. To end this section let us note that

$$\langle \Psi | H | \Psi \rangle = \sum_{s, \kappa_\lambda} \hbar\omega_\lambda |\Phi_{s, \kappa_\lambda}|^2 \left(|\alpha_{s, \kappa_\lambda}|^2 + \frac{1}{2} \right) \quad (40)$$

$$\langle \Psi | \vec{P} | \Psi \rangle = \sum_{s, \kappa_\lambda} \hbar\vec{\kappa}_\lambda |\Phi_{s, \kappa_\lambda}|^2 \left(|\alpha_{s, \kappa_\lambda}|^2 + \frac{1}{2} \right). \quad (41)$$

The contribution from the vacuum fluctuations is nonzero but *finite*. One can phrase the latter property also as follows. The noncanonical algebra of creation-annihilation operators satisfies the resolution of identity

$$\sum_{s, \kappa_\lambda} [a_{s, \vec{\kappa}_\lambda}, a_{s, \vec{\kappa}_\lambda}^\dagger] = \mathbf{1} \quad (42)$$

wheras the canonical algebra would imply

$$\sum_{s, \kappa_\lambda} [a_{s, \vec{\kappa}_\lambda}, a_{s, \vec{\kappa}_\lambda}^\dagger] = \infty \mathbf{1}. \quad (43)$$

IV. SPONTANEOUS AND STIMULATED EMISSION: FIRST VERSION

Some typically quantum optical phenomena occur already at the one-oscillator level. Below we shall see that spontaneous and stimulated emissions are a property of a single-oscillator description, although to have a more complete picture we need the multi-oscillator extension discussed in subsequent sections.

Beginning with the dipole and rotating wave approximations (RWA) we arrive at the Hamiltonian

$$H = \frac{1}{2} \hbar\omega_0 \sigma_3 + \frac{1}{2} \sum_{s, \vec{\kappa}_\lambda} \hbar\omega_\lambda \left(a_{s, \vec{\kappa}_\lambda}^\dagger a_{s, \vec{\kappa}_\lambda} + a_{s, \vec{\kappa}_\lambda} a_{s, \vec{\kappa}_\lambda}^\dagger \right) + \hbar\omega_0 d \sum_{s, \vec{\kappa}_\lambda} \left(g_{s, \vec{\kappa}_\lambda} a_{s, \vec{\kappa}_\lambda} \sigma_+ + g_{s, \vec{\kappa}_\lambda}^* a_{s, \vec{\kappa}_\lambda}^\dagger \sigma_- \right) \quad (44)$$

where $d\vec{u} = \langle +|\hat{d}|- \rangle$ is the matrix element of the dipole moment evaluated between the excited and ground states, and $g_{s,\vec{k}_\lambda} = i\sqrt{\frac{1}{2\hbar\omega_\lambda V}}\vec{e}_{s,\vec{k}_\lambda} \cdot \vec{u}$. The Hamiltonian represents a two-level atom located at $\vec{x}_0 = 0$.

The Hamiltonian in the interaction picture has the well known form

$$H_I = \hbar\omega_0 d \sum_{s,\vec{k}_\lambda} \left(g_{s,\vec{k}_\lambda} e^{i(\omega_0 - \omega_\lambda)t} a_{s,\vec{k}_\lambda} \sigma_+ + g_{s,\vec{k}_\lambda}^* e^{-i(\omega_0 - \omega_\lambda)t} a_{s,\vec{k}_\lambda}^\dagger \sigma_- \right). \quad (45)$$

Consider the initial state

$$\begin{aligned} |\Psi(0)\rangle &= \sum_{s',\vec{k}_{\lambda'},m} \Psi_{s',\vec{k}_{\lambda'},m} |s',\vec{k}_{\lambda'},m,+\rangle \\ &= \sum_{s',\vec{k}'_0} \Psi_{s',\vec{k}'_0,0} |s',\vec{k}'_0,0,+\rangle + \sum_{s',\vec{k}'_n} \Psi_{s',\vec{k}'_n,n} |s',\vec{k}'_n,n,+\rangle. \end{aligned} \quad (46)$$

The states corresponding to $n = 0$ play a role of a vacuum. As a consequence the vacuum is not represented here by a unique vector, but rather by a subspace of the Hilbert space of states. Energy of the general vacuum state

$$|\Psi\rangle = \sum_{s,\vec{k}_\lambda,\pm} \Psi_{s,\vec{k}_\lambda,0,\pm} |s,\vec{k}_\lambda,0,\pm\rangle \quad (47)$$

is related to the density of modes $\rho(\vec{k}_\lambda) = \sum_{s,\pm} |\Psi_{s,\vec{k}_\lambda,0,\pm}|^2$ and is, therefore, state dependent.

In order to estimate the probabilities of spontaneous and stimulated emissions we can use the first-order time-dependent perturbation theory [8] and arrive at

$$\begin{aligned} |\Psi(t)\rangle &= |\Psi(0)\rangle \\ &+ \omega_0 d \sum_{s,\vec{k}_0} \frac{e^{-i(\omega_0 - \omega_{\lambda_0})t} - 1}{\omega_0 - \omega_{\lambda_0}} \Psi_{s,\vec{k}_{\lambda_0},0} g_{s,\vec{k}_{\lambda_0}}^* |s,\vec{k}_{\lambda_0},1,-\rangle \\ &+ \omega_0 d \sum_{s,\vec{k}_n} \frac{e^{-i(\omega_0 - \omega_{\lambda_n})t} - 1}{\omega_0 - \omega_{\lambda_n}} \Psi_{s,\vec{k}_n,n} \sqrt{n+1} g_{s,\vec{k}_n}^* |s,\vec{k}_n,n+1,-\rangle. \end{aligned} \quad (48)$$

One recognizes here the well known contributions from spontaneous and stimulated emissions. It should be stressed that although the final result looks familiar, the mathematical details behind the calculation are different from what we are accustomed to. For example, instead of

$$a_{s_1,\vec{k}_1}^\dagger |s,\vec{k},m\rangle \sim |s_1,\vec{k}_1,1;s,\vec{k},m\rangle, \quad (49)$$

which would hold in the standard formalism for $\vec{k}_1 \neq \vec{k}$, we get simply

$$a_{s_1,\vec{k}_1}^\dagger |s,\vec{k},m\rangle = 0, \quad (50)$$

a consequence of $a_{s_1,\vec{k}_1}^\dagger a_{s,\vec{k}}^\dagger = 0$.

Let us now look more closely at spontaneous emission (we take $n = 0$). The state vector is

$$\begin{aligned} |\Psi(t)\rangle &= |\Psi(0)\rangle \\ &+ \omega_0 d \sum_{s,\vec{k}_0} \frac{e^{-i(\omega_0 - \omega_{\lambda_0})t} - 1}{\omega_0 - \omega_{\lambda_0}} G_{s,\vec{k}_{\lambda_0}}^* |s,\vec{k}_{\lambda_0},1,-\rangle \end{aligned} \quad (51)$$

where we have introduced the ‘‘effective coupling terms’’

$$G_{s,\vec{k}_{\lambda_0}}^* = \Psi_{s,\vec{k}_{\lambda_0},0} g_{s,\vec{k}_{\lambda_0}}^*. \quad (52)$$

As we can see the result is mathematically equivalent to the standard one but with the coupling constants automatically regularized by the presence of the vacuum amplitude in $G_{s,\vec{k}_{\lambda_0}}^*$.

We apply the standard argument but with g 's replaced by G 's and obtain the probability of emission per time unit

$$P = 2\pi\omega_0^2 d^2 \sum_{s, \vec{k}} |\Psi_{s, \vec{k}, 0} g_{s, \vec{k}}|^2 \delta(\omega_0 - \omega_{\vec{k}}). \quad (53)$$

Assuming for simplicity that density of vacuum modes is isotropic and polarization independent we can write it as a function of frequency only, i.e.

$$|\Psi_{s, \vec{k}, 0}|^2 = F(\omega_{\vec{k}}) \quad (54)$$

and finally

$$P = 2\pi\omega_0^2 d^2 F(\omega_0) \sum_{s, \vec{k}} |g_{s, \vec{k}}|^2 \delta(\omega_0 - \omega_{\vec{k}}) = F(\omega_0) P_{\text{old}}. \quad (55)$$

Here P_{old} is the emission rate obtained in the standard theory. The nontrivial structure of the vacuum influences the lifetime of the atom. We shall return to this and related questions later but first have to extend the formalism in a way allowing to consider entangled states of light.

More reliable estimates of the lifetime require more detailed calculations that we postpone to a forthcoming paper. One should also keep in mind the possibility of an unisotropic vacuum caused by more complicated boundary conditions such as those occurring in measurements of the Casimir force.

V. "SECOND QUANTIZATION"

The Hilbert space of states of the field we have constructed is spanned by vectors $|s, \vec{k}, n\rangle$. Still there is no doubt that both in reality (and the standard formalism) there exist multiparticle entangled states such as those spanned by tensor products of the form

$$|+, \vec{k}_1, 1\rangle |-, \vec{k}_2, 1\rangle, \quad (56)$$

and the similar. It seems that there is no reason to limit our discussion to a *single* Hilbert space of a *single* oscillator. What we have done so far was a quantization of the electromagnetic field at the level of a "one-particle" Hilbert space. Similarly to quantization of other physical systems the next step is to consider many (noninteracting) particles.

The procedure is essentially clear. Having the one-particle energy-momentum operators P_a (i.e. generators of 4-translations in the 1-particle Hilbert space) we define in the standard way their extensions to the Fock-type space

$$\begin{aligned} \mathcal{P}_a &= P_a \\ &\oplus (P_a \otimes \mathbf{1} + \mathbf{1} \otimes P_a) \\ &\oplus (P_a \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes P_a \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes P_a) \\ &\oplus \dots \end{aligned} \quad (57)$$

The x -dependence of fields is introduced similarly to the one-particle level

$$\vec{\mathcal{F}}(t, \vec{x}) = e^{i\mathcal{P}\cdot x/\hbar} \vec{\mathcal{F}} e^{-i\mathcal{P}\cdot x/\hbar} \quad (58)$$

but the field itself has yet to be defined. Assume

$$\begin{aligned} \vec{\mathcal{F}} &= c_1 \vec{F} \\ &\oplus c_2 (\vec{F} \otimes \mathbf{1} + \mathbf{1} \otimes \vec{F}) \\ &\oplus c_3 (\vec{F} \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes \vec{F} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes \vec{F}) \\ &\oplus \dots \end{aligned} \quad (59)$$

where c_k are constants discussed below, and \vec{F} is \vec{A} , \vec{E} , or \vec{B} . The multi-oscillator annihilation operator associated with such fields must be therefore of the form

$$\begin{aligned} \mathbf{a}_{s, \vec{k}} &= c_1 a_{s, \vec{k}} \\ &\oplus c_2 (a_{s, \vec{k}} \otimes \mathbf{1} + \mathbf{1} \otimes a_{s, \vec{k}}) \\ &\oplus c_3 (a_{s, \vec{k}} \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes a_{s, \vec{k}} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes a_{s, \vec{k}}) \\ &\oplus \dots \end{aligned} \quad (60)$$

Having two 1-particle operators, say X and Y , one can easily establish a relation between the 1-particle commutator $[X, Y]$ and the commutator of the extensions \mathcal{X}, \mathcal{Y} :

$$\begin{aligned} [\mathcal{X}, \mathcal{Y}] &= c_1^2[X, Y] \\ &\oplus c_2^2([X, Y] \otimes \mathbf{1} + \mathbf{1} \otimes [X, Y]) \\ &\oplus c_3^2([X, Y] \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes [X, Y] \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes [X, Y]) \\ &\oplus \dots \end{aligned} \quad (61)$$

The annihilation operators so defined satisfy therefore the algebra

$$[\mathbf{a}_{s, \vec{\kappa}}, \mathbf{a}_{s', \vec{\kappa}'}^\dagger] = 0 \quad \text{for } (s, \vec{\kappa}) \neq (s', \vec{\kappa}'), \quad (62)$$

$$[\mathbf{a}_{s, \vec{\kappa}}, \mathbf{a}_{s, \vec{\kappa}}^\dagger] = \mathbf{1}_{s, \vec{\kappa}}, \quad (63)$$

$$[\mathbf{a}_{s, \vec{\kappa}}, \mathbf{a}_{s', \vec{\kappa}'}] = 0 \quad (64)$$

$$[\mathbf{a}_{s, \vec{\kappa}}^\dagger, \mathbf{a}_{s', \vec{\kappa}'}^\dagger] = 0 \quad (65)$$

where the operator $\mathbf{1}_{s, \vec{\kappa}}$ is defined by

$$\begin{aligned} \mathbf{1}_{s, \vec{\kappa}} &= c_1^2 \mathbf{1}_{s, \vec{\kappa}} \\ &\oplus c_2^2 (\mathbf{1}_{s, \vec{\kappa}} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1}_{s, \vec{\kappa}}) \\ &\oplus c_3^2 (\mathbf{1}_{s, \vec{\kappa}} \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1}_{s, \vec{\kappa}} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}_{s, \vec{\kappa}}) \\ &\oplus \dots, \end{aligned} \quad (66)$$

and $\mathbf{1}_{s, \vec{\kappa}}$ is a single-oscillator operator (29).

As opposed to the single-oscillator case

$$\mathbf{a}_{s, \vec{\kappa}} \mathbf{a}_{s', \vec{\kappa}'}^\dagger \neq \delta_{ss'} \delta_{\vec{\kappa}, \vec{\kappa}'} (\mathbf{a}_{s, \vec{\kappa}})^2. \quad (67)$$

An important property of the 1-oscillator description was the resolution of identity (42). The requirement that the same be valid at the multi oscillator level leads to $c_n = 1/\sqrt{n}$. In such a case one finds that

$$\mathbf{1}_{s, \vec{\kappa}}^2 \neq \mathbf{1}_{s, \vec{\kappa}} \quad (68)$$

but nevertheless

$$\sum_{s, \vec{\kappa}} \mathbf{1}_{s, \vec{\kappa}} = \mathbf{1}. \quad (69)$$

Below we shall give another justification of this particular choice of c_n .

We can finally write

$$\vec{\mathcal{A}}(t, \vec{x}) = \sum_{s, \kappa_\lambda} \sqrt{\frac{\hbar}{2\omega_\lambda V}} \left(\mathbf{a}_{s, \kappa_\lambda} e^{-i\omega_\lambda t} \vec{e}_{s, \kappa_\lambda} e^{i\vec{\kappa}_\lambda \cdot \vec{x}} + \mathbf{a}_{s, \kappa_\lambda}^\dagger e^{i\omega_\lambda t} \vec{e}_{s, \kappa_\lambda}^* e^{-i\vec{\kappa}_\lambda \cdot \vec{x}} \right) \quad (70)$$

$$= e^{i\mathcal{P} \cdot \vec{x} / \hbar} \vec{\mathcal{A}} e^{-i\mathcal{P} \cdot \vec{x} / \hbar} \quad (71)$$

$$\vec{\mathcal{E}}(t, \vec{x}) = i \sum_{s, \kappa_\lambda} \sqrt{\frac{\hbar\omega_\lambda}{2V}} \left(\mathbf{a}_{s, \kappa_\lambda} e^{-i\omega_\lambda t} e^{i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda} - \mathbf{a}_{s, \kappa_\lambda}^\dagger e^{i\omega_\lambda t} e^{-i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda}^* \right) \quad (72)$$

$$= e^{i\mathcal{P} \cdot \vec{x} / \hbar} \vec{\mathcal{E}} e^{-i\mathcal{P} \cdot \vec{x} / \hbar} \quad (73)$$

$$\vec{\mathcal{B}}(t, \vec{x}) = i \sum_{s, \kappa_\lambda} \sqrt{\frac{\hbar\omega_\lambda}{2V}} \vec{n}_{\kappa_\lambda} \times \left(\mathbf{a}_{s, \kappa_\lambda} e^{-i\omega_\lambda t} e^{i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda} - \mathbf{a}_{s, \kappa_\lambda}^\dagger e^{i\omega_\lambda t} e^{-i\vec{\kappa}_\lambda \cdot \vec{x}} \vec{e}_{s, \kappa_\lambda}^* \right) \quad (74)$$

$$= e^{i\mathcal{P} \cdot \vec{x} / \hbar} \vec{\mathcal{B}} e^{-i\mathcal{P} \cdot \vec{x} / \hbar}. \quad (75)$$

These operators form a basis of the modified version of nonrelativistic quantum optics.

Let us return for the moment to the case of a general c_n . A straightforward calculation shows that

$$\mathbf{H} = \frac{1}{2} \int_V d^3x \left(\vec{\mathcal{E}}(t, \vec{x}) \cdot \vec{\mathcal{E}}(t, \vec{x}) + \vec{\mathcal{B}}(t, \vec{x}) \cdot \vec{\mathcal{B}}(t, \vec{x}) \right) = \frac{1}{2} \sum_{s, \kappa_\lambda} \hbar \omega_\lambda \left(\mathbf{a}_{s, \kappa_\lambda}^\dagger \mathbf{a}_{s, \kappa_\lambda} + \mathbf{a}_{s, \kappa_\lambda} \mathbf{a}_{s, \kappa_\lambda}^\dagger \right) \quad (77)$$

$$= \frac{1}{2} \sum_{s, \kappa_\lambda} \hbar \omega_\lambda \left[c_1^2 \{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \right. \\ \oplus c_2^2 \left(\{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \otimes \mathbf{1} + \mathbf{1} \otimes \{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} + 2a_{s, \vec{\kappa}}^\dagger \otimes a_{s, \vec{\kappa}} + 2a_{s, \vec{\kappa}} \otimes a_{s, \vec{\kappa}}^\dagger \right) \\ \oplus c_3^2 \left(\{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \otimes \mathbf{1} \otimes \mathbf{1} + 2a_{s, \vec{\kappa}} \otimes a_{s, \vec{\kappa}}^\dagger \otimes \mathbf{1} + 2a_{s, \vec{\kappa}} \otimes \mathbf{1} \otimes a_{s, \vec{\kappa}}^\dagger \right. \\ \left. + 2a_{s, \vec{\kappa}}^\dagger \otimes a_{s, \vec{\kappa}} \otimes \mathbf{1} + \mathbf{1} \otimes \{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \otimes \mathbf{1} + 2\mathbf{1} \otimes a_{s, \vec{\kappa}} \otimes a_{s, \vec{\kappa}}^\dagger \right. \\ \left. + 2a_{s, \vec{\kappa}}^\dagger \otimes \mathbf{1} \otimes a_{s, \vec{\kappa}} + 2\mathbf{1} \otimes a_{s, \vec{\kappa}}^\dagger \otimes a_{s, \vec{\kappa}} + \mathbf{1} \otimes \mathbf{1} \otimes \{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \right) \\ \left. \oplus \dots \right] \quad (78)$$

Comparing this with the generator of time translations

$$\mathcal{H} = c\mathcal{P}_0 = \frac{1}{2} \sum_{s, \kappa_\lambda} \hbar \omega_\lambda \left[\{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \right. \\ \oplus \left(\{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \otimes \mathbf{1} + \mathbf{1} \otimes \{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \right) \\ \oplus \left(\{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes \{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes \{a_{s, \vec{\kappa}}, a_{s, \vec{\kappa}}^\dagger\} \right) \\ \left. \oplus \dots \right] \quad (79)$$

we can see that there is a relation between \mathcal{H} and \mathbf{H} but the latter contains terms describing interactions between the oscillators. The contribution from these interactions vanishes on vacuum states. Below, when we introduce the notion of a generalized coherent state, we will be able to relate averages of \mathcal{H} and \mathbf{H} . In a similar way one can introduce the ‘‘Pointing operator’’

$$\mathbf{P} = \int_V d^3x \vec{\mathcal{E}}(t, \vec{x}) \times \vec{\mathcal{B}}(t, \vec{x}) = \frac{1}{2} \sum_{s, \kappa_\lambda} \hbar \vec{\kappa}_\lambda \left(\mathbf{a}_{s, \kappa_\lambda}^\dagger \mathbf{a}_{s, \kappa_\lambda} + \mathbf{a}_{s, \kappa_\lambda} \mathbf{a}_{s, \kappa_\lambda}^\dagger \right). \quad (80)$$

Its relation to the generator of 3-translations $\vec{\mathcal{P}}$ is similar to this between \mathcal{H} and \mathbf{H} .

In the above construction the only element which is beyond a simple transition to many oscillators is the choice of c_n . For different choices of these constants we obtain different algebras of noncanonical commutation relations and therefore also different quantization schemes. Several different ways of reasoning lead to $c_n = 1/\sqrt{n}$ as we shall also see in the next sections.

VI. SOME PARTICULAR STATES

We assume that all the multi-oscillator states are symmetric with respect to permutations of the oscillators.

A. Generalized coherent states

For general c_n an eigenstate of $\mathbf{a}_{s, \kappa_\lambda}$ corresponding to the eigenvalue $\alpha_{s, \vec{\kappa}}$ is of the form

$$|\alpha_{s, \vec{\kappa}}\rangle = f_1(s, \vec{\kappa}) |s, \vec{\kappa}, \alpha_{s, \vec{\kappa}}/c_1\rangle \\ \oplus f_2(s, \vec{\kappa}) |s, \vec{\kappa}, \alpha_{s, \vec{\kappa}}/(2c_2)\rangle |s, \vec{\kappa}, \alpha_{s, \vec{\kappa}}/(2c_2)\rangle \\ \oplus f_3(s, \vec{\kappa}) |s, \vec{\kappa}, \alpha_{s, \vec{\kappa}}/(3c_3)\rangle |s, \vec{\kappa}, \alpha_{s, \vec{\kappa}}/(3c_3)\rangle |s, \vec{\kappa}, \alpha_{s, \vec{\kappa}}/(3c_3)\rangle \\ \oplus \dots \quad (81)$$

where

$$|s, \vec{\kappa}, \alpha_{s, \vec{\kappa}}\rangle = |s, \vec{\kappa}\rangle |\alpha_{s, \vec{\kappa}}\rangle, \quad (82)$$

$\sum_k |f_k(s, \vec{\kappa})|^2 = 1$, and $a|\alpha_{s, \vec{\kappa}}\rangle = \alpha_{s, \vec{\kappa}}|\alpha_{s, \vec{\kappa}}\rangle$. What is interesting not all f_k have to be nonvanishing. The average ‘‘energies’’ of the field in the above eigenstate are

$$\langle \alpha_{s, \vec{\kappa}} | \mathcal{H} | \alpha_{s, \vec{\kappa}} \rangle = \hbar \omega_{\vec{\kappa}} |\alpha_{s, \vec{\kappa}}|^2 \sum_{k=1}^{\infty} \frac{1}{k c_k^2} |f_k(s, \vec{\kappa})|^2 + \frac{1}{2} \hbar \omega_{\vec{\kappa}} \sum_{k=1}^{\infty} k |f_k(s, \vec{\kappa})|^2 \quad (83)$$

and

$$\langle \alpha_{s, \vec{\kappa}} | \mathbf{H} | \alpha_{s, \vec{\kappa}} \rangle = \hbar \omega_{\vec{\kappa}} |\alpha_{s, \vec{\kappa}}|^2 + \frac{1}{2} \hbar \omega_{\vec{\kappa}} \sum_{k=1}^{\infty} k c_k^2 |f_k(s, \vec{\kappa})|^2. \quad (84)$$

The two averages will differ only by the value of the vacuum contribution if $c_k = 1/\sqrt{k}$ which leads back to the above mentioned choice of c_k . With this choice and taking the general combination of coherent states

$$|\Psi\rangle = \sum_{s, \vec{\kappa}} \Phi_{s, \vec{\kappa}} |\alpha_{s, \vec{\kappa}}\rangle \quad (85)$$

we find

$$\langle \Psi | \mathcal{H} | \Psi \rangle = \sum_{s, \vec{\kappa}} \hbar \omega_{\vec{\kappa}} |\Phi_{s, \vec{\kappa}}|^2 |\alpha_{s, \vec{\kappa}}|^2 + \frac{1}{2} \sum_{s, \vec{\kappa}} \hbar \omega_{\vec{\kappa}} |\Phi_{s, \vec{\kappa}}|^2 \sum_{k=1}^{\infty} k |f_k(s, \vec{\kappa})|^2 \quad (86)$$

and

$$\langle \Psi | \mathbf{H} | \Psi \rangle = \sum_{s, \vec{\kappa}} \hbar \omega_{\vec{\kappa}} |\Phi_{s, \vec{\kappa}}|^2 |\alpha_{s, \vec{\kappa}}|^2 + \frac{1}{2} \sum_{s, \vec{\kappa}} \hbar \omega_{\vec{\kappa}} |\Phi_{s, \vec{\kappa}}|^2. \quad (87)$$

B. Vacuum

Similarly to the one-oscillator case the traditional notion of a vacuum state is replaced in our formalism by a vacuum *subspace* consisting of all the vectors of the form

$$\begin{aligned} |\Psi\rangle &= \sum_{s, \vec{\kappa}_\lambda} \Psi_{s, \vec{\kappa}_\lambda, 0}^{(1)} |s, \vec{\kappa}_\lambda, 0\rangle \\ &\oplus \sum_{s_j, \vec{\kappa}_{\lambda_j}} \Psi_{s_1, s_2, \vec{\kappa}_{\lambda_1}, \vec{\kappa}_{\lambda_2}, 0, 0}^{(2)} |s_1, \vec{\kappa}_{\lambda_1}, 0\rangle |s_2, \vec{\kappa}_{\lambda_2}, 0\rangle \\ &\oplus \sum_{s_j, \vec{\kappa}_{\lambda_j}} \Psi_{s_1, s_2, s_3, \vec{\kappa}_{\lambda_1}, \vec{\kappa}_{\lambda_2}, \vec{\kappa}_{\lambda_3}, 0, 0, 0}^{(3)} |s_1, \vec{\kappa}_{\lambda_1}, 0\rangle |s_2, \vec{\kappa}_{\lambda_2}, 0\rangle |s_3, \vec{\kappa}_{\lambda_3}, 0\rangle \\ &\oplus \dots \end{aligned} \quad (88)$$

It seems that there is no reason for introducing the standard ‘‘vacuum state’’ understood as the cyclic vector of the GNS construction.

In the discussion of various vacuum phenomena (e.g. spontaneous emission) we will assume for simplicity that all the oscillators are ‘‘embedded’’ in identical vacua i.e. the multi-oscillator vacuum is of the form

$$\begin{aligned} |\Psi\rangle &= \sqrt{p_1} |\phi\rangle \\ &\oplus \sqrt{p_2} |\phi\rangle |\phi\rangle \\ &\oplus \sqrt{p_3} |\phi\rangle |\phi\rangle |\phi\rangle \\ &\oplus \dots \end{aligned} \quad (89)$$

The average energy of the free-field vacuum state is therefore

$$\overline{\mathcal{H}} = \langle \Psi | \mathcal{H} | \Psi \rangle = \sum_{n=1}^{\infty} n p_n \langle \phi | H | \phi \rangle = \overline{n} \overline{H} \quad (90)$$

where \overline{n} and \overline{H} are, respectively, the average number of oscillators and the average energy of a single oscillator. Again no problem with infinite vacuum energy is found. Obviously, one can contemplate also other vacua, say, in entangled or mixed states.

C. Multi-oscillator vs multi-photon states

The coherent states we have introduced at the one-oscillator level involve superpositions of different excited states. We know that in the traditional approach the transition between two such states, say,

$$|s, \vec{\kappa}, 2\rangle \rightarrow |s, \vec{\kappa}, 0\rangle \quad (91)$$

is interpreted as an absorption (by some system) of two photons. In the new formulation the problem is more complicated since the “2-photon” absorption may be represented also by

$$|s, \vec{\kappa}, 1\rangle |s, \vec{\kappa}, 1\rangle \rightarrow |s, \vec{\kappa}, 0\rangle |s, \vec{\kappa}, 0\rangle. \quad (92)$$

The two types of transitions do not represent the same process and the two final states are physically distinguishable since their energies are different. Indeed,

$$\mathcal{H} |s, \vec{\kappa}, 0\rangle = H |s, \vec{\kappa}, 0\rangle = \frac{1}{2} \hbar \omega_{\vec{\kappa}} |s, \vec{\kappa}, 0\rangle \quad (93)$$

whereas

$$\begin{aligned} \mathcal{H} |s, \vec{\kappa}, 0\rangle |s, \vec{\kappa}, 0\rangle &= (H \otimes \mathbf{1} + \mathbf{1} \otimes H) |s, \vec{\kappa}, 0\rangle |s, \vec{\kappa}, 0\rangle \\ &= \hbar \omega_{\vec{\kappa}} |s, \vec{\kappa}, 0\rangle |s, \vec{\kappa}, 0\rangle. \end{aligned} \quad (94)$$

The notion of a 2-photon state becomes therefore somewhat ambiguous. To make it more precise one has to formulate a photodetection theory within the new framework. In what follows we shall try to avoid the use of the word “photon” and will talk about “light quanta” and “multi-oscillator” (or n -oscillator) and “higher-excited” (or n -th excited) states of light.

The 2-oscillator states

$$|\Psi_{\pm}\rangle = \sum_{\vec{\kappa}_1, \vec{\kappa}_2} \Psi_{\vec{\kappa}_1, \vec{\kappa}_2, n}^{(2)} \left(|+, \vec{\kappa}_1, n\rangle |-, \vec{\kappa}_2, n\rangle \pm |-, \vec{\kappa}_1, n\rangle |+, \vec{\kappa}_2, n\rangle \right), \quad (95)$$

satisfying

$$\Psi_{\vec{\kappa}_1, \vec{\kappa}_2, n}^{(2)} = \pm \Psi_{\vec{\kappa}_2, \vec{\kappa}_1, n}^{(2)} \quad (96)$$

are (for any $n > 0$) perfectly justified generalizations of the standard 2-photon maximally entangled state. We shall later see that although such “higher excited photons” (i.e. $n > 1$) are in principle possible, they are not produced in a two-photon spontaneous emission (at least up to second-order perturbative effects). The technical reason for this is the same as in the ordinary formalism and is related to properties of the annihilation operator a .

VII. SPONTANEOUS EMISSION OF A “SINGLE PHOTON”

In this section we shall again consider the spontaneous emission of light within the two-level-atom approximation. The example illustrates some peculiarities of the multi-oscillator formulation.

Denote by \mathcal{H}_F the multi-oscillator Hamiltonian of the free field we have discussed in the previous two sections. The dipole and RWA Hamiltonian of the 2-level atom interacting with quantized electromagnetic field is now

$$\mathcal{H} = \frac{1}{2}\hbar\omega_0\sigma_3 + \mathcal{H}_F + \hbar\omega_0d \sum_{s,\vec{\kappa}_\lambda} \left(g_{s,\vec{\kappa}_\lambda} \mathbf{a}_{s,\vec{\kappa}_\lambda} \sigma_+ + g_{s,\vec{\kappa}_\lambda}^* \mathbf{a}_{s,\vec{\kappa}_\lambda}^\dagger \sigma_- \right). \quad (97)$$

Similarly to the one-oscillator case one has

$$e^{i\mathcal{H}_F t/\hbar} \mathbf{a}_{s,\vec{\kappa}} e^{-i\mathcal{H}_F t/\hbar} = e^{-i\omega_{\vec{\kappa}} t} \mathbf{a}_{s,\vec{\kappa}} \quad (98)$$

and therefore the interaction-picture Hamiltonian is

$$\mathcal{H}_I = \hbar\omega_0d \sum_{s,\vec{\kappa}_\lambda} \left(g_{s,\vec{\kappa}_\lambda} e^{i(\omega_0 - \omega_\lambda)t} \mathbf{a}_{s,\vec{\kappa}_\lambda} \sigma_+ + g_{s,\vec{\kappa}_\lambda}^* e^{-i(\omega_0 - \omega_\lambda)t} \mathbf{a}_{s,\vec{\kappa}_\lambda}^\dagger \sigma_- \right). \quad (99)$$

The first peculiarity we encounter is the fact that the Hamiltonian is *block diagonal* with respect to \oplus and therefore does not have nonvanishing matrix elements between spaces corresponding to different numbers of oscillators. As a result the interaction cannot change the number of oscillators, a property of crucial importance for statistical properties of light as we shall see in the context of the Planck blackbody radiation law.

Consider the initial state

$$\begin{aligned} |\Psi(0)\rangle &= \sqrt{p_1}|+\rangle \sum_{s,\vec{\kappa}} \phi_{s,\vec{\kappa}} |s,\vec{\kappa},0\rangle \\ &\oplus \sqrt{p_2}|+\rangle \sum_{s_1,\vec{\kappa}_1} \sum_{s_2,\vec{\kappa}_2} \phi_{s_1,\vec{\kappa}_1} \phi_{s_2,\vec{\kappa}_2} |s_1,\vec{\kappa}_1,0\rangle |s_2,\vec{\kappa}_2,0\rangle \\ &\oplus \sqrt{p_3}|+\rangle \sum_{s_1,\vec{\kappa}_1} \sum_{s_2,\vec{\kappa}_2} \sum_{s_3,\vec{\kappa}_3} \phi_{s_1,\vec{\kappa}_1} \phi_{s_2,\vec{\kappa}_2} \phi_{s_3,\vec{\kappa}_3} |s_1,\vec{\kappa}_1,0\rangle |s_2,\vec{\kappa}_2,0\rangle |s_3,\vec{\kappa}_3,0\rangle \end{aligned} \quad (100)$$

The first-order perturbation theory yields (it is instructive to keep again the constants c_n arbitrary)

$$\begin{aligned} |\Psi(t)\rangle &= |\Psi(0)\rangle + \omega_0 d c_1 \sqrt{p_1} \sum_{s,\vec{\kappa}} \frac{e^{-i(\omega_0 - \omega_{\vec{\kappa}})t} - 1}{\omega_0 - \omega_{\vec{\kappa}}} g_{s,\vec{\kappa}}^* \phi_{s,\vec{\kappa}} |-\rangle |s,\vec{\kappa},1\rangle \\ &\oplus \omega_0 d c_2 \sqrt{p_2} |-\rangle \left(\sum_{s_1,\vec{\kappa}_1} \frac{e^{-i(\omega_0 - \omega_{\vec{\kappa}_1})t} - 1}{\omega_0 - \omega_{\vec{\kappa}_1}} g_{s_1,\vec{\kappa}_1}^* \phi_{s_1,\vec{\kappa}_1} |s_1,\vec{\kappa}_1,1\rangle |\phi\rangle + \sum_{s_2,\vec{\kappa}_2} \frac{e^{-i(\omega_0 - \omega_{\vec{\kappa}_2})t} - 1}{\omega_0 - \omega_{\vec{\kappa}_2}} g_{s_2,\vec{\kappa}_2}^* \phi_{s_2,\vec{\kappa}_2} |\phi\rangle |s_2,\vec{\kappa}_2,1\rangle \right) \\ &\oplus \dots \end{aligned} \quad (101)$$

As we can see, the “single-photon” emission can be realized in an infinite number of different ways. In the 1-oscillator subspace the oscillator simply gets excited to the 1-st excited state. The probability amplitude for this process is proportional to the probability amplitude that the field is found in a 1-oscillator state. In the 2-oscillator subspace there are two possibilities: Either the first oscillator gets excited and the second one remains in the ground state, or the other way around. The probability amplitude for this process is proportional to the probability amplitude that the field is found in a 2-oscillator state. And so on.

Repeating the argument given for a single-oscillator description, assuming the isotropy and polarization-independence of the vacuum mode density, we arrive at the spontaneous emission rate of the form

$$P = F(\omega_0) P_{\text{old}} \sum_{n=1}^{\infty} n c_n^2 p_n \quad (102)$$

where $F(\omega_{\vec{\kappa}}) = |\phi_{s,\vec{\kappa}}|^2$. As we can see the choice $c_n = 1/\sqrt{n}$ plays again a special role since then

$$P = F(\omega_0) P_{\text{old}}, \quad (103)$$

that is, the result is the same as in the single-oscillator description.

VIII. SPONTANEOUS EMISSION OF “TWO PHOTONS”

In what follows we will start with the Hamiltonian $H = H_0 + V$, where

$$H_0 = H_A + \mathcal{H}_F \quad (104)$$

$$V = -\frac{e}{m} \vec{\mathcal{A}}(\vec{x}) \cdot \vec{p} \\ = -\frac{e}{m} \sum_{s, \vec{\kappa}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}} V}} \left(\mathbf{a}_{s, \vec{\kappa}} e^{i\vec{\kappa} \cdot \vec{x}} \vec{e}_{s, \vec{\kappa}} \cdot \vec{p} + \mathbf{a}_{s, \vec{\kappa}}^\dagger e^{-i\vec{\kappa} \cdot \vec{x}} \vec{e}_{s, \vec{\kappa}}^* \cdot \vec{p} \right). \quad (105)$$

H_A is the full (i.e. infinite-level) Hamiltonian describing an atom and \mathcal{H}_F is the free-field Hamiltonian discussed in Sec. V and obtained by the multi-oscillator extension of the one-oscillator Hamiltonian introduced in Sec. III. To simplify notation we shall denote the sum and the integral over, respectively, the discrete and the continuous parts of the spectrum of H_A by the sum \sum_c . We are not making the rotating wave approximation. In the dipole approximation we set $\vec{x} = 0$. We shall also keep the constants c_n arbitrary.

A. Two different light-quanta in 2-oscillator subspace

In this subsection we will use the second-order perturbation theory to compute the amplitude

$$\langle b | \langle s_1, \vec{\kappa}_1, 1 | \langle s_2, \vec{\kappa}_2, 1 | U(t_f, t_i) | a \rangle | \Psi \rangle \quad (106)$$

where the states $|s_k, \vec{\kappa}_k, 1\rangle$, $k = 1, 2$, are orthogonal, $U(t_f, t_i)$ is the evolution operator mapping the initial state at time t_i into the final state at time t_f , $|\Psi\rangle$ is a vacuum state (89), and $|a\rangle$, $|b\rangle$ are two bound states of the atomic Hamiltonian H_A .

The fact that the interaction term does not change the number of oscillators reduces the above amplitude to its 2-oscillator counterpart

$$\sqrt{p_2} \langle b | \langle s_1, \vec{\kappa}_1, 1 | \langle s_2, \vec{\kappa}_2, 1 | U(t_f, t_i) | a \rangle | \phi \rangle | \phi \rangle \quad (107)$$

Using the standard perturbative techniques we obtain the second-order approximation [9] (see Appendix XI C)

$$\langle b | \langle s_1, \vec{\kappa}_1, 1 | \langle s_2, \vec{\kappa}_2, 1 | U^{(2)}(t_f, t_i) | a \rangle | \phi \rangle | \phi \rangle \\ = -c_2^2 \frac{2\pi i e^2}{m^2} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_2} V}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_1} V}} \phi_{s_1, \vec{\kappa}_1} \phi_{s_2, \vec{\kappa}_2} \sum_c \frac{(\vec{e}_{s_2, \vec{\kappa}_2}^* \cdot \vec{p}_{bc})(\vec{e}_{s_1, \vec{\kappa}_1}^* \cdot \vec{p}_{ca})}{E_{a, \vec{\kappa}_1, 0, \vec{\kappa}_2, 0} - E_{c, \vec{\kappa}_1, 1, \vec{\kappa}_2, 0} + i0_+} \delta^{(T)}(E_{a, \vec{\kappa}_1, \vec{\kappa}_2} - E_{b, \vec{\kappa}_1, 1, \vec{\kappa}_2, 1}) \\ - c_2^2 \frac{2\pi i e^2}{m^2} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_2} V}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_1} V}} \phi_{s_1, \vec{\kappa}_1} \phi_{s_2, \vec{\kappa}_2} \sum_c \frac{(\vec{e}_{s_1, \vec{\kappa}_1}^* \cdot \vec{p}_{bc})(\vec{e}_{s_2, \vec{\kappa}_2}^* \cdot \vec{p}_{ca})}{E_{a, \vec{\kappa}_1, 0, \vec{\kappa}_2, 0} - E_{c, \vec{\kappa}_1, 0, \vec{\kappa}_2, 1} + i0_+} \delta^{(T)}(E_{a, \vec{\kappa}_1, \vec{\kappa}_2} - E_{b, \vec{\kappa}_1, 1, \vec{\kappa}_2, 1}) \quad (108)$$

where $\delta^{(T)}(E - E') = [\pi(E - E')]^{-1} \sin((E - E')T/2\hbar)$, $\vec{p}_{bc} = \langle b | \vec{p} | c \rangle$, and $\vec{p}_{ca} = \langle c | \vec{p} | a \rangle$.

The energies occurring in the above expression are (ground-state energies are *not* removed!)

$$E_{a, \vec{\kappa}_1, 0, \vec{\kappa}_2, 0} = E_a + \frac{1}{2} \hbar \omega_{\vec{\kappa}_1} + \frac{1}{2} \hbar \omega_{\vec{\kappa}_2} \quad (109)$$

$$E_{b, \vec{\kappa}_1, 1, \vec{\kappa}_2, 1} = E_b + \left(1 + \frac{1}{2}\right) \hbar \omega_{\vec{\kappa}_1} + \left(1 + \frac{1}{2}\right) \hbar \omega_{\vec{\kappa}_2} \quad (110)$$

$$E_{c, \vec{\kappa}_1, 1, \vec{\kappa}_2, 0} = E_c + \left(1 + \frac{1}{2}\right) \hbar \omega_{\vec{\kappa}_1} + \frac{1}{2} \hbar \omega_{\vec{\kappa}_2} \quad (111)$$

$$E_{c, \vec{\kappa}_1, 0, \vec{\kappa}_2, 1} = E_c + \frac{1}{2} \hbar \omega_{\vec{\kappa}_1} + \left(1 + \frac{1}{2}\right) \hbar \omega_{\vec{\kappa}_2} \quad (112)$$

The net result is the following

$$\langle b | \langle s_1, \vec{\kappa}_1, 1 | \langle s_2, \vec{\kappa}_2, 1 | U(t_f, t_i) | a \rangle | \Psi \rangle \\ \approx -c_2^2 \sqrt{p_2} \frac{2\pi i e^2}{m^2} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_2} V}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_1} V}} \phi_{s_1, \vec{\kappa}_1} \phi_{s_2, \vec{\kappa}_2} \sum_c \frac{(\vec{e}_{s_2, \vec{\kappa}_2}^* \cdot \vec{p}_{bc})(\vec{e}_{s_1, \vec{\kappa}_1}^* \cdot \vec{p}_{ca})}{E_a - E_c - \hbar \omega_{\vec{\kappa}_1} + i0_+} \delta^{(T)}(E_a - E_b - \hbar \omega_{\vec{\kappa}_1} - \hbar \omega_{\vec{\kappa}_2}) \\ - c_2^2 \sqrt{p_2} \frac{2\pi i e^2}{m^2} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_2} V}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_1} V}} \phi_{s_1, \vec{\kappa}_1} \phi_{s_2, \vec{\kappa}_2} \sum_c \frac{(\vec{e}_{s_1, \vec{\kappa}_1}^* \cdot \vec{p}_{bc})(\vec{e}_{s_2, \vec{\kappa}_2}^* \cdot \vec{p}_{ca})}{E_a - E_c - \hbar \omega_{\vec{\kappa}_2} + i0_+} \delta^{(T)}(E_a - E_b - \hbar \omega_{\vec{\kappa}_1} - \hbar \omega_{\vec{\kappa}_2})$$

Let us note that the amplitude is symmetric with respect to permutation of states of the two oscillators:

$$\langle b | \langle s_1, \vec{\kappa}_1, 1 | \langle s_2, \vec{\kappa}_2, 1 | U(t_f, t_i) | a \rangle | \Psi \rangle = \langle b | \langle s_2, \vec{\kappa}_2, 1 | \langle s_1, \vec{\kappa}_1, 1 | U(t_f, t_i) | a \rangle | \Psi \rangle$$

B. Two different light-quanta in 3-oscillator subspace

Consider the amplitudes

$$\langle b|\langle s_1, \vec{\kappa}_1, 1|\langle s_2, \vec{\kappa}_2, 1|\langle \phi|U(t_f, t_i)|a\rangle|\Psi\rangle \quad (113)$$

$$\langle b|\langle s_1, \vec{\kappa}_1, 1|\langle \phi|\langle s_2, \vec{\kappa}_2, 1|U(t_f, t_i)|a\rangle|\Psi\rangle \quad (114)$$

$$\langle b|\langle \phi|\langle s_1, \vec{\kappa}_1, 1|\langle s_2, \vec{\kappa}_2, 1|U(t_f, t_i)|a\rangle|\Psi\rangle \quad (115)$$

In the framework we propose it is necessary to include the contributions of this type arising from all the possible numbers of oscillators.

It is again sufficient to restrict the analysis to

$$\sqrt{p_3}\langle b|\langle s_1, \vec{\kappa}_1, 1|\langle s_2, \vec{\kappa}_2, 1|\langle \phi|U(t_f, t_i)|a\rangle|\phi\rangle|\phi\rangle|\phi\rangle \quad (116)$$

and similarly with the other two amplitudes. In second-order perturbation theory (see Appendix XIX)

$$c_3^{-2}\langle b|\langle s_1, \vec{\kappa}_1, 1|\langle s_2, \vec{\kappa}_2, 1|\langle \phi|U^{(2)}(t_f, t_i)|a\rangle|\phi\rangle|\phi\rangle|\phi\rangle = c_2^{-2}\langle b|\langle s_1, \vec{\kappa}_1, 1|\langle s_2, \vec{\kappa}_2, 1|U^{(2)}(t_f, t_i)|a\rangle|\phi\rangle|\phi\rangle. \quad (117)$$

The result is therefore essentially identical to the one obtained for the 2-oscillator subspace.

A closer look at the derivation of the 3-oscillator contribution shows that (i) exactly the same will happen for any number of oscillators and (ii) the second-order amplitude describes an emission of at most two quanta.

C. Comparison with the standard formalism

It is instructive to compare the result we have obtained with the second-order calculation performed by means of the ordinary quantum optics formalism. Let

$$|s_1, \vec{\kappa}_1; s_2, \vec{\kappa}_2\rangle = |s_2, \vec{\kappa}_2; s_1, \vec{\kappa}_1\rangle = \mathbf{a}_{s_1, \vec{\kappa}_1}^\dagger \mathbf{a}_{s_2, \vec{\kappa}_2}^\dagger |0\rangle \quad (118)$$

be the two-photon state of the standard formalism, $\mathbf{a}_{s, \vec{\kappa}}^\dagger$ the standard creation operator and $|0\rangle$ the vacuum state. Then

$$\begin{aligned} & \langle b|\langle s_1, \vec{\kappa}_1; s_2, \vec{\kappa}_2|U^{(2)}(t_f, t_i)|a\rangle|0\rangle \\ &= -\frac{2\pi i e^2}{m^2} \sum_c \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_1} V}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_2} V}} (\vec{e}_{s_1, \vec{\kappa}_1}^* \cdot \vec{p}_{bc}) (\vec{e}_{s_2, \vec{\kappa}_2}^* \cdot \vec{p}_{ca}) \frac{\delta^{(T)}(E_a - E_b - \hbar\omega_{\vec{\kappa}_1} - \hbar\omega_{\vec{\kappa}_2})}{E_a - E_c - \hbar\omega_{\vec{\kappa}_2} + i0_+} \\ & - \frac{2\pi i e^2}{m^2} \sum_c \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_2} V}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}_1} V}} (\vec{e}_{s_2, \vec{\kappa}_2}^* \cdot \vec{p}_{bc}) (\vec{e}_{s_1, \vec{\kappa}_1}^* \cdot \vec{p}_{ca}) \frac{\delta^{(T)}(E_a - E_b - \hbar\omega_{\vec{\kappa}_1} - \hbar\omega_{\vec{\kappa}_2})}{E_a - E_c - \hbar\omega_{\vec{\kappa}_1} + i0_+}. \end{aligned}$$

This is precisely the same expression that occurs in the modified amplitudes.

D. Probability of spontaneous emission of two quanta

In the subspace corresponding to n oscillators the ‘‘two-photon’’ emission can take place in

$$\binom{n}{2} = \frac{n(n-1)}{2} \quad (119)$$

different ways. Taking into account probability amplitudes associated with all the n -oscillator subspaces, $n > 1$, and the fact that the two quanta can be emitted in two different orders, we obtain

$$p(s_1, \vec{\kappa}_1, s_2, \vec{\kappa}_2) = 2 \sum_{n=2}^{\infty} \frac{n(n-1)}{2} c_n^A p_n |\phi_{s_1, \vec{\kappa}_1}|^2 |\phi_{s_2, \vec{\kappa}_2}|^2 p(s_1, \vec{\kappa}_1, s_2, \vec{\kappa}_2)_{\text{old}} \quad (120)$$

where $p(s_1, \vec{\kappa}_1, s_2, \vec{\kappa}_2)_{\text{old}}$ is the standard result obtained by means of ordinary quantum optics. Taking, as before, $c_n = 1/\sqrt{n}$ we find

$$p(s_1, \vec{\kappa}_1, s_2, \vec{\kappa}_2) = \left(1 - \sum_{n=1}^{\infty} \frac{p_n}{n}\right) |\phi_{s_1, \vec{\kappa}_1}|^2 |\phi_{s_2, \vec{\kappa}_2}|^2 p(s_1, \vec{\kappa}_1, s_2, \vec{\kappa}_2)_{\text{old}} \quad (121)$$

Under such assumptions the angular distribution of the two-photon emission is the same as in the standard theory. The probability of the 2-photon spontaneous emission is thus a product of four terms. It may be difficult to distinguish between $F(\omega_{\vec{\kappa}_1})F(\omega_{\vec{\kappa}_2})(1 - \langle \frac{1}{n} \rangle)$ and analogous factors arising in real experiments from detector inefficiency. The above result may have therefore nontrivial implications for the problem of testing quantum mechanics versus local hidden-variables theories and is very closely related to the so-called detector inefficiency loophole in Bell's theorem (cf. [10,11]). The reason is that the presence of $F(\omega_{\vec{\kappa}_1})F(\omega_{\vec{\kappa}_2})(1 - \langle \frac{1}{n} \rangle)$ will necessarily lower two-photon coincidence rates, whereas it is known that in order to violate the Bell inequality the rates must exceed certain thresholds. The problem is worth further studies.

IX. BLACKBODY RADIATION

One of the possible tests of the new formalism is the problem of blackbody radiation. Planck's famous formula [12]

$$\varrho(\omega) = \frac{\hbar}{\pi^2 c^3} \frac{\omega^3}{e^{\beta \hbar \omega} - 1} = \frac{\hbar}{\pi^2 c^3} \omega^3 \overline{N}_\omega, \quad (122)$$

where \overline{N}_ω is the average number of excitations of an oscillator in inverse temperature β , is one of the first great successes of quantum radiation theory and marks the beginning of quantum mechanics. Contemporary measurements of $\varrho(\omega)$ [13,14] performed by means of COBE (Cosmic Background Explorer) are in a very good agreement with the Planck law. The data have been carefully analyzed in the context of nonextensive statistics [15,16] in search of possible deviations from extensivity. The result that comes out systematically is $|q - 1| < 10^{-4}$ where q is the Tsallis parameter. The case $q = 1$ corresponds to the exact Planck formula. If there are any corrections whatever, they must be quite small.

The standard derivation of the formula consists basically of two steps. First, one counts the number of different wave vectors \vec{k} such that $c|\vec{k}| \in [\omega, \omega + \Delta\omega]$. Second, one associates with each such a vector an oscillator and counts the average number of its excitations assuming the Boltzmann-Gibbs probability distribution at temperature T and chemical potential $\mu = 0$. The latter assumption is justified by the fact that the number of excitations of the electromagnetic field is not conserved in atom-light interactions.

In the new model the situation is slightly different since there exists an additional conserved quantum number: The number of *oscillators*. As we have seen in previous calculations the Hamiltonian is block-diagonal with respect to \oplus but changes the number of excitations in each N -oscillator subspace of the direct sum. The state vectors at the multi-oscillator level are symmetric with respect to permutations of the oscillators and therefore the oscillators themselves have to be regarded as bosons whose number is conserved and their chemical potential is $\mu \neq 0$. However, their excitations should be regarded as bosons with vanishing chemical potential.

The energy eigenvalues

$$E_{m,n} = m\hbar\omega \left(n + \frac{1}{2} \right) \quad (123)$$

corresponding to the oscillator whose frequency is ω are parametrized by two natural numbers: m (the number of oscillators) and n (the number of excitations). Assuming the standard Boltzmann-Gibbs statistics we obtain the probabilities

$$p_{m,n} = Z^{-1} e^{-\beta[m\hbar\omega(n+\frac{1}{2})-m\mu]} \quad (124)$$

where

$$Z = \sum_{m=1}^{\infty} e^{\beta m(\mu + \hbar\omega/2)} \frac{e^{-\beta m\hbar\omega}}{1 - e^{-\beta m\hbar\omega}}. \quad (125)$$

The Lambert series [17]

$$\sum_{m=1}^{\infty} a_m \frac{x^m}{1 - x^m} \quad (126)$$

is convergent for any x if $\sum_{m=1}^{\infty} a_m$ is convergent. Otherwise (126) converges for exactly those x for which the power series $\sum_{m=1}^{\infty} a_m x^m$ does. In (125) $a_m = e^{\beta m(\mu + \hbar\omega/2)}$ and $\sum_{m=1}^{\infty} a_m < \infty$ if $\mu + \hbar\omega/2 < 0$. If $\mu + \hbar\omega/2 \geq 0$ we still have convergence of (125) as long as $\sum_{m=1}^{\infty} e^{-\beta m[\frac{1}{2}\hbar\omega - \mu]} < \infty$. The upper limit imposed on μ by the finiteness of Z is therefore $\frac{1}{2}\hbar\omega - \mu > 0$. In what follows we assume that μ is ω -independent and therefore $\mu \leq 0$.

The appropriate average number of excitations is

$$\bar{n}_\omega = Z^{-1} \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} mn e^{-\beta[m\hbar\omega(n+\frac{1}{2})-m\mu]} \quad (127)$$

and the Planck formula is replaced by

$$\varrho_{\text{new}}(\omega) = \frac{\hbar}{\pi^2 c^3} \omega^3 \bar{n}_\omega. \quad (128)$$

It is easy to show that $\varrho_{\text{new}}(\omega)$ tends to the Planck distribution with $\mu \rightarrow -\infty$. To see this consider a more general series

$$Z^{-1} \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} mn q_m e^{-\beta m\hbar\omega(n+\frac{1}{2})} \quad (129)$$

where Z is the normalization factor and $\sum_{m=1}^{\infty} q_m < \infty$. If $q_1 = 1$ and $q_m = 0$ for $m > 1$ then (129) is just the exact Planckian formula. Factoring out $e^{-\beta|\mu|}$ in both the numerator and the denominator of \bar{n}_ω we obtain $q_1 = 1$ and $q_m = e^{-\beta|\mu|(m-1)}$ for $m > 1$. For $|\mu| \rightarrow \infty$ all q_m , for $m > 1$, vanish and the limiting distribution is Planckian.

This proves that an experimental agreement with the ordinary Planck's $\varrho(\omega)$ cannot rule out our modification but can, at most, set a lower bound on an admissible value of $|\mu|$. However, assuming that μ has some finite and fixed value it should be in principle measurable. The plots show that the modifications become visible around $\mu \approx -3k_B T$. Assuming that the chemical potential is temperature independent, say $\mu = -k_B T_0$, we obtain a kind of critical temperature $T_{\text{critical}} \approx T_0/3$ above which the ratio $\mu/(k_B T)$ is small enough to make the modifications of the distribution observable. For $T < T_{\text{critical}}$ the distribution should be given by the Planck law; for $T > T_{\text{critical}}$ the distribution should approach the $\mu = 0$ distribution, i.e. this would be a Planck-type curve but with the maximum lowered and shifted towards higher energies.

Fig. 1 shows the plots of $\varrho_{\text{new}}(\omega)$ for $\mu = 0$ (lower dotted), $\mu = -0.8k_B T$ (upper dotted), and $\mu = -10k_B T$ (solid). The thick dashed curve is the Planck distribution. The curve obtained for $\mu = -10k_B T$ is indistinguishable from the Planck distribution. The plot does not change if one takes $\mu < -10k_B T$ and differences are not visible even if one plots the distributions in logarithmic scales (not shown here). This is a numerical proof that the distribution we have obtained on the basis of the modified quantization tends very quickly to the Planck one as $\mu \rightarrow -\infty$. It is instructive to compare the modification we have predicted with those arising from nonextensive statistics. The two thin dashed lines represent Tsallis distributions resulting from nonextensive formalism for $q = 0.95$ (lower) and $q = 1.05$ (upper). The modifications we have derived are therefore qualitatively different from those resulting from Tsallis statistics.




FIG. 1. $\varrho_{\text{new}}(\omega)$ for $\mu = 0$ (lower dotted), $\mu = -0.8k_B T$ (upper dotted), and $\mu = -10k_B T$ (solid). The energy range is $0.01k_B T < \hbar\omega < 10k_B T$. The thick dashed curve is the Planck distribution. The curve obtained for $\mu = -10k_B T$ is indistinguishable from the Planck distribution. The two thin dashed lines represent Tsallis distributions resulting from the Tsallis formalism for $q = 0.95$ (lower) and $q = 1.05$ (upper). Since $\varrho_{\text{new}} < \varrho$ at least in the neighborhood of the maximum, the new distribution has to be compared with $q < 1$ statistics. The curves are qualitatively different. In particular, all $q < 1$ distributions require an energy cut-off which does not occur for ϱ_{new} .




FIG. 2. $\varrho_{\text{new}}(\omega)$ for $-10k_B T \leq \mu \leq 0$. The cut through $\mu = -10k_B T$ is practically indistinguishable from the Planck distribution.




FIG. 3. Contour plot of $\varrho_{\text{new}}(\omega)$ for $-10k_B T \leq \mu \leq 0$. The fast convergence to Planckian $\varrho(\omega)$ (as $\mu \rightarrow -\infty$) is clearly seen.

X. CONCLUSIONS

“A theory that is as spectacularly successful as quantum electrodynamics has to be more or less correct, although we may not be formulating it in just the right way” [7]. The above quotation from Weinberg could serve as a motto opening our paper. The main idea we have tried to advocate was that the standard canonical quantization procedure is, in certain sense, *too classical* to be good.

The reasons for such a choice of quantization could be both historical and sociological and may be rooted in the fact that the idea of quantizing the field was formulated before the real development of modern quantum mechanics. In oscillations of a simple pendulum it may be justified to treat ω as an external parameter defining the system (via, say, the length of the pendulum). But oscillations of the electromagnetic field do not seem to have such a “mechanical” origin and it is more natural to think of the spectrum of frequencies as eigenvalues of some Hamiltonian. That is exactly what happens with other quantum wave equations.

We have defined the quantum electromagnetic field as an oscillator that can exist in a superposition of different frequencies (or, rather, wave vectors). This should not be confused with the classical superpositions of frequencies created by, say, a guitar string. The superpositions we have in mind disappear at the classical level.

Once one accepts this viewpoint it becomes clear how to quantize the field at the level of a single oscillator. We do not need many oscillators to perform the field quantization. But there is no reason to believe that all the possible fields can be described by the same single oscillator. And even more: We know that the structure of the one-oscillator Hilbert space is not rich enough to describe multi-particle entangled states and there is no doubt that such states are physical. The next step, performed already *after* the quantization, is to consider fields consisting of 1, 2, 3 and more oscillators, and even existing in superpositions of different numbers of them. The resulting structure is analogous to the Fock space but so the procedure can be (although somewhat misleadingly) referred to as “second quantization”. What is essential we do not need the vacuum state understood as the cyclic vector of the GNS construction. In the new framework such an object seems rather artificial.

On the other hand, there exist vacuum *states*. These are all the states describing ground states of the oscillators. They correspond to concrete *finite* average values of energy. A general vacuum state is therefore a superposition of different eigenstates of a free Hamiltonian and is not, in itself, an eigenstate of the Hamiltonian.

No further assumptions are made. The system is described by laws of ordinary quantum mechanics so that to compute concrete problems we can use standard methods. Perturbation theory leads to structures we know from the standard Feynmann diagrams. The blackbody radiation is calculated by means of the standard Boltzmann-Gibbs statistics.

Let us close these remarks with another quotation: “Present quantum electrodynamics contains many very important ‘elements of truth’, but also some clear ‘elements of nonsense’. Because of the divergences and ambiguities, there is general agreement that a rather deep modification of the theory is needed, but in some forty years of theoretical work, nobody has seen how to disentangle the truth from the nonsense. In such a situation, one needs more experimental evidence, but during that same forty years we have found no clues from the laboratory as to what specific features of QED might be modified. Even worse, in the absence of any alternative theory whose predictions differ from those of QED in known ways, we have no criterion telling us *which* experiments would be relevant ones to try.

It seems useful, then, to examine the various disturbing features of QED, which give rise to mathematical or conceptual difficulties, to ask whether present empirical evidence demands their presence, and to explore the consequences of the modified (although perhaps rather crude and incomplete) theories in which these features are removed. Any difference between the predictions of QED and some alternative theory, corresponds to an experiment which might distinguish between them; if it appears untried but feasible, then we have the opportunity to subject QED to a new test in which we know just what to look for, and which we would be very unlikely to think of without the alternative theory. For this purpose, the alternative theory need not be worked out as completely as QED; it is sufficient if we know in what way their predictions will differ in the area of interest. Nor does the alternative theory need to be free of defects in all other respects; for if experiment should show that it contains just a single ‘element of truth’ that is *not* in QED, then the alternative theory will have served its purpose; we would have the long-missing clue showing in what way QED must be modified, and electrodynamics (and, I suspect, much more of theoretical physics along with it) could get moving again “ [3].

ACKNOWLEDGMENTS

This work was done partly during my stays in Arnold Sommerfeld Institute in Clausthal. I gratefully acknowledge a support from the Alexander von Humboldt Foundation. I’m indebted to Prof. Iwo Białynicki-Birula, Robert Alicki, Jan Naudts, and Wolfgang Luecke for critical comments, and Paweł Syty for a stimulating discussion on small ω ’s.

XI. APPENDIX: TECHNICAL DIFFERENCES AND SIMILARITIES WITH RESPECT TO THE STANDARD FORMALISM

A. Proof of Eq. (11)

This calculation is elementary but very important, so we give it explicitly:

$$a_{\omega_k}(t) = e^{iHt/\hbar} a_{\omega_k} e^{-iHt/\hbar} \quad (130)$$

$$= e^{i\Omega \otimes (a^\dagger a + \frac{1}{2}\mathbf{1})t} |\omega_k\rangle \langle \omega_k| \otimes a e^{-i\Omega \otimes (a^\dagger a + \frac{1}{2}\mathbf{1})t} \quad (131)$$

$$= |\omega_k\rangle \langle \omega_k| \otimes e^{i\omega_k (a^\dagger a + \frac{1}{2}\mathbf{1})t} a e^{-i\omega_k (a^\dagger a + \frac{1}{2}\mathbf{1})t} \quad (132)$$

$$= |\omega_k\rangle \langle \omega_k| \otimes e^{-i\omega_k t} a = e^{-i\omega_k t} a_{\omega_k}(0) \quad (133)$$

B. Energy-momentum operators for free fields: 1-oscillator formulas

To see how (36) and (37) arise let us first note that

$$a_{s_1, \vec{\kappa}_1} a_{s_2, \vec{\kappa}_2} = 0 \quad (134)$$

$$a_{s_1, \vec{\kappa}_1} a_{s_2, \vec{\kappa}_2}^\dagger = 0 \quad (135)$$

$$a_{s_1, \vec{\kappa}_1}^\dagger a_{s_2, \vec{\kappa}_2}^\dagger = 0 \quad (136)$$

unless $s_1 = s_2$ and $\vec{\kappa}_1 = \vec{\kappa}_2$ [cf. Eqs. (27) and (28)]. The terms involving $(a_{s, \vec{\kappa}})^2$ and $(a_{s, \vec{\kappa}}^\dagger)^2$ disappear due to $\vec{e}_{s, \vec{\kappa}} \cdot \vec{e}_{s, \vec{\kappa}} = 0$ and its complex conjugate. As a result

$$\begin{aligned} \frac{1}{2} \left(\hat{\vec{E}}(t, \vec{x}) \cdot \hat{\vec{E}}(t, \vec{x}) + \hat{\vec{B}}(t, \vec{x}) \cdot \hat{\vec{B}}(t, \vec{x}) \right) &= e^{iP \cdot x / \hbar} \sum_{s, \kappa_\lambda} \frac{\hbar \omega_\lambda}{2V} |s, \kappa_\lambda\rangle \langle s, \kappa_\lambda| \otimes (aa^\dagger + a^\dagger a) e^{-iP \cdot x / \hbar} \\ &= \sum_{s, \kappa_\lambda} \frac{\hbar \omega_\lambda}{2V} |s, \kappa_\lambda\rangle \langle s, \kappa_\lambda| \otimes (aa^\dagger + a^\dagger a) = H/V. \end{aligned}$$

To find the relation between the Pointing vector and linear momentum we first have to show that

$$\hat{\vec{E}} \times \hat{\vec{B}} = -\hat{\vec{B}} \times \hat{\vec{E}}. \quad (137)$$

The relevant formula is

$$[\hat{E}_\alpha, \hat{B}_\beta] = \sum_{s, \kappa_\lambda} i\hbar \omega_\lambda \frac{s}{2} \left(\delta_{\alpha\beta} - (n_{\vec{\kappa}_\lambda})_\alpha (n_{\vec{\kappa}_\lambda})_\beta \right) |s, \kappa_\lambda\rangle \langle s, \kappa_\lambda| \otimes \mathbf{1}. \quad (138)$$

The remaining calculations are similar to those for H .

C. Derivation of the ‘‘2-photon’’ amplitude: 2 oscillators

We employ the standard second-order time dependent perturbation theory and notation from [9].

$$\begin{aligned} &c_2^{-2} \langle b | \langle s_1, \vec{\kappa}_1, 1 | \langle s_2, \vec{\kappa}_2, 1 | U^{(2)}(t_f, t_i) | a \rangle | \phi \rangle | \phi \rangle \\ &= -\frac{2\pi i e^2}{m^2} \sum_{c, S_1, S_2, \vec{K}_1, \vec{K}_2, n_1, n_2} \sum_{r, \vec{k}, r', \vec{k}'} \phi_{r, \vec{k}} \phi_{r', \vec{k}'} \frac{\delta^{(T)}(E_{a, \vec{\kappa}_1, 0, \vec{\kappa}_2, 0} - E_{b, \vec{\kappa}_1, 1, \vec{\kappa}_2, 1})}{E_{a, \vec{k}, 0, \vec{k}, 0'} - E_{c, \vec{K}_1, n_1, \vec{K}_2, n_2} + i0_+} \\ &\quad \times \langle b | \langle s_1, \vec{\kappa}_1, 1 | \langle s_2, \vec{\kappa}_2, 1 | \sum_{s, \vec{\kappa}} \sqrt{\frac{\hbar}{2\omega_{\vec{\kappa}} V}} \left(\mathbf{a}_{s, \vec{\kappa}} \vec{e}_{s, \vec{\kappa}} \cdot \vec{p} + \mathbf{a}_{s, \vec{\kappa}}^\dagger \vec{e}_{s, \vec{\kappa}}^* \cdot \vec{p} \right) | c \rangle | S_1, \vec{K}_1, n_1 \rangle | S_2, \vec{K}_2, n_2 \rangle \end{aligned}$$

$$\begin{aligned}
& \times \langle c | \langle S_1, \vec{K}_1, n_1 | \langle S_2, \vec{K}_2, n_2 | \sum_{s', \vec{k}'} \sqrt{\frac{\hbar}{2\omega_{\vec{k}'}}} \left(\mathbf{a}_{s', \vec{k}'} \vec{e}_{s', \vec{k}'} \cdot \vec{p} + \mathbf{a}_{s', \vec{k}'}^\dagger \vec{e}_{s', \vec{k}'}^* \cdot \vec{p} \right) | a \rangle | r, \vec{k}, 0 \rangle | r', \vec{k}', 0 \rangle \\
& = -\frac{2\pi i e^2}{m^2} \sum_{c, S_1, S_2, \vec{K}_1, \vec{K}_2, n_1, n_2} \sum_{r, \vec{k}, r', \vec{k}'} \phi_{r, \vec{k}} \phi_{r', \vec{k}'} \frac{\delta^{(T)}(E_{a, \vec{k}_1, 0, \vec{k}_2, 0} - E_{b, \vec{k}_1, 1, \vec{k}_2, 1})}{E_{a, \vec{k}, 0, \vec{k}', 0} - E_{c, \vec{K}_1, n_1, \vec{K}_2, n_2} + i0_+} \\
& \times \langle s_1, \vec{k}_1, 1 | \langle s_2, \vec{k}_2, 1 | \sum_{s, \vec{k}} \sqrt{\frac{\hbar}{2\omega_{\vec{k}}}} \left((a_{s, \vec{k}} \otimes \mathbf{1} + \mathbf{1} \otimes a_{s, \vec{k}}) \vec{e}_{s, \vec{k}} \cdot \vec{p}_{bc} + (a_{s, \vec{k}}^\dagger \otimes \mathbf{1} + \mathbf{1} \otimes a_{s, \vec{k}}^\dagger) \vec{e}_{s, \vec{k}}^* \cdot \vec{p}_{bc} \right) \\
& \times | S_1, \vec{K}_1, n_1 \rangle | S_2, \vec{K}_2, n_2 \rangle \langle S_1, \vec{K}_1, n_1 | \langle S_2, \vec{K}_2, n_2 | \\
& \times \sum_{s', \vec{k}'} \sqrt{\frac{\hbar}{2\omega_{\vec{k}'}}} \left((a_{s', \vec{k}'} \otimes \mathbf{1} + \mathbf{1} \otimes a_{s', \vec{k}'}) \vec{e}_{s', \vec{k}'} \cdot \vec{p}_{ca} + (a_{s', \vec{k}'}^\dagger \otimes \mathbf{1} + \mathbf{1} \otimes a_{s', \vec{k}'}^\dagger) \vec{e}_{s', \vec{k}'}^* \cdot \vec{p}_{ca} \right) | r, \vec{k}, 0 \rangle | r', \vec{k}', 0 \rangle.
\end{aligned}$$

The block-diagonal property of the interaction Hamiltonian has been used twice. The remaining calculations are standard. It is remarkable that although the result we obtain is essentially the same as in the standard formalism, the technical reasons for this are completely different.

D. Derivation of the “2-photon” amplitude: 3 oscillators

Here we sketch the proof of the 3-oscillator amplitude. In second-order perturbation theory

$$\begin{aligned}
& c_3^{-2} \langle b | \langle s_1, \vec{k}_1, 1 | \langle s_2, \vec{k}_2, 1 | \langle \phi | U^{(2)}(t_f, t_i) | a \rangle | \phi \rangle | \phi \rangle \\
& = -\frac{2\pi i e^2}{m^2} \sum_{c, S_1, S_2, \vec{K}_1, \vec{K}_2, \vec{K}_3, n_1, n_2, n_3} \sum_{r_0, \vec{k}_0, r, \vec{k}, r', \vec{k}', r'', \vec{k}''} \phi_{r_0, \vec{k}_0}^* \phi_{r, \vec{k}} \phi_{r', \vec{k}'} \phi_{r'', \vec{k}''} \\
& \times \langle b | \langle s_1, \vec{k}_1, 1 | \langle s_2, \vec{k}_2, 1 | \langle r_0, \vec{k}_0, 0 | \sum_{s, \vec{k}} \sqrt{\frac{\hbar}{2\omega_{\vec{k}}}} \left(\mathbf{a}_{s, \vec{k}} \vec{e}_{s, \vec{k}} \cdot \vec{p} + \mathbf{a}_{s, \vec{k}}^\dagger \vec{e}_{s, \vec{k}}^* \cdot \vec{p} \right) | c \rangle | S_1, \vec{K}_1, n_1 \rangle | S_2, \vec{K}_2, n_2 \rangle | S_3, \vec{K}_3, n_3 \rangle \\
& \times \langle c | \langle S_1, \vec{K}_1, n_1 | \langle S_2, \vec{K}_2, n_2 | \langle S_3, \vec{K}_3, n_3 | \sum_{s', \vec{k}'} \sqrt{\frac{\hbar}{2\omega_{\vec{k}'}}} \left(\mathbf{a}_{s', \vec{k}'} \vec{e}_{s', \vec{k}'} \cdot \vec{p} + \mathbf{a}_{s', \vec{k}'}^\dagger \vec{e}_{s', \vec{k}'}^* \cdot \vec{p} \right) | a \rangle | r, \vec{k}, 0 \rangle | r', \vec{k}', 0 \rangle | r'', \vec{k}'', 0 \rangle \\
& \times \frac{\delta^{(T)}(E_{a, \vec{k}, 0, \vec{k}', 0, \vec{k}'', 0} - E_{b, \vec{k}_1, 1, \vec{k}_2, 1, \vec{k}_0, 0})}{E_{a, \vec{k}, 0, \vec{k}', 0, \vec{k}'', 0} - E_{c, \vec{K}_1, n_1, \vec{K}_2, n_2, \vec{K}_3, n_3} + i0_+} \\
& = -\frac{2\pi i e^2}{m^2} \sum_{c, S_1, S_2, \vec{K}_1, \vec{K}_2, \vec{K}_3, n_1, n_2, n_3} \sum_{r_0, \vec{k}_0, r, \vec{k}, r', \vec{k}', r'', \vec{k}''} \phi_{r_0, \vec{k}_0}^* \phi_{r, \vec{k}} \phi_{r', \vec{k}'} \phi_{r'', \vec{k}''} \\
& \times \langle b | \langle s_1, \vec{k}_1, 1 | \langle s_2, \vec{k}_2, 1 | \langle r_0, \vec{k}_0, 0 | \sum_{s, \vec{k}} \sqrt{\frac{\hbar}{2\omega_{\vec{k}}}} \left(\mathbf{a}_{s, \vec{k}} \vec{e}_{s, \vec{k}} \cdot \vec{p} + \mathbf{a}_{s, \vec{k}}^\dagger \vec{e}_{s, \vec{k}}^* \cdot \vec{p} \right) \\
& \times | c \rangle | S_1, \vec{K}_1, n_1 \rangle | S_2, \vec{K}_2, n_2 \rangle | S_3, \vec{K}_3, n_3 \rangle \langle c | \langle S_1, \vec{K}_1, n_1 | \langle S_2, \vec{K}_2, n_2 | \langle S_3, \vec{K}_3, n_3 | \\
& \times \sum_{s', \vec{k}'} \sqrt{\frac{\hbar}{2\omega_{\vec{k}'}}} \left(a_{s', \vec{k}'}^\dagger \otimes \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes a_{s', \vec{k}'}^\dagger \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1} \otimes a_{s', \vec{k}'}^\dagger \right) \vec{e}_{s', \vec{k}'}^* \cdot \vec{p} | a \rangle | r, \vec{k}, 0 \rangle | r', \vec{k}', 0 \rangle | r'', \vec{k}'', 0 \rangle \\
& \times \frac{\delta^{(T)}(E_{a, \vec{k}, 0, \vec{k}', 0, \vec{k}'', 0} - E_{b, \vec{k}_1, 1, \vec{k}_2, 1, \vec{k}_0, 0})}{E_{a, \vec{k}, 0, \vec{k}', 0, \vec{k}'', 0} - E_{c, \vec{K}_1, n_1, \vec{K}_2, n_2, \vec{K}_3, n_3} + i0_+}
\end{aligned}$$

The remaining part of the proof is standard. In the course of the computation one recognizes the elements known from standard Feynman diagrams, in particular the self-energy corrections due to emission and reabsorption of virtual photons. A general property of the perturbation series we find is its better convergence due to the presence of the vacuum amplitudes $\phi_{s, \vec{k}}$.

-
- [1] M. Born, W. Heisenberg, and P. Jordan, *Z. Phys.* **35**, 557 (1925)
- [2] E. Schrödinger, *Ann. Phys.* **79**, 361 (1926); *ibid.* **79**, 489 (1926).
- [3] E. T. Janes, in *Coherence and Quantum Optics III*, edited by L. Mandel and E. Wolf (Plenum, New York, 1973).
- [4] A. Casado, T. W. Marshall, R. Risco Delgado, and E. Santos, *Phys. Rev. A* **55**, 3879 (1997); *ibid.* *Phys. Rev. A* **56**, 2477 (1997).
- [5] P. A. M. Dirac, *Proc. Roy. Soc. A* **112**, 661 (1926); *ibid.* **114**, 243 (1927).
- [6] J. D. Cohn, astro-ph/9807128.
- [7] S. Weinberg, *Dreams of a final theory* (Vintage, New York, 1994).
- [8] H. Haken, *Light. Waves, photons, atoms* (Elsevier Science, Amsterdam, 1981).
- [9] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Atom-photon interactions: Basic processes and applications* (Wiley, New York, 1992).
- [10] S. Aerts, P. Kwiat, J-A. Larsson, and M. Żukowski, *Phys. Rev. Lett.* **83**, 2872 (1999).
- [11] N. Gisin, e-print quant-ph/9905062.
- [12] M. Planck, *Verh. Deut. Phys. Gessellsch.* **2**, 237 (1900).
- [13] L. Page and D. Wilkinson, *Rev. Mod. Phys.* **71**, S173 (1999).
- [14] G. F. Smoot, astro-ph/9902027.
- [15] C. Tsallis, F. C. Sa Bareto, and E. D. Loh, *Phys. Rev. E* **52**, 1447 (1995).
- [16] A. R. Plastino, A. Plastino, and H. Vucetich, *Phys. Lett. A* **207**, 42 (1995).
- [17] G. M. Fichtenholz, *Course of Differential and Integral Calculus*, vol. 2 (Fizmatgiz, Moscow, 1959) (in Russian).