

Covariant geometric quantization of non-relativistic Hamiltonian mechanics

GIOVANNI GIACHETTA^{†1}, LUIGI MANGIAROTTI^{†2} AND GENNADI SARDANASHVILY^{‡3}

[†] Department of Mathematics and Physics, University of Camerino, 62032 Camerino (MC), Italy

[‡] Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia

Abstract. We provide geometric quantization of the vertical cotangent bundle V^*Q equipped with the canonical Poisson structure. This is a momentum phase space of non-relativistic mechanics with the configuration bundle $Q \rightarrow \mathbf{R}$. The goal is the Schrödinger representation of V^*Q . We show that this quantization is equivalent to the fibrewise quantization of symplectic fibres of $V^*Q \rightarrow \mathbf{R}$ that makes the quantum algebra of non-relativistic mechanics an instantwise algebra. Quantization of the classical evolution equation defines a connection on this instantwise algebra, which provides quantum evolution in non-relativistic mechanics as a parallel transport along time.

1 Introduction

We study covariant geometric quantization of non-relativistic Hamiltonian mechanics subject to time-dependent transformations.

Its configuration space is a fibre bundle $Q \rightarrow \mathbf{R}$ equipped with bundle coordinates (t, q^k) , $k = 1, \dots, m$, where t is the Cartesian coordinate on the time axis \mathbf{R} with affine transition functions $t' = t + \text{const}$. Different trivializations $Q \cong \mathbf{R} \times M$ of Q correspond to different non-relativistic reference frames. In contrary to all the existent quantizations of non-relativistic mechanics (e.g., [14, 19]), we do not fix a trivialization of Q .

The momentum phase space of non-relativistic mechanics is the vertical cotangent bundle V^*Q of $Q \rightarrow \mathbf{R}$, endowed with the holonomic coordinates (t, q^k, p_k) . It is provided with the canonical Poisson structure

$$\{f, f'\}_V = \partial^k f \partial_k f' - \partial_k f \partial^k f', \quad f, f' \in C^\infty(V^*Q), \quad (1)$$

¹E-mail address: giachetta@campus.unicam.it

²E-mail address: mangiaro@camserv.unicam.it

³E-mail address: sard@grav.phys.msu.su

whose symplectic foliation coincides with the fibration $V^*Q \rightarrow \mathbf{R}$ [8, 12]. Given a trivialization

$$V^*Q \cong \mathbf{R} \times T^*M, \quad (2)$$

the Poisson manifold $(V^*Q, \{, \}_V)$ is isomorphic to the direct product of the Poisson manifold \mathbf{R} with the zero Poisson structure and the symplectic manifold T^*M . An important peculiarity of the Poisson structure (1) is that the Poisson algebra $C^\infty(V^*Q)$ of smooth real functions on V^*Q is a Lie algebra over the ring $C^\infty(\mathbf{R})$ of functions of time alone.

Our goal is the geometric quantization of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$, but it is not sufficient for quantization of non-relativistic mechanics.

The problem is that non-relativistic mechanics can not be described as a Poisson Hamiltonian system on the momentum phase space V^*Q . Indeed, a non-relativistic Hamiltonian \mathcal{H} is not an element of the Poisson algebra $C^\infty(V^*Q)$. Its definition involves the cotangent bundle T^*Q of Q . Coordinated by $(q^0 = t, q^k, p_0 = p, p_k)$, the cotangent bundle T^*Q plays the role of the homogeneous momentum phase space of non-relativistic mechanics. It is equipped with the canonical Liouville form $\Xi = p_\lambda dq^\lambda$, the symplectic form $\Omega = d\Xi$, and the corresponding Poisson bracket

$$\{f, g\}_T = \partial^\lambda f \partial_\lambda g - \partial_\lambda f \partial^\lambda g, \quad f, g \in C^\infty(T^*Q).$$

Due to the one-dimensional canonical fibration

$$\zeta : T^*Q \rightarrow V^*Q, \quad (3)$$

the cotangent bundle T^*Q provides the symplectic realization of the Poisson manifold V^*Q , i.e.,

$$\zeta^* \{f, g\}_V = \{ \zeta^* f, \zeta^* g \}_T$$

for all $f, g \in C^\infty(V^*Q)$. A Hamiltonian on V^*Q is defined as a global section

$$h : V^*Q \rightarrow T^*Q, \quad p \circ h = -\mathcal{H}(t, q^j, p_j), \quad (4)$$

of the one-dimensional affine bundle (3) [8, 12]. As a consequence (see Section 6), the evolution equation of non-relativistic mechanics is expressed into the Poisson bracket $\{, \}_T$ on T^*Q . It reads

$$\vartheta_{H^*}(\zeta^* f) = \{ \mathcal{H}^*, \zeta^* f \}_T, \quad (5)$$

where ϑ_{H^*} is the Hamiltonian vector field of the function

$$\mathcal{H}^* = \partial_t](\Xi - \zeta^* h^* \Xi)) = p + \mathcal{H} \quad (6)$$

on T^*Q .

Therefore, we need the compatible geometric quantizations both of the cotangent bundle T^*Q and the vertical cotangent bundle V^*Q such that the monomorphism

$$\zeta^* : (C^\infty(V^*Q), \{, \}_V) \rightarrow (C^\infty(T^*Q), \{, \}_T) \quad (7)$$

of the Poisson algebra on V^*Q to that on T^*Q is prolonged to a monomorphism of quantum algebras of V^*Q and T^*Q .

Recall that the geometric quantization procedure falls into three steps: prequantization, polarization and metaplectic correction (e.g., [3, 14, 19]). Given a symplectic manifold (Z, Ω) and the corresponding Poisson bracket $\{, \}$, prequantization associates to each element f of the Poisson algebra $C^\infty(Z)$ on Z a first order differential operator \widehat{f} in the space of sections of a complex line bundle C over Z such that the Dirac condition

$$[\widehat{f}, \widehat{f}'] = -i\{\widehat{f}, \widehat{f}'\} \quad (8)$$

holds. Polarization of a symplectic manifold (Z, Ω) is defined as a maximal involutive distribution $\mathbf{T} \subset TZ$ such that $\text{Orth}_\Omega \mathbf{T} = \mathbf{T}$, i.e.,

$$\Omega(\vartheta, v) = 0, \quad \forall \vartheta, v \in \mathbf{T}_z, \quad z \in Z. \quad (9)$$

Given the Lie algebra $\mathbf{T}(Z)$ of global sections of $\mathbf{T} \rightarrow Z$, let $\mathcal{A}_T \subset C^\infty(Z)$ denote the subalgebra of functions f whose Hamiltonian vector fields ϑ_f fulfill the condition

$$[\vartheta_f, \mathbf{T}(Z)] \subset \mathbf{T}(Z). \quad (10)$$

Elements of this subalgebra are only quantized. Metaplectic correction provides the pre-Hilbert space E_T where the quantum algebra \mathcal{A}_T acts by symmetric operators. This is a certain subspace of sections of the tensor product $C \otimes \mathcal{D}_{1/2}$ of the prequantization line bundle $C \rightarrow Z$ and a bundle $\mathcal{D}_{1/2} \rightarrow Z$ of half-densities on Z . The geometric quantization procedure has been extended to Poisson manifolds [16, 17] and to Jacobi manifolds [7].

We show that standard prequantization of the cotangent bundle T^*Q (e.g., [3, 14, 19]) provides the compatible prequantization of the Poisson manifold V^*Q such that the monomorphism ζ^* (7) is prolonged to a monomorphism of prequantum algebras.

In contrast with the prequantization procedure, polarization of T^*Q need not imply a compatible polarization of V^*Q , unless it includes the vertical cotangent bundle $V_\zeta T^*Q$

of the fibre bundle ζ (3), i.e., spans over vectors ∂^0 . The canonical real polarization of T^*Q , satisfying the condition

$$V_\zeta T^*Q \subset \mathbf{T}, \quad (11)$$

is the vertical polarization. It coincides with the vertical tangent bundle VT^*Q of T^*Q , i.e., spans over all the vectors ∂^λ . We show that this polarization and the corresponding metaplectic correction of T^*Q induces the compatible quantization of the Poisson manifold V^*Q such that the monomorphism of Poisson algebras ζ^* (7) is prolonged to a monomorphism of quantum algebras of V^*Q and T^*Q . The quantum algebra \mathcal{A}_V of V^*Q consists of functions on V^*Q which are at most affine in momenta p_k and have the Schrödinger representation in the space of half-densities on Q . It is essential that, since these operators does not contain the derivative with respect to time, the quantum algebra \mathcal{A}_V is a $C^\infty(\mathbf{R})$ -algebra.

We prove that the Schrödinger quantization of V^*Q yields geometric quantization of symplectic fibres of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ such that any quantum operator \hat{f} on V^*Q , restricted to the fibre V_t^*Q , $t \in \mathbf{R}$, coincides with the quantum operator on the symplectic manifold V_t^*Q of the function $f|_{V^*Q}$. Thus, the quantum algebra \mathcal{A}_V of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ can be seen as the instantwise $C^\infty(\mathbf{R})$ -algebra of its symplectic fibres. This agrees with the instantwise quantization of symplectic fibres $\{t\} \times T^*M$ of the direct product (2) in [14].

A fault of the Schrödinger representation on half-densities is that the Hamiltonian function \mathcal{H}^* (6) does not belong to the quantum algebra \mathcal{A}_T of T^*Q in general. A rather sophisticated solution of this problem for quadratic Hamiltonians has been suggested in [14]. However, the Laplace operator constructed in [14] does not fulfill the Dirac condition (8). If a Hamiltonian \mathcal{H} is a polynomial of momenta p_k , one can represent it as an element of the universal enveloping algebra of the Lie algebra \mathcal{A}_T , but this representation is not necessarily globally defined.

In order to include a Hamiltonian function \mathcal{H}^* (6) to the quantum algebra, one can choose the Hamiltonian polarization of T^*Q which contains the Hamiltonian vector field $\vartheta_{\mathcal{H}^*}$ of \mathcal{H}^* . However, it does not satisfy the condition (11) and does not define any polarization of the Poisson manifold V^*Q . This polarization is the necessary ingredient in a different variant of geometric quantization of V^*Q which is seen as a presymplectic manifold $(V^*Q, h^*\Omega)$. Given a trivialization (2), this quantization has been studied in [19]. In Section 5, its frame-covariant form is discussed.

Finally, since the quantum algebra \mathcal{A}_V of the Poisson manifold V^*Q is a $C^\infty(\mathbf{R})$ -algebra and since $\hat{p} = -i\partial/\partial t$, quantization of the classical evolution equation (5) defines

the connection

$$\nabla \widehat{f} = i[\widehat{\mathcal{H}}^*, \widehat{f}] \quad (12)$$

on the enveloping quantum algebra $\overline{\mathcal{A}}_V$ and describes quantum evolution in non-relativistic mechanics as a parallel transport along time.

2 Prequantization

Basing on the standard prequantization of the cotangent bundle T^*Q , we here construct the compatible prequantizations of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ and its symplectic leaves.

Recall the prequantization of T^*Q (e.g., [3, 14, 19]). Since its symplectic form Ω is exact and belongs to the zero de Rham cohomology class, the prequantization bundle is the trivial complex line bundle

$$C = T^*Q \times \mathbf{C} \rightarrow T^*Q, \quad (13)$$

whose Chern class c_1 is zero. Coordinated by $(q^\lambda, p_\lambda, c)$, it is provided with the admissible linear connection

$$A = dp_\lambda \otimes \partial^\lambda + dq^\lambda \otimes (\partial_\lambda + ip_\lambda c \partial_c) \quad (14)$$

with the strength form $F = -i\Omega$ and the Chern form

$$c_1 = \frac{i}{2\pi} F = \frac{1}{2\pi} \Omega.$$

The A -invariant Hermitian fibre metric on C is $g(c, c) = c\bar{c}$. The covariant derivative of sections s of the prequantization bundle C (13) relative to the connection A (14) along the vector field u on T^*Q takes the form

$$\nabla_u(s) = (u^\lambda \partial_\lambda - iu^\lambda p_\lambda) s. \quad (15)$$

Given a function $f \in C^\infty(T^*Q)$, the covariant derivative (15) along the Hamiltonian vector field

$$\vartheta_f = \partial^\lambda f \partial_\lambda - \partial_\lambda f \partial^\lambda, \quad \vartheta_f \lrcorner \Omega = -df$$

of f reads

$$\nabla_{\vartheta_f} = \partial^\lambda f (\partial_\lambda - ip_\lambda) - \partial_\lambda f \partial^\lambda.$$

Then, in order to satisfy the Dirac condition (8), one assigns to each element f of the Poisson algebra $C^\infty(T^*Q)$ the first order differential operator

$$\widehat{f}(s) = -i(\nabla_{\partial_f} + if)s = [-i\vartheta_f + (f - p_\lambda\partial^\lambda f)]s \quad (16)$$

on sections $s \in C(T^*Q)$ of the prequantization bundle C (13). For instance, the prequantum operators (16) for local functions $f = p_\lambda$, $f = q^k$, a global function $f = t$, and the constant function $f = 1$ read

$$\widehat{p}_\lambda = -i\partial_\lambda, \quad \widehat{q}^\lambda = i\partial^\lambda + q^\lambda, \quad \widehat{1} = 1.$$

For elements f of the Poisson subalgebra $C^\infty(V^*Q) \subset C^\infty(T^*Q)$, the Kostant–Souriau formula (16) takes the form

$$\widehat{f}(s) = [-i(\partial^k f \partial_k - \partial_\lambda f \partial^\lambda) + (f - p_k \partial^k f)]s. \quad (17)$$

Turn now to prequantization of the Poisson manifold $(V^*Q, \{, \}_V)$. The Poisson bivector w of the Poisson structure on V^*Q reads

$$w = \partial^k \wedge \partial_k = -[w, u]_{\text{SN}}, \quad (18)$$

where $[\cdot, \cdot]_{\text{SN}}$ is the Schouten–Nijenhuis bracket and $u = p_k \partial^k$ is the Liouville vector field on the vertical cotangent bundle $V^*Q \rightarrow Q$. The relation (18) shows that the Poisson bivector w is exact and, consequently, has the zero Lichnerowicz–Poisson cohomology class [8, 17]. Therefore, let us consider the trivial complex line bundle

$$C_V = V^*Q \times \mathbf{C} \rightarrow V^*Q \quad (19)$$

such that the zero Lichnerowicz–Poisson cohomology class of w is the image of the zero Chern class c_1 of C_V under the cohomology homomorphisms

$$H^*(V^*Q, \mathbf{Z}) \rightarrow H_{\text{deRh}}^*(V^*Q) \rightarrow H_{\text{LP}}^*(V^*Q).$$

Since the line bundles C (13) and C_V (19) are trivial, C can be seen as the pull-back ζ^*C_V of C_V , while C_V is isomorphic to the pull-back h^*C of C with respect to a section h (4) of the affine bundle (3). Since $C_V = h^*C$ and since the covariant derivative of the connection A (14) along the fibres of ζ (3) is trivial, let us consider the pull-back

$$h^*A = dp_k \otimes \partial^k + dq^k \otimes (\partial_k + ip_k c \partial_c) + dt \otimes (\partial_t - i\mathcal{H}c \partial_c) \quad (20)$$

of the connection A (14) onto $C_V \rightarrow V^*Q$ [9]. This connection defines the contravariant derivative

$$\nabla_\phi s_V = \nabla_{w^\sharp \phi} s_V \quad (21)$$

of sections s_V of $C_V \rightarrow V^*Q$ along one-forms ϕ on V^*Q , which corresponds to a contravariant connection A_V on the line bundle $C_V \rightarrow V^*Q$ [17]. It is readily observed that this contravariant connection does not depend on the choice of a section h . By virtue of the relation (21), the curvature bivector of A_V equals to $-iw$ [18], i.e., A_V is an admissible connection for the canonical Poisson structure on V^*Q . Then the Kostant–Souriau formula

$$\widehat{f}_V(s_V) = (-i\nabla_{\partial_{Vf}} + f)s_V = [-i(\partial^k f \partial_k - \partial_k f \partial^k) + (f - p_k \partial^k f)]s_V \quad (22)$$

defines prequantization of the Poisson manifold V^*Q .

In particular, the prequantum operators of functions $f \in C^\infty(\mathbf{R})$ of time alone reduces simply to multiplication $\widehat{f}_V s_V = f s_V$ by these functions. Consequently, the prequantum algebra $\widehat{C}^\infty(V^*Q)$ inherits the structure of a $C^\infty(\mathbf{R})$ -algebra.

It is immediately observed that the prequantum operator \widehat{f}_V (22) coincides with the prequantum operator $\widehat{\zeta^* f}$ (17) restricted to the pull-back sections $s = \zeta^* s_V$ of the line bundle C . Thus, prequantization of the Poisson algebra $C^\infty(V^*Q)$ on the Poisson manifold $(V^*Q, \{, \})$ is equivalent to its prequantization as a subalgebra of the Poisson algebra $C^\infty(T^*Q)$ on the symplectic manifold T^*Q .

The above prequantization of the Poisson manifold V^*Q yields prequantization of its symplectic leaves as follows.

Since $w^\sharp \phi = \phi^k \partial_k - \phi_k \partial^k$ is a vertical vector field on $V^*Q \rightarrow \mathbf{R}$ for any one-form ϕ on V^*Q , the contravariant derivative (21) defines a connection along each fibre V_t^*Q , $t \in \mathbf{R}$, of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$. This is the pull-back

$$A_t = i_t^* h^* A = dp_k \otimes \partial^k + dq^k \otimes (\partial_\lambda + ip_k c \partial_c)$$

of the connection $h^* A$ (20) on the pull-back bundle $i_t^* C_V \rightarrow V_t^*Q$ with respect to the imbedding $i_t : V_t^*Q \rightarrow V^*Q$. It is readily observed that this connection is admissible for the symplectic structure

$$\Omega_t = dp_k \wedge dq^k$$

on V_t^*Q , and provides prequantization of the symplectic manifold (V_t^*Q, Ω_t) . The corresponding prequantization formula is given by the expression (22) where functions f and sections s_V are restricted to V_t^*Q . Thus, the prequantization (22) of the Poisson manifold V^*Q is a leafwise prequantization [18].

3 Polarization

Given compatible prequantizations of the cotangent bundle T^*Q , the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ and its symplectic fibres, let us now construct their compatible polarizations.

Recall that, given a polarization \mathbf{T} of a prequantum symplectic manifold (Z, Ω) , the subalgebra $\mathcal{A}_T \subset C^\infty(Z)$ of functions f obeying the condition (10) is only quantized. Moreover, after further metaplectic correction, one consider a representation of this algebra in a quantum space E_T such that

$$\nabla_u e = 0, \quad \forall u \in \mathbf{T}(Z), \quad e \in E_T. \quad (23)$$

Recall that by a polarization of a Poisson manifold $(Z, \{, \})$ is meant a sheaf \mathbf{T}^* of germs of complex functions on Z whose stalks \mathbf{T}_z^* , $z \in Z$, are Abelian algebras with respect to the Poisson bracket $\{, \}$ [18]. One can also require that the algebras \mathbf{T}_z^* are maximal, but this condition need not hold under pull-back and push-forward operations. Let $\mathbf{T}^*(Z)$ be the structure algebra of global sections of the sheaf \mathbf{T}^* ; it is also called a Poisson polarization [16, 17]. A quantum algebra \mathcal{A}_T associated to the Poisson polarization \mathbf{T}^* is defined as a subalgebra of the Poisson algebra $C^\infty(Z)$ which consists of functions f such that

$$\{f, \mathbf{T}^*(Z)\} \subset \mathbf{T}^*(Z).$$

Polarization of a symplectic manifold yields its maximal Poisson polarization, and *vice versa*.

There are different polarizations of the cotangent bundle T^*Q . We will consider those polarizations of T^*Q whose direct image as Poisson polarizations onto V^*Q with respect to the morphism ζ (3) are polarizations of the Poisson manifold V^*Q . This takes place if the germs of polarization \mathbf{T}^* of the Poisson manifold $(T^*Q, \{, \}_T)$ are constant along the fibres of the fibration ζ (3) [18], i.e., are germs of functions independent of the momentum coordinate $p_0 = p$. It means that the corresponding polarization \mathbf{T} of the symplectic manifold T^*Q is vertical with respect to the fibration $T^*Q \rightarrow \mathbf{R}$, i.e., obeys the condition (11). A short calculation shows that, in this case, the associated quantum algebra \mathcal{A}_T consists of functions $f \in C^\infty(T^*Q)$ which are at most affine in the momentum coordinate p_0 . Moreover, given such a polarization, the equality (23) implies the equality

$$\nabla_{u_0 \partial^0} e = 0, \quad e \in E_T,$$

for any vertical vector field $u_0 \partial^0$ on the fibre bundle $T^*Q \rightarrow V^*Q$. Then the prequantization formulas (17) and (22) for the Poisson algebra $C^\infty(V^*Q)$ coincide on quantum

spaces, i.e., the monomorphism ζ^* (7) is prolonged to monomorphism of quantum algebras of V^*Q and T^*Q .

The vertical polarization VT^*Q of T^*Q obeys the condition (11). It is a strongly admissible polarization, and its integral manifolds are fibres of the cotangent bundle $T^*Q \rightarrow Q$. One can verify easily that the associated quantum algebra \mathcal{A}_T consists of functions on T^*Q which are at most affine in momenta p_λ . The quantum space E_T associated to the vertical polarization obeys the condition

$$\nabla_{u^\lambda \partial_\lambda} e = 0, \quad \forall e \in E_T. \quad (24)$$

Therefore, the operators of the quantum algebra \mathcal{A}_T on this quantum space read

$$f = a^\lambda(q^\mu)p_\lambda + b(q^\mu), \quad \hat{f} = -i\nabla_{a^\lambda \partial_\lambda} + b. \quad (25)$$

This is the Schrödinger representation of T^*Q .

The vertical polarization of T^*Q defines the maximal polarization \mathbf{T}^* of the Poisson manifold V^*Q which consists of germs of functions constant on the fibres of $V^*Q \rightarrow Q$. The associated quantum space E_V obeys the condition

$$\nabla_{u^k \partial_k} e = 0, \quad \forall e \in E_V. \quad (26)$$

The quantum algebra \mathcal{A}_V corresponding to this polarization of V^*Q consists of functions on V^*Q which are at most affine in momenta p_k . Their quantum operators read

$$f = a^\lambda(q^\mu)p_k + b(q^\mu), \quad \hat{f} = -i\nabla_{a^k \partial_k} + b. \quad (27)$$

This is the Schrödinger representation of V^*Q .

In turn, each symplectic fibre V_t^*Q , $t \in \mathbf{R}$, of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ is provided with the pull-back polarization $\mathbf{T}_t^* = i_t^* \mathbf{T}^*$ with respect to the Poisson morphism $i_t : V_t^*Q \rightarrow V^*Q$. The corresponding distribution \mathbf{T}_t coincides with the vertical tangent bundle of the fibre bundle $V_t^*Q \rightarrow Q_t$. The associated quantum algebra \mathcal{A}_t consists of functions on $V_t^*Q_t$ which are at most affine in momenta p_k , while the quantum space E_t obeys the condition similar to (26). Therefore, the representation of the quantum algebra \mathcal{A}_t takes the form (27). It follows that the Schrödinger representation the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ is a fibrewise representation.

4 Metaplectic correction

To complete the geometric quantization procedure of V^*Q , let us consider the metaplectic correction of the Schrödinger representations of T^*Q and V^*Q .

The representation of the quantum algebra \mathcal{A}_T (25) can be defined in the subspace of sections of the line bundle $C \rightarrow T^*Q$ which fulfill the relation (24). This representation reads

$$f = a^\lambda(q^\mu)p_\lambda + b(q^\mu), \quad \hat{f} = -ia^\lambda\partial_\lambda + b. \quad (28)$$

Therefore, it can be restricted to the sections s of the pull-back line bundle $C_Q = \hat{0}^*C \rightarrow Q$ where $\hat{0}$ is the canonical zero section of the cotangent bundle $T^*Q \rightarrow Q$. However, this is not yet a representation in a Hilbert space.

Let Q be an oriented manifold. Applying the general metaplectic technique [3, 19], we come to the vector bundle $\mathcal{D}_{1/2} \rightarrow Q$ of complex half-densities on Q with the transition functions $\rho' = J^{-1/2}\rho$, where J is the Jacobian of the coordinate transition functions on Q . Since $C_Q \rightarrow Q$ is a trivial bundle, the tensor product $C_Q \otimes \mathcal{D}_{1/2}$ is isomorphic to $\mathcal{D}_{1/2}$. Therefore, the quantization formula (28) can be extended to sections of the half-density bundle $\mathcal{D}_{1/2} \rightarrow Q$ as

$$f = a^\lambda(q^\mu)p_\lambda + b(q^\mu), \quad \hat{f}\rho = (-i\mathbf{L}_{a^\lambda\partial_\lambda} + b)\rho = (-ia^\lambda\partial_\lambda - \frac{i}{2}\partial_\lambda(a^\lambda) + b)\rho, \quad (29)$$

where \mathbf{L} denotes the Lie derivative. The second term in the right-hand side of this formula is a metaplectic correction. It makes the operator \hat{f} (29) symmetric with respect to the Hermitian form

$$\langle \rho_1 | \rho_2 \rangle = \left(\frac{1}{2\pi} \right)^m \int_Q \rho_1 \bar{\rho}_2$$

on the pre-Hilbert space E_T of sections of $\mathcal{D}_{1/2}$ with compact support. The completion \overline{E}_T of E_T provides a Hilbert space of the Schrödinger representation of the quantum algebra \mathcal{A}_T , where the operators (29) are essentially self-adjoint, but not necessarily bounded. Of course, functions of compact support on the time axis \mathbf{R} have a limited physical application, but we can always restrict our consideration to some bounded interval of \mathbf{R} .

Since, in the case of the vertical polarization, there is a monomorphism of the quantum algebra \mathcal{A}_V to the quantum algebra \mathcal{A}_T , one can define the Schrödinger representation of \mathcal{A}_V by the operators

$$f = a^k(q^\mu)p_k + b(q^\mu), \quad \hat{f}\rho = (-ia^k\partial_k - \frac{i}{2}\partial_k(a^k) + b)\rho \quad (30)$$

in the same space of complex half-densities on Q as that of \mathcal{A}_T . Moreover, this representation preserves the structure of \mathcal{A}_V as a $C^\infty(\mathbf{R})$ -algebra.

The metaplectic correction of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ also provides the metaplectic corrections of its symplectic leaves as follows. It is readily observed that the Jacobian J restricted to a fibre Q_t , $t \in \mathbf{R}$, is a Jacobian of coordinate transformations on Q_t . Therefore, any half-density ρ on Q , restricted to Q_t , is a half-density on Q_t . Then the representation (30) restricted to a fibre V_t^*Q is exactly the metaplectic correction of the Schrödinger representation of the symplectic manifold V_t^*Q on half-densities on Q_t .

Thus, the quantum algebra \mathcal{A}_V of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$, given by the operators (30), can be seen as the instantwise algebra of operators on its symplectic fibres.

The representation (30) can be extended locally to functions on T^*Q which are polynomials of momenta p_λ . These functions can be represented by elements of the universal enveloping algebra $\overline{\mathcal{A}}_T$ of the Lie algebra \mathcal{A}_T , but this representation is not necessarily globally defined. For instance, a generic quadratic Hamiltonian

$$\mathcal{H} = a^{jk}(q^\lambda)p_j p_k + b^k(q^\lambda)p_k + c(q^\lambda) \quad (31)$$

leads to the Hamiltonian function $\mathcal{H}^* = p + \mathcal{H}$ which is not an element of $\overline{\mathcal{A}}_T$ because of the quadratic term. This term can be quantized only locally, unless the Jacobian of the coordinate transition functions on Q is independent of fibre coordinates q^k on Q .

5 Presymplectic quantization

As was mentioned above, to include a Hamiltonian function \mathcal{H}^* (6) to the quantum algebra, one can choose the Hamiltonian polarization of T^*Q . This polarization accompanies a different approach to geometric quantization of the momentum phase space of non-relativistic mechanics V^*Q which is considered as a presymplectic manifold. In comparison with the above one, this quantization does not lead to quantization of the Poisson algebra of functions on V^*Q as follows.

Every global section h (4) of the affine bundle ζ (3) yields the pull-back Hamiltonian form

$$H = h^*\Xi = p_k dq^k - \mathcal{H}dt \quad (32)$$

on V^*Q . With respect to a trivialization (2), the form H is the well-known integral invariant of Poincaré–Cartan [1]. Given a Hamiltonian form H (32), there exists a unique Hamiltonian connection

$$\gamma_H = \partial_t + \partial^k \mathcal{H} \partial_k - \partial_k \mathcal{H} \partial^k \quad (33)$$

on the fibre bundle $V^*Q \rightarrow \mathbf{R}$ such that

$$\gamma_H]dH = 0 \tag{34}$$

[8, 12]. It defines the Hamilton equations on V^*Q .

A glance at the equation (34) shows that one can think of the Hamiltonian connection γ_H as being the Hamiltonian vector field of a zero Hamiltonian with respect to the presymplectic form dH on V^*Q . Therefore, one can study geometric quantization of the presymplectic manifold (V^*Q, dH) .

Usually, geometric quantization is not applied directly to a presymplectic manifold (Z, ω) , but to a symplectic manifold (Z', ω') such that the presymplectic form ω is a pull-back of the symplectic form ω' . Such a symplectic manifold always exists. The following two possibilities are usually considered: (i) (Z', ω') is a reduction of (Z, ω) along the leaves of the characteristic distribution of the presymplectic form ω of constant rank [6, 15], and (ii) there is a coisotropic imbedding of (Z, ω) to (Z', ω') [4, 5].

In application to (V^*Q, dH) , the reduction procedure however meets difficulties. Since the kernel of dH is generated by the vectors $(\partial_t, \partial_k \mathcal{H} \partial^k - \partial^k \mathcal{H} \partial_k, k = 1, \dots, m)$, the presymplectic form dH in physical models is almost never of constant rank. Therefore, one has to provide an exclusive analysis of each physical model, and to cut out a certain subset of V^*Q in order to use the reduction procedure.

The second variant of geometric quantization of the presymplectic manifold (V^*Q, dH) seems more attractive because the section h (4) is a coisotropic imbedding. Indeed, the tangent bundle TN_h of the closed imbedded submanifold $N_h = h(V^*Q)$ of T^*Q consists of the vectors

$$u = -(u^\mu \partial_\mu \mathcal{H} + u_j \partial^j \mathcal{H}) \partial^0 + u_k \partial^k + u^\mu \partial_\mu. \tag{35}$$

Let us prove that the orthogonal distribution $\text{Orth}_\Omega TN_h$ of TN_h with respect to the symplectic form Ω belongs to TN_h . By definition, it consists of the vectors $u \in T_z T^*Q$, $z \in T^*Q$, such that

$$u]v]\Omega = 0, \quad \forall v \in T_z N_h.$$

A simple calculation shows that these vectors obey the conditions

$$-u^0 \partial^i \mathcal{H} + u^i = 0, \quad u^0 \partial_\mu - u_\mu = 0$$

and, consequently, take the form (35).

The image $N_h = h(V^*Q)$ of the coisotropic imbedding h is given by the constraint

$$\mathcal{H}^* = p + \mathcal{H}(t, q^k, p_k) = 0.$$

Then the geometric quantization of the presymplectic manifold (V^*Q, dH) consists in geometric quantization of the cotangent bundle T^*Q and setting the quantum constraint condition

$$\widehat{\mathcal{H}}^* \psi = 0$$

on physically admissible quantum states. This condition implies that $\widehat{\mathcal{H}}^*$ belongs to the quantum algebra of T^*Q . It takes place if the above mentioned Hamiltonian polarization of T^*Q , which contains the Hamiltonian vector field

$$\vartheta_{\mathcal{H}^*} = \partial_0 + \partial^k \mathcal{H} \partial_k - \partial_k \mathcal{H} \partial^k, \quad (36)$$

is used.

Such a polarization of T^*Q always exists. Indeed, any section h (4) of the affine bundle $T^*Q \rightarrow V^*Q$ defines the splitting

$$a_\lambda \partial^\lambda = a_k (\partial^k - \partial^k \mathcal{H} \partial^0) + (a_0 + a_k \partial^k \mathcal{H}) \partial^0$$

of the vertical tangent bundle VT^*Q of $T^*Q \rightarrow Q$. One can justify this fact, e.g., by inspection of the coordinate transformation law. Then elements $(\partial^k - \partial^k \mathcal{H} \partial^0, k = 1, \dots, m)$, and the values of the Hamiltonian vector field $\vartheta_{\mathcal{H}^*}$ (36) obey the polarization condition (9) and generates a polarization of T^*Q . It is clear that the Hamiltonian polarization does not satisfy the condition (11), and does not define any polarization of the Poisson manifold V^*Q .

Nevertheless, given a trivialization (2), symplectic fibres V_t^*Q , $t \in \mathbf{R}$, of the Poisson bundle $V^*Q \rightarrow \mathbf{R}$ can be provided with the Hamiltonian polarization \mathbf{T}_t generated by vectors $\partial_k \mathcal{H} \partial^k - \partial^k \mathcal{H} \partial_k$, $k = 1, \dots, m$, except the points where

$$d\mathcal{H} = \partial_k \mathcal{H} dq^k + \partial^k \mathcal{H} dp_k = 0.$$

This is a standard polarization in conservative Hamiltonian mechanics of one-dimensional systems, but it requires an exclusive analysis of each model.

6 Classical and quantum evolution equations

Turn now to the evolution equation in classical and quantum non-relativistic mechanics.

Given a Hamiltonian connection γ_H on the momentum phase space V^*Q , let us consider the Lie derivative

$$\mathbf{L}_{\gamma_H} f = \gamma_H \rfloor df = (\partial_t + \partial^k \mathcal{H} \partial_k - \partial_k \mathcal{H} \partial^k) f \quad (37)$$

of a function $f \in C^\infty(V^*Q)$ along γ_H . This equality is the evolution equation in classical non-relativistic Hamiltonian mechanics. Substituting a solution of the Hamilton equations in its right-hand side, one obtains the time evolution of f along this solution. Given a trivialization (2), the evolution equation (37) can be written as

$$\mathbf{L}_\gamma f = \partial_t f + \{\mathcal{H}, f\}_V.$$

However, taken separately, the terms in its right-hand side are ill-behaved under time-dependent transformations. Let us bring the evolution equation into the frame-covariant form.

The affine bundle ζ (3) is modelled over the trivial line bundle $V^*Q \times \mathbf{R} \rightarrow V^*Q$. Therefore, Hamiltonian forms H constitute an affine space modelled over the vector space $C^\infty(V^*Q)$. To choose a centre of this affine space, let us consider a connection

$$\Gamma = \partial_t + \Gamma^k \partial_k \quad (38)$$

on the configuration bundle $Q \rightarrow \mathbf{R}$. By definition, it is a section of the affine bundle (3), and yields the Hamiltonian form

$$H_\Gamma = \Gamma^* \Xi = p_k dq^k - \mathcal{H}_\Gamma dt, \quad \mathcal{H}_\Gamma = p_k \Gamma^k dt.$$

Its Hamiltonian connection is the canonical lift

$$V^* \Gamma = \partial_t + \Gamma^i \partial_i - p_i \partial_j \Gamma^i \partial^j$$

of the connection Γ onto $V^*Q \rightarrow \mathbf{R}$. Then any Hamiltonian form H (32) on the momentum phase space V^*Q admits the splittings

$$H = H_\Gamma - \widetilde{\mathcal{H}}_\Gamma dt, \quad \widetilde{\mathcal{H}}_\Gamma = \mathcal{H} - p_k \Gamma^k, \quad (39)$$

where $\widetilde{\mathcal{H}}_\Gamma$ is a function on V^*Q . The physical meaning of this splitting becomes clear due to the fact that every trivialization of $Q \rightarrow \mathbf{R}$ yields a complete connection Γ on Q , and *vice versa* [8, 9]. From the physical viewpoint, the vertical part of this connection Γ

(38) can be seen as a velocity of an "observer", and Γ characterizes a reference frame in non-relativistic time-dependent mechanics [8, 11, 12]. Then one can show that $\widetilde{\mathcal{H}}_\Gamma$ in the splitting (39) is the energy function with respect to this reference frame [2, 8, 12].

Given the splitting (39), the evolution equation can be written in the frame-covariant form

$$\mathbf{L}_{\gamma_H} f = V^*\Gamma \rfloor H + \{\widetilde{\mathcal{H}}_\Gamma, f\}_V.$$

However, the first term in its right-hand side is not reduced to the Poisson bracket on V^*Q , and is not quantized in the framework of geometric quantization of the Poisson manifold V^*Q . To bring the right-hand side of the evolution equation into a Poisson bracket alone, let us consider the pull-back ζ^*H of the Hamiltonian form $H = h^*\Xi$ onto the cotangent bundle T^*Q . It is readily observed that the difference $\Xi - \zeta^*H$ is a horizontal 1-form on $T^*Q \rightarrow \mathbf{R}$, and we obtain the function (6) on T^*Q . Then the relation

$$\zeta^*(\mathbf{L}_{\gamma_H} f) = \vartheta_{H^*}(\zeta^* f) = \{\mathcal{H}^*, \zeta^* f\}_T,$$

holds for any element f of the Poisson algebra $C^\infty(V^*Q)$ [10].

Since the quantum algebra \mathcal{A}_V of the Poisson manifold V^*Q can be seen as the instantwise algebra, one can quantize the evolution equation (5) as follows.

Given the Hamiltonian function \mathcal{H}^* (6), let $\widehat{\mathcal{H}}^*$ be the corresponding quantum operator written, e.g., as an element of universal enveloping algebra of the quantum algebra \mathcal{A}_T . The bracket (12) defines a derivation of the enveloping algebra $\overline{\mathcal{A}}_V$ of the quantum algebra \mathcal{A}_V , which is also a $C^\infty(\mathbf{R})$ -algebra. Moreover, since $\widehat{p} = -i\partial/\partial t$, the derivation (12) obeys the Leibniz rule

$$\nabla(r(t)\widehat{f}) = \partial_t r(t)\widehat{f} + r(t)\nabla\widehat{f}.$$

Therefore, it is a connection on the $C^\infty(\mathbf{R})$ -algebra $\overline{\mathcal{A}}_V$, and defines quantum evolution of $\overline{\mathcal{A}}_V$ as a parallel transport along time [9, 13].

References

- [1] V.Arnold, *Mathematical Methods of Classical Mechanics* (Springer, Berlin, 1978).
- [2] A.Echeverría Enríquez, M.Muñoz Lecanda and N.Román Roy, Non-standard connections in classical mechanics, *J. Phys. A* **28** (1995) 5553.

- [3] A.Echeverría Enríquez, M.Muñoz Lecanda, N.Román Roy, and C.Victoria-Monge, Mathematical foundations of geometric quantization, *Extracta Math.* **13** (1998) 135.
- [4] M.Gotay and J.Śniatycki, On the quantization of presymplectic dynamical systems via coisotropic imbeddings, *Commun. Math. Phys.* **82** (1981) 377.
- [5] M.Gotay, On coisotropic imbeddings of presymplectic manifolds, *Proc. Amer. Math. Soc.* **84** (1982) 111.
- [6] M.Gotay, Constraints, reduction and quantization, *J. Math. Phys.* (1986) 2051.
- [7] M. de León, J.Marrero and E.Padron, On the geometric quantization of Jacobi manifolds, *J. Math. Phys.* **38** (1997) 6185.
- [8] L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).
- [9] L.Mangiarotti and G.Sardanashvily, *Connections in Classical and Quantum Field Theory* (World Scientific, Singapore, 2000).
- [10] L.Mangiarotti and G.Sardanashvily, Constraints in Hamiltonian time-dependent mechanics, *J. Math. Phys.* **41** (2000) 2858
- [11] E.Massa and E.Pagani, Jet bundle geometry, dynamical connections and the inverse problem of Lagrangian mechanics, *Ann. Inst. Henri Poincaré* **61** (1994) 17.
- [12] G.Sardanashvily, Hamiltonian time-dependent mechanics, *J. Math. Phys.* **39** (1998) 2714.
- [13] G.Sardanashvily, Classical and quantum mechanics with time-dependent parameters, *J. Math. Phys.* **41** (2000) 5245.
- [14] J.Śniatycki, *Geometric Quantization and Quantum Mechanics* (Springer-Verlag, Berlin, 1980).
- [15] I.Vaisman, Geometric quantization on presymplectic manifold, *Monatsh. Math.* **96** (1983) 293.
- [16] I.Vaisman, On the geometric quantization of Poisson manifolds, *J. Math. Phys.* **32** (1991) 3339.

- [17] I.Vaisman, *Lectures on the Geometry of Poisson Manifolds* (Birkhäuser Verlag, Basel, 1994).
- [18] I.Vaisman, On the geometric quantization of the symplectic leaves of Poisson manifolds, *Diff. Geom. Appl.* **7** (1997) 265.
- [19] N.Woodhouse, *Geometric Quantization* (Clarendon Press, Oxford, 1980) (2nd ed. 1992).