

Laser-Induced Giant Quasi-Molecules in Optical Lattices

Bimalendu Deb and Gershon Kurizki

Department of Chemical Physics, Weizmann Institute of Science, 76100 Rehovot, Israel
(December 2, 2024)

We analyze atom-atom interactions in optical lattices due to a laser-induced long-range interatomic force which prevails over the usual London-van der-Waals forces. This force, which can be generated by an intense laser field at a wavelength other than that of the lattice-generating laser, is shown to bind pairs of cold atoms trapped at different lattice sites, and cause their translational quantum correlations (spatial entanglement).

PACS numbers: 34.10.+x, 33.80.-b, 42.50.Ct, 42.50.Fx

The best-known laser-induced force acting upon or between atoms is the "dipole force", which results from the action of the gradient of the laser field. It is used to confine cold atoms in optical traps [1] and optical lattices [2], and can modify cold-atom collisions and photoassociation [3]. Recently, the mechanical effect of another laser-induced force on cold atoms has come to the fore, that of dipole-dipole interactions generated by off-resonant lasers [4]. It has been shown that in the near-zone region of mutually-detuned orthogonal triads of laser-beams the time-average of the dipole-dipole interaction produces a gravity-like interatomic potential ($1/r$ attraction), which can create a self-bound Bose condensate [5]. Partial condensate confinement (in certain directions), which still requires an external trap, may also occur due to the $1/r^3$ near-zone dipole-dipole interaction induced by a single laser [6].

Here we put forward a new effect of the laser-induced dipole-dipole interaction: a mechanism for binding together pairs of neutral atoms localized at different sites of an optical lattice while repelling them from the same site. We show that this mechanism can form novel diatomic or multi-atomic states with submicron interatomic separations: "giant quasi-molecules (diatoms)" with translational quantum correlations (spatial entanglement). An optical lattice provides appropriate conditions for the formation of such states: it accommodates cold atoms whose mean separations r are large enough to be in the Casimir regime [4] ($r \geq 2\pi \times 137a_0$, a_0 being the Bohr radius). At separations compatible with the Casimir regime, an additional plane-wave field with a moderate intensity would suffice to induce long-range forces that prevail over the ordinary van der Waals-London forces arising from vacuum fluctuations [4]. By contrast, at smaller separations ($r \leq 2\pi \times 137a_0$) laser-induced interatomic forces cannot overcome the ordinary van der Waals forces, since this would require a field intensity large enough to ionize the atoms. Another advantage of an optical lattice is that it allows the fixing of the polarization direction of the additional plane wave field relative to the interatomic axis. The optical lattice can first be used to create a periodic array of cold atoms. Once the intense plane-wave field is applied, it can form giant diatoms, or a chain of such diatoms, by modifying the long-range interatomic forces at separations which are of the order of lattice period. Then the atoms will remain bound by the field-modified dipole-dipole forces even if the lattice potential is switched off.

The essential effect analyzed here stems from the induced dipole-dipole interaction potential between two unexcited neutral atoms A and B in the presence of a monochromatic, circularly-polarized, electromagnetic plane wave at an extremely off-resonant frequency. It is given by [4]

$$V_{AB} = -\frac{2\pi\alpha_A\alpha_B I}{c} \cos(kr \cos \theta) \times \left[\frac{\{\cos(kr) + kr \sin(kr)\}(1 - 3\cos^2 \theta)}{r^3} + \frac{k^2(1 + \cos^2 \theta) \cos(kr)}{r} \right]. \quad (1)$$

Here θ is the angle between the interatomic separation and the wave-vector $k = \omega/c$ of the plane-wave field, I is the intensity of the field and $\alpha_C = \frac{2\omega_A \mathbf{d}^2}{\hbar(\omega_C^2 - \omega^2)}$ is the dynamic polarizability of the atom C (=A,B), with $\mathbf{d}$ and ω_C being the dipole moment and the transition frequency of the atom, respectively [we shall set $\alpha_A = \alpha_B = \alpha$]. Two situations are of interest: (i) In a one-dimensional (1D) lattice, depending on whether the circularly polarized wave is parallel or perpendicular to the interatomic separation, this interatomic potential is denoted by $V^{\parallel}$ or $V^{\perp}$, respectively. The $V^{\parallel}$ potential provides on-site ($kr \ll 1$) repulsion, $V^{\parallel} \simeq \frac{4\pi I \alpha^2}{cr^3}$, and adjacent-site attraction $V^{\parallel} \simeq -\frac{4\pi I \alpha^2 k^2}{cr}$ for $kr \simeq n\pi$ ($n = 1, 2, \dots$), with $k \simeq k_L$ being close to the wavevector k_L of the lattice potential $V_L(r) = V_0 \sin^2 k_L r$. These properties allow the formation of stable giant molecules. By contrast, $V^{\perp}$ is unsuitable, yielding on-site attraction (Fig.1). (ii) In a 2D lattice, a *partly-stable quasi-binding effect* is achieved by the time-averaged sum of the two potentials, $V_+ = V^{\parallel} + V^{\perp}$, corresponding to two circularly polarized laser beams propagating along the orthogonal X and Y axes, whose phase is fast oscillating due to their frequency difference [5,7]. When $kr \simeq 2n\pi$, where $n \geq 1$,

pairs of ultracold atoms can be trapped in the periodic long-range interatomic potential-wells of $V_+(r)$ whose depth decreases with n . Both cases (i) and (ii) can lead to a crystalline structure of bound atom-pairs.

For pair binding to occur predominantly at adjacent sites at a temperature T , the following conditions need be fulfilled: (a) $V(r) \geq k_B T$, which constrains the intensity to $I > \frac{4\pi k_B T}{\alpha^2 k^3}$. (b) In order to effectively suppress the photon absorption and the resulting atomic excitation by the additional plane-wave laser field, the saturation parameter $S(= \frac{2\Omega^2}{4\delta^2 + \Gamma^2})$, with Ω being the laser Rabi frequency, $\delta = \omega - \omega_A$ its detuning and Γ the transition line width, should be much less than unity, $S \ll 1$. (c) The frequency and intensity of the applied field should be below the photoionization threshold. Conditions (a) and (b) can be combined to require $\frac{4\pi k_B T}{\alpha^2 k^3 I} < S \ll 1$ in what follows (Fig.1).

We proceed with the general theoretical framework that allows one to treat atom-atom scattering in an optical lattice for any interatomic potential. The Hamiltonian for a system of two interacting atoms A and B in an optical lattice is $H = H_0 + V_{AB}$. Here $V_{AB}(r)$ is the interaction potential, which is assumed to be a function of the interatomic separation $r = |\mathbf{r}_A - \mathbf{r}_B|$ only, and

$$H_0 = -\frac{\hbar^2}{2m_A} \nabla_A^2 - \frac{\hbar^2}{2m_B} \nabla_B^2 + U_A(r_A) + U_B(r_B), \quad (2)$$

$U_{A(B)}$ is the periodic potential of the lattice experienced by the atom $A(B)$.

The formation of the quasi-bound pair is revealed by the analysis of the transition matrix

$$\mathbf{T} = V_{AB}(\mathbf{1} - G V_{AB})^{-1} \quad (3)$$

G being the unperturbed Green function (associated with H_0). Let D be the determinant of the matrix $(\mathbf{1} - G V_{AB})$. When $\text{Re}(D) = 0$ for a positive energy $E = E_r$, a resonance may occur, implying the existence of two-atom quasi-bound states in the lattice, but only if its width (its reciprocal lifetime) is given by the positive value [8]

$$\Gamma_r = 2 \frac{\text{Im}(D)}{\frac{d}{dE} \text{Re}(D)}. \quad (4)$$

Information about the two-atom scattering can be inferred from the measurement of the transport properties [8] and light scattering spectra, which depend on the atomic density of states in the lattice. The change in the two-atom density of states due to the interatomic interaction potential V_{AB} is given by

$$\delta\rho = -\frac{1}{\pi N} \text{Im} \left[\frac{d}{dE} (\ln D) \right], \quad (5)$$

N being the normalization constant. In the vicinity of a genuine resonance located at $E = E_r$ (i.e., if the conditions leading to Eq. (4) are satisfied), Eq. (5) reduces to the Lorentzian shape [8]

$$\delta\rho = \frac{\Gamma_r/2\pi}{(E - E_r)^2 + \Gamma_r^2/4}. \quad (6)$$

This expression measures *the rapid enhancement* of the two-atom density of states relative to that of a perfect optical lattice (without atom-atom interaction),

$$\rho^{(0)}(E) = -\frac{1}{\pi N} \text{Im}[\text{Tr}(G)]. \quad (7)$$

We now consider that the atoms are located near the minima of the periodic potential wells of the optical lattice, for which the tight-binding approximation is valid. As the basis states, we choose the Wannier functions $\chi_n(r_C - R_\mu)$ for the atom $C = A, B$. Here n refers to the energy band index and μ denotes the site index. The Wannier function has the property: $\int d\vec{r} \chi_n(\vec{r} - \vec{R}_\mu) H_0 \chi_n(\vec{r} - \vec{R}_{\mu'}) = \epsilon_n (\vec{R}_\mu - \vec{R}_{\mu'})$, ϵ_n being the n th band energy eigenvalue. The symmetrized or antisymmetrized product of the Wannier functions of the two atoms is denoted by

$$|nm; \mu\nu\rangle = \mathcal{N} [\chi_n(\vec{r}_A - \vec{R}_\mu) \chi_m(\vec{r}_B - \vec{R}_\nu) \pm \chi_n(\vec{r}_B - \vec{R}_\mu) \chi_m(\vec{r}_A - \vec{R}_\nu)], \quad (8)$$

where $\mathcal{N}$ is a normalization constant, and $+$ ($-$) refers to bosons (fermions). Here we assume that $\vec{R}_\mu \neq \vec{R}_\nu$, i.e. no lattice-site is occupied by more than one atom. In the plots and estimates given below only bosons are considered.

In a 1D lattice, $R_\mu = \mu a$, where a is the lattice spacing μ is an integer, we then have the dispersion relation

$$\epsilon_n(k) = \lambda_n(0) + 2\lambda_n(1) \cos ka \quad (9)$$

where $\lambda_n(0) = \int dx \chi_n^*(x) H_0^A \chi_n(x)$ and $\lambda_n(1) = \int dx \chi_n^*(x) H_0^A \chi_n(x-a)$. In the harmonic approximation of the potential, $\chi_n(x)$ is the n th level harmonic oscillator eigenfunction. Under these conditions, the unperturbed Green function can be expressed in the chosen basis as [9]

$$\begin{aligned} G_{nm}(\mu\nu, \mu'\nu') &= \langle nm; \mu\nu | \frac{1}{E^+ - H_0} | nm; \mu'\nu' \rangle \\ &= -iC \int_0^\infty dt \exp(-iE't) [J_{n+}(t) + J_{n-}(t)]. \end{aligned} \quad (10)$$

Here $n_\pm = |(\mu - \nu) \pm (\mu' - \nu')|$, J_n is the n th order Bessel function, $E' = \{E - \lambda_n(0) - \epsilon_m(0)\}/\lambda_{nm}(1)$ and $C = 2\mathcal{N}^2/(2\pi\lambda_{nm}(1))$, with $\lambda_{nm}(1) = 2\{\lambda_n(1) + \epsilon_m(1)\}$.

In a 2D square lattice ($\vec{R}_\mu = \hat{i}\mu_1 a + \hat{j}\mu_2 a$; μ_i being an integer), under the tight-binding condition, we have

$$\epsilon_n(\vec{k}) = \lambda_n(0) + 2\lambda_n(1) [\cos k_x a + \cos k_y a] \quad (11)$$

where $\lambda_n(0) = \int d^2 r_A \chi_n^*(r_A - R_{\mu_1\mu_2}) H_0^A \chi_n(r_A - R_{\mu_1\mu_2})$ and $\lambda_n(1) = \int d^2 r_A \chi_n^*(r_A - R_{\mu_1\mu_2}) H_0^A \chi_n(r_A - R_{\mu_1+1, \mu_2})$. The Green function can be calculated for this 2D lattice similarly to its 1D counterpart (10).

For our numerical analysis, we consider that the two bosonic atoms are in the lowest band, i.e., $n = m = 0$. Since the van der Waals attraction ($-c_6/r^6$) is negligible in comparison to the laser-induced dipole-dipole barrier of $V_{||}$ (Fig.1) in the region $r_0 < r < 1/k$, where r_0 is of the order of several thousand Angstroms, we have imposed a cut-off separation $r_c (> r_0)$ for calculating the low-energy collisional dynamics of the two atoms in the 1D or 2D lattice. Figure 2 shows the resulting variation of the real parts of the determinant D as a function of energy and its vanishing at resonance in 1D and 2D lattices. The corresponding atomic density of states (7) and its near-resonant enhancement (5) or (6) are shown in Fig.3.

In the nearly-free approximation (NFA), which holds for the atoms that occupy the states near the top of the lattice potential wells, the dispersion relation is given by $\epsilon_n(k) = \gamma k^2$ where γ is a constant. The individual atomic state can then be approximated by a plane wave, so that the corresponding Wannier function for a one-dimensional lattice can be expressed as $\chi_n(X_\mu) = \frac{\sqrt{a} \sin(\pi X_\mu/a)}{\pi X_\mu}$, with $X_n = x - \mu a$. The Green function in this case is given by

$$G_{nm}(\mu\nu, \mu'\nu') = -\frac{a}{2\pi\gamma_+} \frac{\exp(ikX_\xi)}{k} \quad (12)$$

where $\gamma_+ = \gamma_A + \gamma_B$, $X_\xi = (X_\mu - X_\nu) - (X_{\mu'} - X_{\nu'})$ and $k = \sqrt{E/\gamma_+}$. This NFA Green function yields a two-atom scattering resonance, but at a much lower energy and with a much larger width than its tightly bound counterpart, which therefore would be hardly observable.

The solution of the Schroedinger equation, $H\Psi = E\Psi$ for the tightly-binding limit and in the presence of V_{AB} can be written as

$$\Psi = \sum_{nm} \sum_{\mu\nu} C_{nm} |nm; \mu\nu\rangle. \quad (13)$$

This is a superposition of the two-atom Wannier functions (8) with coefficients C_{nm} obtainable from the Lippman-Schwinger matrix equation [8]

$$\begin{aligned} \sum_{\mu'\nu'} \langle nm; \mu\nu | (H_0 - E) | nm; \mu'\nu' \rangle C_{nm}(R_\mu, R_\nu) \\ + \sum_{\mu'\nu'} V_{nm; \mu\nu, \mu'\nu'} C_{nm}(R_{\mu'}, R_{\nu'}) = 0, \end{aligned} \quad (14)$$

where $V_{nm; \mu\nu, \mu'\nu'} = \langle nm; \mu\nu | V_{AB} | nm; \mu'\nu' \rangle$. Equation(14) can be solved by using the unperturbed Green's function Eq.(10). The solution is given by

$$C_{nm}(R_\mu, R_\nu) = C_{nm}^{(0)}(R_\mu, R_\nu) + \sum_{\mu''\nu'', \mu'''\nu'''} G_{nm}(\mu\nu, \mu'\nu') V_{nm; \mu'\nu', \mu''\nu''} C_{nm}(R_{\mu''}, R_{\nu''}), \quad (15)$$

where $C_{nm}^{(0)}$ pertains to the uncoupled superposition of two-atom Wannier functions satisfying the equation

$$\sum_{\mu'\nu'} \langle nm; \mu\nu | (H_0 - E) | nm; \mu'\nu' \rangle C_{nm}^{(0)}(R_{\mu'}, R_{\nu'}) = 0. \quad (16)$$

The formation of a quasibound state on resonance gives rise to *quantum translational correlations* (*spatial entanglement*) of the two atoms. This entanglement is manifest by strong deviations of the C_{nm} in Eq.(15) from the uncoupled coefficients $C_{nm}^{(0)}$, at E near the scattering resonance calculated above. It should be possible to obtain multi-atom correlated states, by the interaction of adjacent pairs, using a multi-dimensional extension of Eqs(13)-(16).

To conclude, we have demonstrated the possibility of observing a novel effect: the quasi-binding of cold atoms residing at adjacent sites of an optical lattice, by applying a plane-wave, off-resonant laser that induces an attractive dipole-dipole interaction between these sites. This quasi-binding is a long-range quasi-molecular collision capable of producing translational two-atom or multi-atom states with intriguing quantum correlations.

ACKNOWLEDGMENTS

This work has been supported by the US-Israel Foundation (BSF).

-
- [1] C. Cohen-Tannoudji, Rev.Mod.Phys. **70**, 707 (1998); S.Chu, Rev.Mod.Phys. **70**,685 (1998); W.D. Philips, Rev.Mod.Phys. **70**, 721 (1998).
 - [2] P.S. Jessen and I.H. Deutsch, Adv.At.Mol.Opt.Phys. **37**, 94 (1996); S. Rolston, Physics World, 27 (Oct.,1998); G. Grynberg et al., Phys.Rev.Lett. **70**, 2249 (1993); A. Hemmerich and T.W. Hansch, Phys.Rev.Lett. **70**, 410 (1993);
 - [3] D. Kleppner, Physics Today, 11 (July, 1995); P.D. Lett et al., Annu.Rev.Phys.Chem.**46**, 423 (1995); J. Weiner, Adv.At.Mol.Opt.Phys. **35**, 45 (1995); A. Fioretti et al., Phys.Rev.Lett. **80**, 4402 (1998).
 - [4] T. Thirunamachandran, Molecular Physics **40**, 393 (1980); M.M. Burns et al., Phys.Rev.Lett. **63**, 1233 (1989); M.M. Burns et al., Science **249**, 749 (1990); P. Grossel and D.V. Labeke, Phys.Rev. A **49**, 1177 (1994); P.B. Lerner, Aus.J.Phys. **48**, 849 (1995); P.W. Milonni and A. Smith, Phys.Rev. A **53**, 3484 (1996);
 - [5] D. O'Dell, S. Giovanazzi, G. Kurizki and V.M. Akulin, Phys.Rev.Lett. **84**, 5687 (2000).
 - [6] K. Góral, K. Rzążewski and T. Pfau, Phys. Rev. A **61**, 051601(R) (2000).
 - [7] As shown by S. Giovanazzi, D. O'Dell and G. Kurizki [subm. to Phys. Rev. A (Rapid Commun.), quant-ph/0010045 v3], the rate of atom ejection from the binding potential due to interference between the mutually-detuned laser beams can be comparable to the Rayleigh scattering rate, under conditions similar to those in Fig. 1. Both of these loss mechanisms could significantly reduce the lifetime of the two-atom bound state, as calculated here [Eq. (4)].
 - [8] J. Callaway, Quantum Theory of the Solid State, (Academic,1974); J. Callaway, J.Math.Phys. **5**, 783 (1964).
 - [9] G.F. Koster and J.C. Slater, Phys.Rev. **96**, 1208 (1954).

Figure Captions

FIG.1 (a) The laser-induced dipole-dipole potential for two Cs atoms in their ground states ($6S_{1/2}$) as a function of kr (Eq. (1)). The dipole-dipole potential is induced by a circularly polarized laser field with intensity 5 W/cm^2 , incident parallel (solid line) or perpendicular (dashed line) to the lattice axis, detuned by 1.0 GHz from the atomic transition $6S_{1/2} \rightarrow 6P_{3/2}$ of wavelength $\lambda_L = 852.113 \text{ nm}$ and linewidth $\Gamma/2\pi = 5.21 \text{ MHz}$. The saturation parameter is 0.0485 . (b) One dimensional periodic lattice potential of the form $V_L(X) = V_0 \sin^2 k_L X$ (see text). The lattice generating lasers (intensity 50 W/cm^2) are detuned by 5 GHz from the same transition. For tightly-bound atoms in the lowest band the frequency of the lowest harmonic oscillation is 15.7 kHz .

FIG.2 (a). The variation of the real part of the matrix determinant D as a function of energy for a 1D lattice under the tight binding approximation, using the parameters of Fig.1. Both atoms are in the lowest band, whose energy ranges from 14.906 to 16.499 kHz . A resonance occurs at energy 15.715 kHz with linewidth 1.331 kHz . (b) Same as in (a) for 2D square lattice under the tight binding approximation. The interatomic potential is $V_+ = V_{\parallel} + V_{\perp}$ with field intensity 50 W/cm^2 . Both atoms are in the lowest band whose energy ranges from 29.813 kHz to 32.998 kHz . A resonance occurs at energy 32.504 kHz with linewidth 717 Hz .

Fig.3 (a) Atomic density of states (ρ) and its change ($\delta\rho$) due to laser field induced interatomic interactions in a 1D lattice, for the parameters of Fig. 2a. The density of states under the influence of the field-induced interactions increases rapidly near the resonance energy. (b) Same as in Fig.3 (a), in a 2D lattice for the parameters of Fig. 2(b).





