Entangling many atomic ensembles through laser manipulation

L.-M. Duan 1,2*

¹Institute for quantum information, mc 107-81, California Institute of Technology, Pasadena, CA 91125-8100 ²Laboratory of quantum information, USTC, Hefei 230026, China

We propose an experimentally feasible scheme to generate Greenberger-Horne-Zeilinger (GHZ) type of maximal entanglement between many atomic ensembles based on laser manipulation and single-photon detection. The scheme, with inherent fault tolerance to the dominant noise and efficient scaling of the efficiency with the number of ensembles, allows to maximally entangle many atomic ensemble within the reach of current technology. Such a maximum entanglement of many ensembles has wide applications in demonstration of quantum nonlocality, high-precision spectroscopy, and quantum information processing.

PACS numbers: 03.65.Ud, 03.67.-a, 42.50.Gy, 42.50.-p

Quantum entanglement links two or more distant subsystems in a profound quantum mechanical way. Such a link has found wide applications in demonstration of quantum nonlocality [1,2], high-precision spectroscopy [3], and quantum information processing including computation, communication and cryptography [4,5]. There are great experimental efforts recently to get more and more subsystems entangled [6–9], since with more subsystems entangled, quantum nonlocality becomes more striking [1,2], and the entanglement is more useful for various applications [3–5]. In most of the experimental efforts, the subsystems are take as single-particles, and up to now three to four atoms or photons have been entangled with a linear ion-trap [8], with a spontaneous parametric down converter [6,9], or with a high-Q cavity [7]. There are also proposals to entangle indistinguishable atoms in Bose-Einstein condensates [10], or to weakly entangle two macroscopic atomic ensembles [11], and the latter has been demonstrated in a recent exciting experiment [12].

In all the experimental efforts, it is hard to continuously increase the number of the entangled subsystems due to the fast exponential decrease of the preparation efficiency [6,7,9] or due to noise and imperfections in the setup [7,8]. Here, we propose a scheme to generate GHZ type of maximal entanglement between many atomic ensembles with the following features: firstly, the scheme has built-in fault-tolerance and is robust to realistic noise and imperfections. As a result, the physical requirements of the scheme are moderate and well fit the experimental technique. Secondly, the preparation efficiency of the GHZ entanglement only decreases with the number of ensembles by a slow polynomial law. Such an efficient scaling makes it possible to maximally entangle many (such as tens of) ensembles with the current technology. Our scheme is based on Raman type laser manipulation of the ensembles and single-photon detection which postselects the desired entangled state in a probabilistic fashion. In contrast to the belief that entangling schemes based on postselections will necessarily suffer from the fast exponential degradation of the efficiency, we design a scheme which circumvents this problem by making use of quantum memory available in atomic internal levels.

The basic element of our system is an ensemble of many identical alkali atoms, whose experimental realization can be either a room-temperature atomic gas [12,13] or a sample of cold trapped atoms [14,15]. The relevant level structure of the atom is shown by Fig. 1. From the three levels $\left|g\right\rangle,\left|h\right\rangle,\left|v\right\rangle$, we can define two collective atomic operators $s = (1/\sqrt{N_a}) \sum_{i=1}^{N_a} |g\rangle_i \langle s|$ with s = h, v, where $N_a \gg 1$ is the total atom number. The atoms are initially prepared through optical pumping to the ground state $|q\rangle$, which is effectively a vacuum state $|vac\rangle$ of the operators h, v. The h, v behave like independent bosonic mode operators as long as most of the atoms remain in the state $|g\rangle$. A basis of the "polarization" qubit (in analogy to the language for photons) can be defined from the states $|H\rangle = h^{\dagger} |\text{vac}\rangle$ and $|V\rangle = v^{\dagger} |\text{vac}\rangle$, which have an experimentally demonstrated long coherence time [12-15]. Single-bit rotations in this basis can be done with high precision by shining Raman pulses or radio-frequency pulses on all the atoms. The excitations in the mode h can be transferred to optical excitations [16] and then detected by single-photon detectors. Such a transfer has a high efficiency even for a free-space ensemble due to the collectively enhanced coherent interaction as has been demonstrated both in theory [16] and in experiments [13,14].

The first step for generation of many-party entanglement is to share an excitation between the modes h_i, h_j in two distant ensembles i, j. This can be readily done through a scheme in the recent quantum repeater proposal [17], where one prepares the state $\left(h_i^{\dagger} + e^{i\phi_{ij}}h_j^{\dagger}\right)/\sqrt{2}|\text{vac}\rangle$ [18], with $\phi_{ij} = \phi_j - \phi_i$, an unknown phase difference fixed by the optical channel connecting the i, j ensembles. This state, after a single-bit rotation, can be transferred to the useful form

$$|\Psi_{i,j}\rangle = \left(h_i^{\dagger} + e^{i\phi_{i,j}}v_j^{\dagger}\right)/\sqrt{2} |\text{vac}\rangle.$$
 (1)

The basic idea of the preparation scheme in Ref. [17] is as follows: one excites the ensembles i and j respectively through a short weak Raman pulse applied to the transi-

^{*}Email: lmduan@caltech.edu

tion $|g\rangle \rightarrow |e\rangle$ so that the forward scattered Stokes light from the transition $|e\rangle \rightarrow |h\rangle$ has a mean photon number much smaller than 1. The forward-scattered Stokes lights from the two ensembles are then interfered at a beam splitter and further detected by two single photon detectors. If we successfully get a detector click, we do not know from which ensemble this registered photon comes from, and due to this indistinguishability, the accompany collective atomic excitation should be distributed over the ensembles i and j with an equal probability amplitude, and we get the state described before. This preparation scheme has the following two features: first, the preparation only succeeds with a controllable small probability p_0 for each Raman driving pulse, and needs to be repeated in average $1/p_0$ times for the final successful state generation, with the total preparation time $t_0 \sim 1/(p_0 f_p)$, where f_p is the repetition frequency of the Raman pulses. Second, the scheme, with inherent resilience to noise, is well based on the current technology of laser manipulation. We can safely use it as our first step, to generate the state (1) with a fidelity $F = 1 - p_0$ very close to the unity by controlling the probability p_0 . For instance, with a typical repetition frequency $f_p = 10$ MHz, one may prepare the state (1) with a fidelity $F = 1 - p_0 \approx 99\%$ in a time $t_0 \sim 10\mu$ s.

Based on the preparation of the state (1), now we show how to generate effective many-party entanglement between n such atomic ensembles. We prepare the state (1) between the i and i + 1 ensembles for each i from 1 to n, and get the following state

$$|\Psi\rangle = \left(1/\sqrt{2^n}\right) \prod_{i=1}^n \left(h_i^{\dagger} + e^{i\phi_{i,i+1}} v_{i+1}^{\dagger}\right) |\operatorname{vac}\rangle, \qquad (2)$$

where for convenience we have assumed the notation $n + 1 \equiv 1$ for the subscripts, and have used the same symbol $|\text{vac}\rangle$ to denote the vacuum of the whole *n* ensembles. In the expansion of the state (1), there are only two components which have one excitation on each ensemble. This component state is given by

$$|\Psi_{\text{eff}}\rangle = \left(1/\sqrt{2}\right) \left(\prod_{i=1}^{n} h_{i}^{\dagger} + e^{i\phi_{t}} \prod_{i=1}^{n} v_{i}^{\dagger}\right) |\text{vac}\rangle \qquad (3)$$

with $\phi_t = \sum_{i=1}^n \phi_{i,i+1}$, which is exactly the *n*-party GHZ type maximally entangled state in the polarization basis. Note that for any practical application of the GHZ entanglement [1–4], the state preparation should be succeeded by a measurement of the polarization of the excitation on each ensemble, which can be done for our system by combining single-bit rotations, such as Hadamard transformations, with the number detection of the mode h_i through single-photon detectors. If in this measurement we only keep the results for which an excitation appears on each ensemble (i.e., postselect the case when the detector on each side registers a click), the states (2) and (3) become effectively equivalent since the other components in the state (2) have no contributions to the measurement. Through this postselection technique, we can simply prepare the state (2), which, whenever we put it into applications, yields effectively the GHZ entanglement described by the effective state (3). Here and in the following, we call a component of the full state as the effective state if only this component has contributions to the application measurements (which are the measurements required for the detection or application of the generated state). The effective state is the state postselected by the application measurements.

For applications of the GHZ entanglement, we need also to know the phase ϕ_t in the effective state (3), which is fixed by the whole setup and in principle can be measured. However, a better way is to directly cancel this unknown phase ϕ_t with the following method. Assume that we have an even number n of the ensembles. The pair of ensembles i and i' = n+2-i are put in near proximity so that the ensembles i, i+1 and i', (i+1)' can be connected through the same optical channel, which fixes the phases to satisfy the relation $\phi_{i,i+1} = \phi_{i',(i+1)'} = -\phi_{(i+1)',i'}$ [19]. With this relation, the accumulated phase ϕ_t is exactly canceled to zero.

The above preparation scheme of the effective GHZ entanglement is robust to realistic noise and imperfections. The dominant noise in this system is the photon detector inefficiency, the transferring inefficiency (induced by the spontaneous emission loss) of the excitation from the atomic mode h_i to the optical mode, and the small decay of the atomic excitation in each ensemble. All the above noise is well described by loss of excitations with a overall loss probability denoted by η . Note that by including the detector inefficiency, we have automatically taken into account that the single-photon detectors cannot perfectly distinguish between single and two photons. It is easy to see that loss of excitations only has influence on the success probability to register an excitation from each ensemble. Whenever the excitation is registered, its polarization is still perfectly entangled as shown by the effective state (3).

Now we consider the efficiency of this scheme, which can be described by the total time needed to successfully register the effective GHZ entanglement. The preparation of the factor state (1) is probabilistic, however, due to the available quantum memory provided by the metastable atomic modes h, v, the preparation time t_1 of the state (2) is at most nt_0 if its factor states are prepared one after the other, and can be reduced to $t_1 \sim t_0$ (in the case of $n < 1/p_0$ if its factor states are prepared independently at the same time. In contrast to this, in the case of no quantum memory, one would need about $1/p_0^n$ repeats of the Raman pulses for a successful preparation of the state (2), and a total time $t_0/p_0^{n-1} \gg t_0$. After preparation of the state (2), the projection efficiency (success probability) from the state (2) to the effective GHZ state (3) is given by $(1-\eta)^n/2^{n-1}$, where we have assumed the same loss probability η for each ensemble. So the total time for registering the *n*-party GHZ entanglement is $T \sim t_0 2^{n-1} / (1-\eta)^n$, which increases with the number of ensembles exponentially by the factor $2/(1-\eta)$.

Note that this increase has been much slower than the case for spontaneous parametric down conversion where the exponential increasing factor is about 2 orders larger due to the absence of quantum memory [6,9].

We can in fact further improve the scheme to get a much more efficient scaling of the efficiency, with the time T increasing with the party number n only polynomially. The improved scheme is divided into the following three steps:

(i) We start with two pairs of ensembles 1, 2 and 3, 4, prepared in the state $|\Psi_{1,2}\rangle \bigotimes |\Psi_{3,4}\rangle$ with $|\Psi_{i,j}\rangle$ in the form of Eq. (1). We then connect these two disjoint pairs by preparing the state $|\Psi_{2,3}\rangle$. The ensembles 2 and 3 will not be involved any more in the following steps for state preparation, so we can immediately put them into applications by doing the same type of measurements on them as if we had generated *n*-party GHZ entanglement. In these measurements, if one excitation is registered from each ensemble 2 and 3, we succeed and will go on with the next step. Otherwise, we simply repeat the above process until we succeed. Upon success, only the component $|\Psi_{1-4}\rangle$ of the state $|\Psi_{1,2}\rangle \bigotimes |\Psi_{3,4}\rangle \bigotimes |\Psi_{2,3}\rangle$ has contributions to the measurement with

$$|\Psi_{1-4}\rangle = \left(1/\sqrt{2}\right) \left(h_1^{\dagger} h_2^{\dagger} h_3^{\dagger} + v_2^{\dagger} v_3^{\dagger} v_4^{\dagger}\right) |\text{vac}\rangle, \qquad (4)$$

where for simplicity we have neglected the phase $\phi_{i,i+1}$ since they will finally cancel each other with the method described before. If loss of excitations with a loss probability η is taken into account for detections on the ensembles 2, 3, a registered click might result from two excitations, and in this case there will be no excitation in the ensembles 1 and 4. So with the loss, upon success of step (i), instead of $|\Psi_{1-4}\rangle$ the effective state of the ensembles 1-4 is actually described by

$$\rho_{1-4} = (|\Psi_{1-4}\rangle \langle \Psi_{1-4}| + c_1 \rho_{\text{vac}}) / (1 + c_1)$$
(5)

with the vacuum coefficient $c_1 = 2\eta$, where ρ_{vac} stands for the vacuum component with no excitation in the undetected ensembles 1 and 4. The probability of a successful detection on both of the ensembles 2 and 3 is given by $p_1 = (1 - \eta)^2 (1 + 2\eta) / 4$, which means that we need to repeat the process in average $1/p_1$ times for the final success of step (i).

(ii) In step (ii) we further extend the number of entangled ensembles in the effective state (5). Assume that we have applied the method of step (i) in parallel to the two disjoint sets of ensembles 1-4 and 5-8, with their effective states (each in the form of Eq. (5)) denoted by ρ_{1-4} and ρ_{5-8} , respectively. We connect these two sets by first preparing the state $|\Psi_{4,5}\rangle$ (in the form of Eq. (1)) and then putting the ensembles 4, 5 into application measurements as described in step (i). Upon success of these measurements with one excitation registered from each ensemble, the postselected state of the ensembles 1-8 is effectively described by ρ_{1-8} which is similar to Eq. (5), but with an increased vacuum coefficient and with $|\Psi_{1-4}\rangle$ replaced by $|\Psi_{1\cdot k}\rangle = (1/\sqrt{2}) \left(\prod_{i=1}^{k-1} h_i^{\dagger} + \prod_{i=2}^{k} v_i^{\dagger}\right) |\text{vac}\rangle$, (k = 8). Whenever the measurement fails, we repeat the whole state preparation from step (i). The above connection process can be continued with the number n of effectively entangled ensembles doubled for each time of connection. After i times connection, we have $n = 2^{i+1}$. The success probability and the new vacuum coefficient of the ith connection are denoted respectively by p_i and c_i , which satisfy the following recursion relations with the previous vacuum coefficient c_{i-1} through $p_i = (1 - \eta)^2 (1 + 2\eta + 2c_{i-1}) / \left[4 (1 + c_{i-1})^2\right]$, and $c_i = 2c_{i-1} + 2\eta$. From these recursion relations, we have $c_i = 2\eta \left(2^i - 1\right)$, which, after substituted into p_i , yields an explicit expression for the repetition number $1/p_i$ of the ith connection.

(iii) After a desired number $n = 2^{i+1}$ of the ensembles have been entangled in the effective state ρ_{1-n} , we close the loop in the last step by first preparing the state $|\Psi_{n,1}\rangle$ (in the form of Eq. (1)) and then putting the last two ensembles n, 1 into application measurements. As usual, we keep the results only when one excitation appears from each detected ensemble, and this automatically eliminates contributions from the vacuum component in the state ρ_{1-n} . So the effective state of the whole set of ensembles postselected by all the application measurements is still described by the exact GHZ state (3), and the application measurement results should reveal perfect GHZ entanglement between the n ensembles in the polarization degree of freedom. The application measurements on the ensembles n, 1 in the last step succeeds with a probability $p_l = (1 - \eta)^2 / [2(1 + c_i)]$, so the whole process needs to be repeated in average $1/p_l$ times.

Now we calculate in this improved scheme how much time is needed in total for a successful detection of the n-party GHZ entanglement. This time is given by $T_{\rm imp} = t_0 / \left| p_l p_1 \prod_{j=2}^i p_j \right|$, with t_0 , the preparation time of the state (1). We consider two limiting cases. In the first case with a negligible loss probability η for each ensemble, we have $p_l = 2p_j = 1/2$ and $T_{\rm imp} = 2^{2i+1}t_0 = n^2 t_0/2$, which increases with the number n of entangled ensembles by the slow quadratic law. In the second case with a considerably large loss probability η , the total time $T_{\rm imp}$ is approximated by $T_{\rm imp} \sim t_0 \left[2\eta n / (1-\eta)^2 \right] (n/2)^{\log_2 \left[2\eta \sqrt{n} / (1-\eta)^2 \right]}$, which increases with n faster, but still polynomially (or, more accurately, sub-exponentially). The basic reason for the improvement from the exponential scaling to the much slower polynomial scaling is due to that we have divided the whole preparation process into many small steps, checking in each step whether the preparation is successful, and repeating this small step instead of the whole process if it fails.

Finally, we briefly discuss the practical implication of this proposal. With the improved scheme, for example, we can generate high-fidelity GHZ entanglement over n = 16 ensembles in a time $T_{\rm imp} \sim 50$ ms with a notable loss $\eta \approx 1/3$ and a typical choice $t_0 \sim 10 \mu s$. With such a short preparation time $T_{\rm imp}$, the noise that we have not included, such as the non-stationary phase drift induced by the pumping lase or by the optical channel, is negligible. As long as the number n of the ensembles is not huge, we can also safely neglect the single-bit rotation error (below 10^{-4} with the use of accurate polarization techniques for Zeeman sublevels [20] and the dark count probability (about 10^{-5} in a typical detection time window $0.1\mu s$) of single-photon detectors. Due to the efficient scaling of this scheme, one can use it to steadily increase the number of entangled ensembles, and it seems reasonable to generate GHZ entanglement over tens of ensembles with the current technology. Such an extraordinary possibility opens up prospects for many exciting experiments and applications.

Acknowledgments This work was supported in part by the Caltech MURI Center for Quantum Networks under ARO Grant No. DAAD19-00-1-0374, by the National Science Foundation under Grant No. EIA-0086038, and also by the Chinese Science Foundation and Chinese Academy of Sciences.

D. M. Greenberger, M. A. Horne, A. Shimony, and A. Zeilinger, Am. J. Phys. 58, 1131 (1990).

- [2] N. D. Mermin, Phys. Rev. Lett. 65, 1838 (1990).
- [3] J. Bollinger, W. M. Itano, D. Wineland, D. Heinzen, *Phys. Rev. A* 54, 4649 (1996).
- [4] D. Gottesman, I. L. Chuang, Nature 402, 390 (1999).
- [5] M. A. Nielsen, I. L. Chuang, *Quantum Computation and Quantum Information*, (Cambridge University Press, UK, 2000).
- [6] D. Bouwmeester et al., Phys. Rev. Lett. 82, 1345 (1999).
- [7] A. Rauschenbeutel et al., Science 288, 2024 (2000).
- [8] C. A. Sackett et al., Nature 404, 256 (2000).
- [9] J.-W. Pan et al., Phys. Rev. Lett. 86, 4435 (2001).
- [10] A. Sorensen, L. M. Duan, J. I. Cirac, P. Zoller, *Nature* 409, 63 (2001).
- [11] L. M. Duan, J. I. Cirac, P., Zoller, E. S. Polzik, *Phys. Rev. Lett.* 85, 5643 (2000).
- [12] B. Julsgaard, A. Kozhekin, E. S. Polzik, *Nature* **413**, 400 (2001).
- [13] Phillips, D. F. *et al.* Storage of light in atomic vapor. *Phys. Rev. Lett.* **86**, 783-786 (2001).
- [14] C. Liu, Z. Dutton, C. H. Behroozi, L. V. Hau, Nature 409, 490 (2001).
- [15] J.-F. Roch, et al. Phys. Rev. Lett. 78, 634 (1997).
- [16] M. Fleischhauer, M. D. Lukin, *Phys. Rev. Lett.* 84, 5094 (2000).
- [17] L. M. Duan, M. D. Lukin, J. I. Cirac, P. Zoller, Nature 414, 413 (2001).
- [18] Ref. [17] shows that the generated state will be mixed with a small vacuum component if we take into account

the detector dark counts. However, this vacuum component is typically very small since the normal dark count rate (100 Hz) is much smaller than the repetition frequency (10 MHz, for instance) of the Raman pulses, and in addition, this component will be finally automatically eliminated in our scheme since its effect is the same as the effect of detector inefficiency in the application measurements which we will analyze below. So, for simplicity, we write a pure state here by neglecting this small component.

- [19] Remember the notation $n + 1 \equiv 1$. The index *i* and i' = n + 2 i actually denote the same ensemble for i = 1 and n/2 + 1. So as a special case of the relation $\phi_{i,i+1} = -\phi_{(i+1)',i'}$, we have $\phi_{1,2} = -\phi_{n,1}$ and $\phi_{n/2,n/2+1} = -\phi_{n/2+1,n/2+2}$.
- [20] D. Budker, V. Yashuk, M. Zolotorev, *Phys. Rev. Lett.* 81, 5788-5791 (1998).



FIG. 1. The relevant atomic level structure with $|g\rangle$, the ground state, $|e\rangle$, the excited state, and $|h\rangle$, $|v\rangle$ the two metastable states (e.g., Zeeman or hyperfine sublevels) for storing a qubit of information. The three levels $|g\rangle$, $|e\rangle$, and $|h\rangle$ can be coupled through a Raman process which is useful for measurement of the collective atomic excitation in the state $|h\rangle$ [16] and for generating preliminary entanglement between two ensembles [17].