Linear Optics Quantum Communication over Long Distances

Zeng-Bing Chen,^{1*} Huai-Xin Lu,¹ and Yong-De Zhang^{2,1}

¹Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230027, People's Republic of

China

²CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, People's Republic of China

(November 20, 2018)

We propose a feasible scheme for teleporting an arbitrary polarization state or entanglement of photons by requiring only single-photon (SP) sources, simple linear optical elements and SP quantum non-demolition measurements. An unknown SP polarization state can be faithfully teleported either to a duplicate polarization state or to an entangled state. Our proposal can be used to implement long-distance quantum communication in a simple way. The scheme is within the reach of current technology and significantly simplifies the realistc implementation of long-distance high-fidelity quantum communication with photon qubits.

PACS numbers: 03.67.Hk, 03.67.-a, 42.50.-p

Quantum mechanics offers us the surprising capabilities of manipulating quantum information for computational and communicational tasks in a nonclassical way [1]. So far, many quantum information protocols have been proposed to realize the capabilities. Among these protocols, much attention has been paid on quantum communication (QC), e.g., quantum cryptography [2] and quantum teleportation [3–6]. Quantum teleportation is a process that transmits an unknown quantum state from a sender (Alice) to a receiver (Bob) via a quantum channel with the help of some classical information. It may also play a constructive role in quantum computation [7,8]. As is widely believed, many quantum information protocols [1], including QC, heavily rely on the pre-existing Einstein-Podolsky-Rosen (EPR) entanglement [9] as a necessary "resource". Meanwhile, quantum entanglement is also a test ground of various fundamental problems in quantum mechanics. Along this line of thought, a large number of physical systems (e.g., photons [10,11]) has been proposed to generate entangled states, a notoriously difficult task.

QC ultimately aims to transmit quantum states (e.g., the polarization states of photons) or create quantum correlations between remote locations, a long-sought goal in the context of quantum information. For transmitting an unknown quantum state with fidelity 1, the quantum channel is a maximally entangled state [3]. Thus the establishment of high-quality entangled states between distant nodes is essential for large-scale QC. On the one hand, photons still remain the best long-distance carriers of quantum information. On the other hand, the unavoidable noise of the photonic channel degrades the entanglement established between two distant parties. The degradation of quantum entanglement (as well as the communication fidelity) is exponential with respect to the channel length [12] and remains one of the main obstacles for robust long-distance QC. To face this difficulty, various entanglement purification protocol have been proposed [13,14]. However, since entanglement purification, though interesting in its own right, requires an exponentially large number of partially entangled states to obtain one desired entangled state, the quantum repeater protocol is necessary to overcome the exponential decay of the entanglement [14].

Motivated by several linear optics implementations of quantum information processing (quantum computation [8,15], entanglement purification [14] and generation of polarization-entangled photons [11]), this paper demonstrates a dramatically different approach for QC and quantum network over long distances by thoroughly exploiting linear optics, as envisioned in Refs. [8,11]. We first propose a scheme for teleporting an arbitrary polarization state or entanglement of photons. It requires only the single-photon (SP) sources, simple linear optical elements [e.g., the polarizing beam splitters (PBS) and wave plates] and quantum non-demolition measurements (QNDM) with SP resolution [16,17]. An unknown SP polarization state can be teleported either to another photon with the identical polarization state or to an entangled state; both of the possibilities occur with the probability of 50%. We then show that our scheme, together with the concept of quantum repeaters [12], could be used to implement long-distance QC in a simple way. For teleporting with certainty, one requires the complete Bellstate measurement (BSM). Since the first teleportation experiment, which realized the identification of only one, among four, Bell state [4], several feasible schemes have been proposed to achieve the full BSM [18–20]. Combining these BSM schemes and the recent achievement of nearly ideal SP source [21], our long-distance QC scheme is within the reach of current technology.

Very recently, Duan *et al.* [22] proposed a different scheme of long-distance QC, which involves laser manipulation of atomic ensembles, beam splitters and SP detectors. While their proposal represents a novel way of creating long-distance quantum entanglement between two atomic ensembles, we are aiming to establish remote quantum entanglement between photons by linear optics.

In the original proposal of quantum teleportation [3], Alice possesses a polarization state of photon 1

$$\left|\psi\right\rangle_{1} = \alpha \left|H\right\rangle_{1} + \beta \left|V\right\rangle_{1}, \qquad (1)$$

where H(V) denotes the horizontal (vertical) linear polarization; α and β are two arbitrary complex probability amplitudes satisfying $|\alpha|^2 + |\beta|^2 = 1$. The state $|\psi\rangle_1$, completely unknown to Alice, is to be transmitted by Alice to Bob via one of the following quantum channels (the Bell states)

$$\begin{split} \left| \Psi^{\pm} \right\rangle_{23} &= \frac{1}{\sqrt{2}} (\left| H \right\rangle_2 \left| V \right\rangle_3 \pm \left| V \right\rangle_2 \left| H \right\rangle_3), \\ \left| \Phi^{\pm} \right\rangle_{23} &= \frac{1}{\sqrt{2}} (\left| H \right\rangle_2 \left| H \right\rangle_3 \pm \left| V \right\rangle_2 \left| V \right\rangle_3), \end{split}$$
(2)

which can be generated in the process of spontaneous parametric down-conversion in a nonlinear optical crystal [4,10]. In the four Bell states $|\Psi^{\pm}\rangle_{23}$ and $|\Phi^{\pm}\rangle_{23}$, the subscripts label the photon 2 (possessed by Alice) and the photon 3 (possessed by Bob). Photons 2 and 3 (the entangled qubits) are maximally entangled when they are in these Bell states. In the original quantum teleportation protocol, one requires an existing Bell state as a prerequisite resource. Currently, the mostly used reliable source of the polarization-entangled photons is parametric down-conversion in a nonlinear optical crystal [10]. However, the process is spontaneous (or random) and suffers from the low yield. This limits the utility of such an entangled photon source, in particular in long-distance QC, where the resource overheads are rather large.

Here we suggest a different quantum teleportation protocol by exploiting only simple linear optical elements and SP sources, accompanied with appropriate quantum measurements. Consider three independent SP sources emitting three photons: the SP source-1 emits photon 1 with quantum state $|\psi\rangle_1$, while the remaining two SP sources emit photons 2 and 3, whose quantum states are

$$|\psi\rangle_j = \frac{1}{\sqrt{2}} (|H\rangle_j + |V\rangle_j). \quad (j = 2, 3)$$
(3)

These SP states in Eqs. (1) and (3) can be easily generated with high precision since only trivial single-qubit operations are involved. The current technology has realized deterministic and controllable SP sources with high repetition rate and without the requirement of a cryogenic temperature [21], making them very suitable for the long-distance QC task.

Now photons 2 and 3 are superimposed onto the PBS₂₃ simultaneously, resulting in two output 2' and 3'. Then the photon(s) in the output 2' and photon 1 are incident on the PBS_{12'}, with the corresponding output 1' and 2". Since the PBS reflects V-photons and transmits H-photons, the whole state $|\psi\rangle_1 |\psi\rangle_2 |\psi\rangle_3$ of the three photons after the action of the two PBS is transformed into

$$\begin{split} |\psi\rangle_{1'2''3'} &= \frac{1}{2} [\alpha |H\rangle_{1'} |H\rangle_{2''} |H\rangle_{3'} + \beta |V\rangle_{1'} |V\rangle_{2''} |V\rangle_{3'} \\ &+ \alpha |0\rangle_{1'} |HV\rangle_{2''} |V\rangle_{3'} + \beta |HV\rangle_{1'} |0\rangle_{2''} |H\rangle_{3'} \end{split}$$

$$+ (\alpha |V\rangle_{1'} |HV\rangle_{2''} + \beta |HV\rangle_{1'} |V\rangle_{2''}) |0\rangle_{3'} + (\alpha |0\rangle_{1'} |H\rangle_{2''} + \beta |V\rangle_{1'} |0\rangle_{2''}) |HV\rangle_{3'}],$$
(4)

where $|0\rangle$ and $|HV\rangle$ denote the zero-photon and twophoton (one with *H*-polarization and another with *V*polarization) states, respectively. Introduce two new Bell states [18]

$$\left|\Xi^{\pm}\right\rangle = \frac{1}{\sqrt{2}}(\left|0\right\rangle \left|HV\right\rangle \pm \left|HV\right\rangle \left|0\right\rangle),\tag{5}$$

which are also maximally entangled. Then using Eqs. (2) and (5) enables us to rewrite the first two lines of (4) as

$$\begin{split} |\psi'\rangle_{1'2''3'} &= \frac{1}{2} [|\Phi^{+}\rangle_{1'2''} \left(\alpha |V\rangle_{3'} + \beta |H\rangle_{3'}\right) \\ &+ |\Phi^{-}\rangle_{1'2''} \left(\alpha |V\rangle_{3'} - \beta |H\rangle_{3'}\right) \\ &+ |\Xi^{+}\rangle_{1'2''} \left(\alpha |H\rangle_{3'} + \beta |V\rangle_{3'}\right) \\ &+ |\Xi^{-}\rangle_{1'2''} \left(\alpha |H\rangle_{3'} - \beta |V\rangle_{3'}\right)]. \end{split}$$
(6)

Now Bob, who possesses the output photon(s) 3', needs to perform a QNDM of the photon number at his side. His detection results will be informed to Alice, who possesses the output photons 1' and 2'', via a classical channel. If the detected photon number is zero (two), then the form of (4) and the standard theory of quantum measurement imply that Alice's states are projected onto (i.e., teleported to) an entangled state $\alpha |V\rangle_{1'} |HV\rangle_{2''} +$ $\beta |HV\rangle_{1'} |V\rangle_{2''} (\alpha |0\rangle_{1'} |H\rangle_{2''} + \beta |V\rangle_{1'} |0\rangle_{2''})$, whose entanglement is determined by the input state $|\psi\rangle_1$. In this case, Alice obtains an entangled state conditioned on Bob's measurement results. However, if the photon number detected by Bob is one, then the whole state $|\psi\rangle_{1'2''3'}$ will be projected onto $|\psi'\rangle_{1'2''3'}.$ In this case, Alice can teleport the unknown input state $|\psi\rangle_1$ to Bob by performing the BSM that collapses her states onto one of the four Bell states $|\Phi^{\pm}\rangle_{1'2''}$ and $|\Xi^{\pm}\rangle_{1'2''}$, instead of the four Bell states in Eq. (2) in the original proposal [3]. The present BSM can then be accomplished by exploiting two complete BSM schemes proposed recently [18,19].

For convenience, we call the whole setups (the PBS, the SP sources as well as the required measurement setups) that teleport an unknown input quantum state a "quantum teleporter". The quantum teleporter can faithfully transmit an unknown SP polarization state either to another photon with the identical polarization state or to an entangled state; both of the possibilities occur with the probability of 50%. If the latter is discarded, the success probability of the scheme is reduced to 50%. Since the perfection of the simple linear optical elements is extremely high, the quantum teleporters in our scheme are robust and can transmit the polarization states of photons with high fidelity.

Entanglement swapping is in fact teleportation of entanglement [3,5,6]. The above scheme can also be used in entanglement swapping. In this case we consider two EPR pairs $|\Psi^{-}\rangle_{12}$ and $|\Psi^{-}\rangle_{34}$; photons 1 and 3 are distributed to Alice. Alice superimposes her photons onto a PBS (with the corresponding output 1' and 3'). Afterward, the initial state $|\Psi^{-}\rangle_{12} |\Psi^{-}\rangle_{34}$ becomes

$$\begin{split} |\psi\rangle_{1'23'4} &= \frac{1}{2} [|\Psi^{+}\rangle_{1'3'} |\Phi^{+}\rangle_{24} - |\Psi^{-}\rangle_{1'3'} |\Phi^{-}\rangle_{24} \\ &- |\Xi^{+}\rangle_{1'3'} |\Psi^{+}\rangle_{24} - |\Xi^{-}\rangle_{1'3'} |\Psi^{-}\rangle_{24}]. \end{split}$$
(7)

Then Alice performs the BSM. At the end of the entanglement swapping, conditioned on Alice's result photons 2 and 4 are entangled in one of the four Bell states $|\Psi^{\pm}\rangle_{24}$ and $|\Phi^{\pm}\rangle_{24}$, though they never interact in any way [5,6]. In the above entanglement swapping protocol, the two EPR pairs can be created by the SP sources, simple linear optical elements and the QNDM with SP resolution [11]. The whole setups that swap the quantum entanglement are called an "entanglement swapper".

As in all realistic teleportation experiments [4], the BSM relies on the interference of two independently created photons. One thus has to guarantee good spatial and temporal overlap at the PBS and above all, one has to erase all kinds of the which-way information for the two photons. The procedure is crucial to the validity of Eqs. (4) and (7). Thus, SP sources with good coherence and controllable emission time and direction are highly desirable and in fact, can be realized in quantum dots with integrated cavity structure [21]. The SP coherence may also be enhanced by frequency filters.

While the proposed linear optics implementation of quantum teleportation is interesting in its own right, its usefulness lies in realizing long-distance QC simply by combining the basic idea of quantum repeaters [12] and the linear optics entanglement purification protocol [14]. To achieve a successful QC, one can either send directly the qubit, whose information is encoded in the polarization state of a photon, through a photonic channel (the scheme-I) or create the high-quality entanglement over the channel, i.e., the quantum channel (the scheme-II), which can then be used to transmit quantum information by virtue of the quantum teleportation protocol. The quantum repeater protocol [12] can be implemented correspondingly in two distinct ways. In the present longdistance QC model, one divides the whole QC channel into many nodes; the length between two nearest neighboring nodes is comparable to the channel attenuation length. To faithfully transmit an unknown SP polarization state, a quantum teleporter (together with a quantum memory if needed) is used at each node: The quantum states are transmitted by the quantum teleporters as a relay race. Similarly, entanglement swapping can be extended to a long distance by using many entanglement swappers, each of which is placed at one node. Most importantly, the suggested scheme has obvious advantages in terms of its simplicity and integrability into the linear optics entanglement purification protocol proposed by Pan et al. [14]. The integrability implies that the linear optics realization of long-distance QC may also have the built-in entanglement purification, an important feature that is essential to the long-distance QC protocol due to Duan *et al.* [22]. The crucial point here is that our scheme of quantum teleportation eliminates the stringent requirement of entangled photon sources; both the entanglement and its purification are plug-in on demand. Thus we can overcome the problem associated with the exponential decay of the entanglement and communication fidelity in a simple way.

Another basic building block in the long-distance QC is quantum memory. It is essential for locally storing and manipulating quantum information. As proposed in Ref. [23], a quantum network or a QC channel consists of spatially separated nodes, which are connected by photonic channels. When quantum memory located at each node is represented by the internal states of atoms (or ions), one needs to transfer the internal state of an atom at a node to quantum state of an optical mode, and vice versa at the next node. While this proposal is attractive for creating long-distance quantum correlations of the internal states of atoms over a noisy channel, it is preferable in our scheme to store the quantum states of photons within quantum memory directly to establish the polarization entanglement of photons over a long photonic channel. Fortunately, a light pulse can be, with nearly 100% efficiency in principle, stored for a controllable period of time and then released on demand in an optically dense many-atom system [24,25], as some recent experiments demonstrated [26]. In particular, transfer and storage of the polarization state of a photon are also realizable [24]. Storing polarization entanglement of photons over long distances could also be accomplished by a scheme proposed by Lloyd *et al.* [27]. Thus quantum memory suitable for our scheme is feasible already.

Due to dissipation, noise and imperfect measurements, errors always occur in a realistic QC channel. Detailed analyses of the scalability and resource requirements within our scheme are yet to be done. However, we can make the following remarks. In some sense, our scheme is a minimal one, which has a great advantage of tolerating errors and lessening the required overheads. First, only three measurements, two of which are required for the complete BSM [18], are involved in teleporting an unknown SP state, with six possible outcome states to be distinguished. Meanwhile, the complicated experimental task of creating quantum entanglement does not need to be done separately in the present approach. Moreover, the simple linear-optical elements are thoroughly used within our scheme. Their extremely good performance induces only small errors that can be safely ignored. The minimal scheme we proposed is thus essential to a scalable QC with photons.

As in the linear optics quantum computation [8], dominant sources of errors in our linear optics QC come from photon loss, phase errors and detector inefficiency, which are well understood. These errors can be dealt with by using the method described in Ref. [8]. The detector inefficiency only influences the communication efficiency of our scheme, similarly to other linear optics implementations of quantum information processing [8,14]. Under the scheme-I, complicated error detection and correction are necessarily involved [1]. In this respect, it is more advantageous to work under the scheme-II. Within our linear optics realization of the quantum repeaters, the resources of creating high-quality entanglement over a long QC channel might grow polynomially with the channel length as in the original quantum repeater protocol [12].

The QNDM with SP resolution is crucial in our approach. Standard realization of the ONDM of the photon number requires the cross-Kerr nonlinearity [16], which induces the cross-Kerr phase modulation required to implement the complete BSM [18,19]. To obtain good signal to noise ratio, it is desirable to acquire giant cross-Kerr nonlinearity. Though the required nonlinearity could be obtained by ultraslow SP pulses propagating in a coherently prepared atomic gas [25,28], currently an SP-QNDM still remains an experimental challenge. In Ref. [29], the SP-QNDM has indeed been realized by a cavity-QED technique. But such an SP-QNDM is strongly frequency-dependent. Very recently, Kok et al. [17] proposed a probabilistic interferometric SP-QNDM device with linear optics and SP detectors; the device, which can distinguish the zero-photon, single-photon and two-photon evens, is frequency-independent and may be polarization-preserving. Thus it can be naturally integrated into the present QC protocol.

To summarize, we have proposed a linear optics model for long-distance high-fidelity QC. Our proposal eliminates an obstacle of the large-scale QC since the highquality entanglement required for faithful quantum teleportation and entanglement swapping is plug-in on demand. It relies on the beautiful technologies developed for nearly ideal SP sources aiming at the QC tasks, for the QNDM with SP resolution and for quantum memory of light. The proposed scheme simplifies the experimental aspect of realistic long-distance QC and the optical quantum network. The linear optics quantum information processing has several attractive advantages [8]. With several involved technologies working together, our scheme might be realizable in a near future.

We thank Lu-Ming Duan and Jian-Wei Pan for valuable comments. This work was supported by the National Natural Science Foundation of China under Grants No. 10104014, No. 19975043 and No. 10028406 and by the Chinese Academy of Sciences.

- * E-mail address: zbchen@ustc.edu.cn
- The Physics of Quantum Information, edited by D. Bouwmeester, A. Ekert, and A. Zeilinger (Springer-Verlag, Berlin, 2000).
- [2] C.H. Bennett and G. Brassard, in *Proc. IEEE Int. Conf.* on Computers, Systems and Signal Processing (IEEE, New York, 1984); A.K. Ekert, Phys. Rev. Lett. **67**, 661

(1991).

- [3] C.H. Bennett et al., Phys. Rev. Lett. 70, 1895 (1993).
- [4] D. Bouwmeester et al., Nature (London) **390**, 575 (1997).
- [5] M. Żukowski et al., Phys. Rev. Lett. 71, 4287 (1993).
- [6] J.-W. Pan *et al.*, Phys. Rev. Lett. **80**, 3891 (1998).
- [7] D. Gottesman and I.L. Chuang, Nature (London) 402, 390 (1999).
- [8] E. Knill, R. Laflamme, and G.J. Milburn, Nature (London) 409, 46 (2001).
- [9] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. 47, 777 (1935).
- [10] P.G. Kwiat et al., Phys. Rev. Lett. 75, 4337 (1995).
- [11] Z.-B. Chen, J.-W. Pan, and Y.-D. Zhang, e-print quantph/0105100.
- [12] H.-J. Briegel et al., Phys. Rev. Lett. 81, 5932 (1998).
- [13] C.H. Bennett et al., Phys. Rev. Lett. 76, 722 (1996).
- [14] J.-W. Pan *et al.*, Nature (London) **410**, 1067 (2001); T. Yamamoto, M. Koashi, and N. Imoto, Phys. Rev. A **64**, 012304 (2001); Z. Zhao, J.-W. Pan, and M.-S. Zhan, *ibid.* **64**, 014301 (2001).
- [15] M. Koashi, T. Yamamoto, and N. Imoto, Phys. Rev. A 63, 030301 (2001); T.B. Pittman, B.C. Jacobs, and J.D. Franson, *ibid.* 64, 062311 (2001).
- [16] J.C. Howell and J.A. Yeazell, Phys. Rev. A 62, 032311 (2000).
- [17] P. Kok, H. Lee, and J.P. Dowling, e-print quantph/0202046.
- [18] M.G.A. Paris et al., Phys. Lett. A 273, 153 (2000).
- [19] D. Vitali, M. Fortunato, and P. Tombesi, Phys. Rev. Lett. 85, 445 (2000).
- [20] Y.-H. Kim, S.P. Kulik, and Y. Shih, Phys. Rev. Lett. 86, 1370 (2001).
- [21] B. Lounis and W.E. Moerner, Nature (London) 407, 491 (2000); P. Michler *et al.*, Science 290, 2282 (2000); E. Moreau *et al.*, Appl. Phys. Lett. 79, 2865 (2001); Z. Yuan *et al.*, Science 295, 102 (2002).
- [22] L.-M. Duan, M.D. Lukin, J.I. Cirac, and P. Zoller, Nature (London) 414, 413 (2001).
- [23] J.I. Cirac *et al.*, Phys. Rev. Lett. **78**, 3221 (1997); P.
 Zoller, Nature (London) **404**, 340 (2000).
- [24] A. Fleischhauer, S.F. Yelin, and M.D. Lukin, Opt. Commun. 179, 395 (2000); A. Fleischhauer and C. Mewes, e-print quant-ph/0110056, and references therein.
- [25] M.D. Lukin and A. Imamoğlu, Nature (London) 413, 273 (2001).
- [26] D.F. Phillips et al., Phys. Rev. Lett. 86, 783 (2001); C.
 Liu et al., Nature (London) 409, 490 (2001); A. Mair et al., Phys. Rev. A 65, 031802(R) (2002).
- [27] S. Lloyd *et al.*, Phys. Rev. Lett. **87**, 167903 (2001).
- [28] M.D. Lukin and A. Imamoğlu, Phys. Rev. Lett. 84, 1419 (2000).
- [29] G. Nogues et al., Nature (London) 400, 239 (1999).